arXiv:1012.4576v1 [cond-mat.other] 21 Dec 2010

Lectures on Fröhlich Polarons from 3D to 0D – including detailed theoretical derivations

J. T. Devreese*

Theory of Quantum and Complex Systems (TQC), Universiteit Antwerpen,
CDE, Universiteitsplein, 1, B-2610 Antwerpen, Belgium

(Dated: December 22, 2010)

Abstract

Based on a course presented at the International School of Physics Enrico Fermi, CLXI Course,."Polarons in Bulk Materials and Systems with Reduced Dimensionality", Varenna, Italy, 21.6. - 1.7.2005.

In the present course, an overview is presented of the fundamentals of continuum-polaron physics, which provide the basis of the analysis of polaron effects in ionic crystals and polar semiconductors. These Lecture Notes deal with "large", or "continuum", polarons, as described by the Fröhlich Hamiltonian. The emphasis is on the polaron optical absorption, with detailed mathematical derivations.

---

*Electronic address: jozef.devreese@ua.ac.be

Contents

I Single polaron

I. Introduction. The "standard" theories 5

A. The polaron concept 5

B. Intuitive concepts 6

C. The Fröhlich Hamiltonian 10

D. Infinite mass model [“shift”–operators] 11

E. The "standard" theories 12

1. Weak coupling via a perturbation theory 12
2. Weak coupling via a canonical transformation [“shift”-operators] 13
3. Strong coupling via a canonical transformation [“shift”-operators] 15
4. All-coupling theory. The Feynman path integral 24
5. On Monte Carlo calculations of the polaron free energy 26
6. On the contributions of the N-phonon states to the polaron ground state 28

F. Polaron mobility 31

II. Optical Absorption. Weak coupling 33

A. Optical absorption at weak coupling [within the perturbation theory] 33

B. Optical absorption at weak coupling [within the canonical-transformation method [71] (DHL)] 34

III. Optical absorption. Strong coupling 41

IV. Arbitrary coupling 43

A. Impedance function of large polarons: An alternative derivation of FHIP [74] 43

B. Calculation of the memory function (Devreese *et. al.* [48]) 65

C. Discussion of optical absorption of polarons at arbitrary coupling 68

1. Sum rules for the optical conductivity spectra 72

D. Scaling relations 76

1. Derivation of the scaling relations 76

2. Check of the scaling relation for the path integral Monte Carlo result for the polaron free energy 83
Appendix 1. Weak coupling: LLP approach 84
Appendix 2. Expansion in Stieltjes continuous fractions [52] 97

II Many polarons

V. Optical conductivity of an interacting many-polaron gas 101
A. Kubo formula for the optical conductivity of the many-polaron gas 101
B. Force-force correlation function 104
C. Canonical transformation 106
D. Dynamic structure factor 109
1. Calculation of the dynamic structure factor using the retarded Green's functions 113
2. Plasmon-phonon contribution 117
E. Comparison to the infrared spectrum of $\mathrm{Nd}_{2-x}\mathrm{Ce}_x\mathrm{CuO}_{2-y}$ 118
F. Experimental data on the optical absorption in manganites: interpretation in terms of a many-polaron response 119

VI. Interacting polarons in a quantum dot 123
A. The partition function and the free energy of a many-polaron system 123
B. Model system 129
1. Analytical calculation of the model partition function 130
C. Variational functional 134
D. Two-point correlation functions 138
1. The correlation function $\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta)$ 139
E. Many-polaron ground state in a quantum dot: extrapolation to the homogeneous limit and comparison to the results for a polaron gas in bulk [88] 149
F. Optical conductivity 153
1. Selected results: the manifestations of the shell filling in optical conductivity 159

VII. Variational path-integral treatment of a translation invariant N-polaron

system 162

A. The many-polaron system 162

B. Variational principle 163

C. Results 165

VIII. Ripplonic polarons in multielectron bubbles 168

A. Ripplon-phonon modes of a MEB 168

B. Electron-ripplon interaction in the MEB 170

C. Locally flat approximation 171

D. Ripplopolaron in a Wigner lattice: the mean-field approach 171

E. Ripplopolaron Wigner lattice at finite temperature 173

F. Melting of the ripplopolaron Wigner lattice 175

References 179

Part I

Single polaron

I. INTRODUCTION. THE "STANDARD" THEORIES

A. The polaron concept

A charge placed in a polarizable medium is screened. Dielectric theory describes the phenomenon by the induction of a polarization around the charge carrier. The induced polarization will follow the charge carrier when it is moving through the medium. The carrier together with the induced polarization is considered as one entity (see Fig. 1). It was called a *polaron* by L. D. Landau [1]. The physical properties of a polaron differ from those of a band-carrier. A polaron is characterized by its *binding (or self-) energy* E_0, an *effective mass* m^* and by its characteristic *response* to external electric and magnetic fields (e. g. dc mobility and optical absorption coefficient).



FIG. 1: Artist view of a polaron. A conduction electron in an ionic crystal or a polar semiconductor repels the negative ions and attracts the positive ions. A self-induced potential arises, which acts back on the electron and modifies its physical properties. (From [2].)

If the spatial extension of a polaron is large compared to the lattice parameter of the solid, the latter can be treated as a polarizable continuum. This is the case of a *large (Fröhlich)* polaron. When the self-induced polarization caused by an electron or hole becomes of the order of the lattice parameter, a *smll (Holstein)* polaron can arise. As distinct from large polarons, small polarons are governed by short-range interactions.

B. Intuitive concepts

a. The polaron radius. Large polarons vs small polarons Consider the LO phonon field with frequency ω_{LO} interacting with an electron. Denote by $\Delta\nu$ the quadratic mean square deviation of the electron velocity. In the electron-phonon interaction is weak, the electron can travel a distance

$$\Delta x \approx \frac{\Delta\nu}{\omega_{\mathrm{LO}}} \tag{1.1}$$

during a time $\omega_{\mathrm{LO}}^{-1}$,characteristic for the lattice period,because it is the distance within which the electron can be localized using the phonon field as measuring device. From the uncertainty relations it follows

$$\begin{aligned}
\Delta p \Delta x &= \frac{m}{\omega_{\mathrm{LO}}} \left(\Delta\nu\right)^2 \approx \hbar \\
\Delta\nu &\sim \sqrt{\frac{\hbar\omega_{\mathrm{LO}}}{m}}, \\
\Delta x &\sim \sqrt{\frac{\hbar}{m\omega_{\mathrm{LO}}}}. \qquad (1.2)
\end{aligned}$$

At weak coupling Δx is a measure of the polaron radius r_p. To be consistent, the polaron radius r_p must be considerably larger than the lattice parameter a.(this is a criterion of a "large polaron"). Experimental evaluation of the polaron radius leads to the follwing typical values: $r_p \approx 10\mathring{A}$ for alkali halides, $r_p \approx 20\mathring{A}$ for silver halides, $r_p \approx 100\mathring{A}$ for II-VI, II-V semiconductors. The continuum approximation is not satisfied for transition metal oxides (NiO, CaO, MnO), in other oxides (UO_2,NbO_2...). For those solids the "small polaron" concept is used. In some substances (e.g. perovskites) some intermediate region between large and small polarons is realized.

b. The coupling constant [3] Consider the case of *strong electron-phonon interaction* in a polar crystal. The electron of mass m is then localized and can - to a first approximation

- be considered as a static charge distribution within a sphere with radius l_1. The medium is characterized by an average dielectric constant $\bar{\varepsilon}$,which will be defined below.

The potential energy of a sphere of radius $l_1$uniformly charged with the charge e in a vacuum is (see Eq. (8.6) of Ref. [4])

$$U_{vac} = \frac{3}{5}\frac{e^2}{l_1}. \tag{1.3}$$

The potential energy of a uniformly charged sphere in a medium with the high-frequency dielectric constant ε_∞ is

$$U_1 = \frac{3}{5}\frac{e^2}{\varepsilon_\infty l_1}. \tag{1.4}$$

This is the potential energy of the *self-interaction* of the charge e uniformly spread over the sphere of radius l_1 in a medium with the dielectric constant ε_∞. In a medium with an inertial polarization field (due to LO phonons), the potential energy of the uniformly charged sphere is

$$U_2 = \frac{3}{5}\frac{e^2}{\varepsilon_0 l_1}, \tag{1.5}$$

where ε_0 is the static dielectric constant. The polaron effect is then related to the change of the potential energy of the interaction of the charged sphere due to the inertial polarization field. This change is the potential energy U_2 of the uniformly charged sphere in the presence of the inertial polarization field minus the potential energy of the *self-interaction* $U_1$of the charge e uniformly spread over the sphere in a medium without the inertial polarization:

$$U_{pol} \equiv U_2 - U_1 = \frac{3}{5}\frac{e^2}{l_1}\left(\frac{1}{\varepsilon_0} - \frac{1}{\varepsilon_\infty}\right) = -\frac{3}{5}\frac{e^2}{\bar{\varepsilon} l_1}, \tag{1.6}$$

with

$$\frac{1}{\bar{\varepsilon}} = \frac{1}{\varepsilon_\infty} - \frac{1}{\varepsilon_0}.$$

The electron distribution in a sphere may be non-uniform, what may influence the numerical coefficient in Eqs. (1.4) to (1.6). In this connection one can use the estimate [3]

$$U_{pol} \sim -\frac{e^2}{\bar{\varepsilon} l_1}. \tag{1.7}$$

The restriction of the electron in space requires its de Broglie wave length to be of the order l_1,so that its kinetic energy is of the order $4\pi^2\hbar^2/2ml_1^2$.Minimizing the total energy with respect to $l_1$leads to

$$\frac{\partial}{\partial l_1}\left(-\frac{e^2}{l_1\bar{\varepsilon}}+\frac{4\pi^2\hbar^2}{2ml_1^2}\right)=0\Longrightarrow\frac{1}{l_1}=\frac{e^2m}{4\pi^2\hbar^2\bar{\varepsilon}},$$

wherefrom the binding energy is

$$U_1=-\frac{e^4m}{8\pi^2\hbar^2\bar{\varepsilon}^2}. \tag{1.8}$$

For *weak coupling*, one can neglect the kinetic energy of the electron. Taking the polaron radius according to (1.2), $r_p=\sqrt{2\hbar/m\omega_{\mathrm{LO}}}$, the binding energy is

$$U_2=-\frac{e^2}{r_p\bar{\varepsilon}}=-\frac{e^2}{\bar{\varepsilon}}\sqrt{\frac{m\omega_{\mathrm{LO}}}{2\hbar}}. \tag{1.9}$$

We note that

$$\frac{U_1}{\hbar\omega_{\mathrm{LO}}}=-\frac{e^4m}{8\pi^2\hbar^3\bar{\varepsilon}^2\omega_{\mathrm{LO}}}=-\frac{1}{4\pi^2}\left(\frac{U_2}{\hbar\omega_{\mathrm{LO}}}\right)^2. \tag{1.10}$$

Following the conventions of the field theory, the self energy at weak coupling is written as

$$U_2=-\alpha\hbar\omega_{\mathrm{LO}}.$$

Therefore the so-called Fröhlich polaron coupling constant is

$$\begin{aligned}\alpha&=\frac{e^2}{\bar{\varepsilon}}\sqrt{\frac{m}{2\hbar^3\omega_{\mathrm{LO}}}}\\&\equiv\frac{e^2}{\hbar c}\sqrt{\frac{mc^2}{2\hbar\omega_{\mathrm{LO}}}}\frac{1}{\bar{\varepsilon}}.\end{aligned} \tag{1.11}$$

For the average dielectric constant one shows that

$$\frac{1}{\bar{\varepsilon}}=\frac{1}{\varepsilon_\infty}-\frac{1}{\varepsilon_0},$$

where ε_∞ and ε_0 are, respectively, the electronic and the static dielectric constant of the polar crystal. The difference $1/\varepsilon_\infty-1/\varepsilon_0$ arises because the ionic vibrations occur in the infrared spectrum and the electrons in the shells can follow the conduction electron adiabatically.

c. Polaron mobility Here we give a simple derivation leading to the gross features of the mobility behaviuor, especially its temperature dependence. The key idea is that the mobility will change because the number of phonons in the lattice, with which the polaron interacts, is changing with temperature.

The phonon density is given by

$$n = \frac{1}{e^{\frac{\hbar\omega_{\mathrm{LO}}}{kT}} - 1}.$$

The mobility for large polaron is proportional to the inverse of the number of phonons:

$$\mu \approx \frac{1}{n} = e^{\frac{\hbar\omega_{\mathrm{LO}}}{kT}} - 1$$

and for low temperatures $kT \ll \hbar\omega_{\mathrm{LO}}$

$$\mu \approx e^{\frac{\hbar\omega_{\mathrm{LO}}}{kT}}. \tag{1.12}$$

The mobility of continuum poarons decreases with increasing temperature following an exponential law. The slope of the straight line in $\ln\mu$ vs $1/T$ is characterized by the LO phonon frequesncy. Systematic study performed, in particular, by Fröhlich and Kadanoff, gives

$$\mu = \frac{e}{2m\omega_{\mathrm{LO}}} e^{\frac{\hbar\omega_{\mathrm{LO}}}{kT}}. \tag{1.13}$$

The small polaron will jump from ion to ion under the influence of optical phonons. The lerger the numver of phonons, the lerger the mobility. The behaviuor of the small polaron is the opposite of that of the large polaron. One expects:

$$\mu \approx n = \frac{1}{e^{\frac{\hbar\omega_{\mathrm{LO}}}{kT}} - 1}.$$

For low temperatures $kT \ll \hbar\omega_{\mathrm{LO}}$ one has:

$$\mu \approx e^{-\frac{\hbar\omega_{\mathrm{LO}}}{kT}} : \tag{1.14}$$

the mobility of small polaron is thermally activated. Systematic analysis within the small-polaron theory shows that

$$\mu \approx e^{-\gamma \frac{\hbar\omega_{\rm LO}}{kT}} : \tag{1.15}$$

with $\gamma \sim 5$.

C. The Fröhlich Hamiltonian

Fröhlich proposed a model Hamiltonian for the "large" polaron through which its dynamics is treated quantum mechanically ("Fröhlich Hamiltonian"). The polarization, carried by the longitudinal optical (LO) phonons, is represented by a set of quantum oscillators with frequency $\omega_{\rm LO}$, the long-wavelength LO-phonon frequency, and the interaction between the charge and the polarization field is linear in the field [3]:

$$H = \frac{\mathbf{p}^2}{2m_b} + \sum_{\mathbf{k}} \hbar\omega_{\rm LO} a_{\mathbf{k}}^{+} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a_{\mathbf{k}}^{\dagger} e^{-i\mathbf{k}\cdot\mathbf{r}}), \tag{1.16}$$

where $\mathbf{r}$ is the position coordinate operator of the electron with band mass m_b, $\mathbf{p}$ is its canonically conjugate momentum operator; $a_{\mathbf{k}}^{\dagger}$ and $a_{\mathbf{k}}$ are the creation (and annihilation) operators for longitudinal optical phonons of wave vector $\mathbf{k}$ and energy $\hbar\omega_{\rm LO}$. The V_k are Fourier components of the electron-phonon interaction

$$V_k = -i\frac{\hbar\omega_{\rm LO}}{k}\left(\frac{4\pi\alpha}{V}\right)^{\frac{1}{2}}\left(\frac{\hbar}{2m_b\omega_{\rm LO}}\right)^{\frac{1}{4}}. \tag{1.17}$$

The strength of the electron–phonon interaction is expressed by a dimensionless coupling constant α, which is defined as:

$$\alpha = \frac{e^2}{\hbar}\sqrt{\frac{m_b}{2\hbar\omega_{\rm LO}}}\left(\frac{1}{\varepsilon_\infty} - \frac{1}{\varepsilon_0}\right). \tag{1.18}$$

In this definition, ε_∞ and ε_0 are, respectively, the electronic and the static dielectric constant of the polar crystal.

In Table I the Fröhlich coupling constant is given for a few solids[1].

In deriving the form of V_k, expressions (1.17) and (1.18), it was assumed that (i) the spatial extension of the polaron is large compared to the lattice parameter of the solid ("continuum" approximation), (ii) spin and relativistic effects can be neglected, (iii) the band-electron has parabolic dispersion, (iv) in line with the first approximation it is also assumed that the LO-phonons of interest for the interaction, are the long-wavelength phonons with constant frequency ω_{LO}.

---

[1] In some cases, due to lack of reliable experimental data to determine the electron band mass, the values of α are not well established.

The model, represented by the Hamiltonian (1.16) (which up to now could not been solved exactly) has been the subject of extensive investigations, see, e. g., Refs. [21–28]. In what follows the key approaches of the Fröhlich-polaron theory are briefly reviewed with indication of their relevance for the polaron problems in nanostructures.

D. Infinite mass model [“shift”–operators]

Here some insight will be given in the type of transformation that might be useful to study the Fröhlich Hamiltonian (1.16). For this purpose the Hamiltonian will be treated for a particle with infinite mass $m_b \to \infty$, (which is at $\mathbf{r} = 0$) :

$$H^{\infty} = \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^{+}_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} + V^{*}_{k} a^{\dagger}_{\mathbf{k}}), \tag{1.19}$$

TABLE I: *Electron-phonon coupling constants* (After Ref. [2])

Material	α	Ref.	Material	α	Ref.
InSb	0.023	[5]	AgCl	1.84	[11]
InAs	0.052	[5]	KI	2.5	[5]
GaAs	0.068	[5]	TlBr	2.55	[5]
GaP	0.20	[5]	KBr	3.05	[5]
CdTe	0.29	[6]	$Bi_{12}SiO_{20}$	3.18	[19]
ZnSe	0.43	[5]	CdF_2	3.2	[5]
CdS	0.53	[5]	KCl	3.44	[5]
α-Al_2O_3	1.25	[7]	CsI	3.67	[5]
AgBr	1.53	[11]	$SrTiO_3$	3.77	[20]
α-SiO_2	1.59	[12]	RbCl	3.81	[5]

which can be transformed into the following expression with "shifted" phonon operators:

$$H^{\infty}=\sum_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}\left(a_{\mathbf{k}}^{\dagger}+\frac{V_k}{\hbar\omega_{\mathrm{LO}}}\right)\left(a_{\mathbf{k}}+\frac{V_k^*}{\hbar\omega_{\mathrm{LO}}}\right)-\sum_{\mathbf{k}}\frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}}}. \tag{1.20}$$

To determine the eigenstates of this Hamiltonian, one can perform a unitary transformation which produces the following "shift" of the phonon operators:

$$a_{\mathbf{k}}\to b_{\mathbf{k}}=a_{\mathbf{k}}+\frac{V_k^*}{\hbar\omega_{\mathrm{LO}}},\, a_{\mathbf{k}}^{\dagger}\to b_{\mathbf{k}}^{\dagger}=a_{\mathbf{k}}^{\dagger}+\frac{V_k}{\hbar\omega_{\mathrm{LO}}}.$$

The transformation

$$S=\exp\left[-\sum_{\mathbf{k}}a_{\mathbf{k}}^{\dagger}\frac{V_k^*}{\hbar\omega_{\mathrm{LO}}}+\sum_{\mathbf{k}}\frac{V_k}{\hbar\omega_{\mathrm{LO}}}a_{\mathbf{k}}\right] \tag{1.21}$$

is canonical:

$$S^{\dagger}=\exp\left[-\sum_{\mathbf{k}}a_{\mathbf{k}}\frac{V_k}{\hbar\omega_{\mathrm{LO}}}+\sum_{\mathbf{k}}\frac{V_k^*}{\hbar\omega_{\mathrm{LO}}}a_{\mathbf{k}}^{\dagger}\right]=S^{-1}$$

and has the desired property:

$$S^{-1}a_{\mathbf{k}}S=a_{\mathbf{k}}-\frac{V_k^*}{\hbar\omega_{\mathrm{LO}}},\, S^{-1}a_{\mathbf{k}}^{\dagger}S=a_{\mathbf{k}}^{\dagger}-\frac{V_k}{\hbar\omega_{\mathrm{LO}}}.$$

The transformed Hamiltonian is now:

$$S^{-1}H^{\infty}S=\sum_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}-\sum_{\mathbf{k}}\frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}}}.$$

The eigenstates of the Hamiltonian contain an integer number of phonons $(|n_{\mathbf{k}}\rangle)$ .The eigenenergies are evidently:

$$E=\sum_{\mathbf{k}}n_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}-\sum_{\mathbf{k}}\frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}}}.$$

This expression is divergent at it is often the case in field theory of point charges are considered. A transformation of the type S has been of great interest in developing weak coupling theory as shown below.

E. The "standard" theories

1. *Weak coupling via a perturbation theory*

For actual crystals α-values typically range from $\alpha=0.02$ (InSb) up to $\alpha\sim 3$ to 4 (alkali halides, some oxides), see Table 1. A weak-coupling theory of the polaron was developed originally by Fröhlich [3]. He derived the first weak-coupling perturbation-theory results:

$$E_0 = -\alpha\hbar\omega_{\mathrm{LO}} \tag{1.22}$$

and

$$m^* = \frac{m_b}{1-\alpha/6}. \tag{1.23}$$

Expressions (1.22) and (1.23) are rigorous to order α.

2. Weak coupling via a canonical transformation ["shift"-operators]

Inspired by the work of Tomonaga on quantum electrodynamics (Q. E. D.), Lee, Low and Pines (LLP) [29] analyzed the properties of a weak-coupling polaron starting from a formulation based on canonical transformations (cp. the results of the subsection I D).As hown by them, the unitary transformation

$$U = \exp\left\{\frac{i}{\hbar}\left(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k} a^\dagger_{\mathbf{k}} a_{\mathbf{k}}\right)\cdot\mathbf{r}\right\}, \tag{1.24}$$

where $\mathbf{P}$ is a "c"-number representing the *total system momentum* allows to eliminate the electron co-ordinates from the system. Intuitively one might guess this transformation by writing the exact wave function in the form

$$\Psi_{\mathrm{total}\ H} = \exp\left(\frac{i}{\hbar}\mathbf{p}\cdot\mathbf{r}\right)|\Phi\rangle\,.$$

It is plausible that the "Bloch" factor $\exp\left(i/\hbar\mathbf{p}\cdot\mathbf{r}\right)$ attaches the system to the electron as origin of the co-cordinates. After this transformation the Hamiltonian (1.16) becomes:

$$\mathcal{H} = U^{-1}HU = \frac{\left(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k} a^\dagger_{\mathbf{k}} a_{\mathbf{k}}\right)^2}{2m_b} + \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^\dagger_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} + V_k^* a^\dagger_{\mathbf{k}}). \tag{1.25}$$

If, for the sake of simplicity, the case of total momentum equal to zero is considered, this expression becomes:

$$\mathcal{H} = \sum_{\mathbf{k},\mathbf{k}'} \frac{\hbar^2 \mathbf{k}\cdot\mathbf{k}' a^\dagger_{\mathbf{k}} a^\dagger_{\mathbf{k}'} a_{\mathbf{k}} a_{\mathbf{k}'}}{2m_b} + \sum_{\mathbf{k}} \left(\hbar\omega_{\mathrm{LO}} + \frac{\hbar^2 k^2}{2m_b}\right) a^\dagger_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} + V_k^* a^\dagger_{\mathbf{k}}). \tag{1.26}$$

The first term of this Hamiltonian is the correlation energy term involving different values for $\mathbf{k}$ and $\mathbf{k}'$.If one diagonalizes the second and the trird term of the Hamiltonian (1.26) (this can be done exactly by means of the "shifted-oscillator canonical transformation" S (1.21)), the result of LLP is found. The expectation value of the first term is zero for the wave function $S|0\rangle$. Therefore one is sure to obtain a variational result. It is remarkable that merely extracting the $\mathbf{k} = \mathbf{k}'$ term from the expression

$$\sum_{\mathbf{k},\mathbf{k}'} \frac{\hbar^2 \mathbf{k} \cdot \mathbf{k}' a_{\mathbf{k}}^\dagger a_{\mathbf{k}'}^\dagger a_{\mathbf{k}} a_{\mathbf{k}'}}{2m_b} \tag{1.27}$$

eliminates the divergency from the problem (cp. with the case $m_b \to \infty$) and is equivalent to the sophisticated theory by Lee, Low and Pines (LLP), which corresponds thus to neglect of the term (1.27). The details of the LLP theory are given in Appendix 1. The explicit form for the energy is now

$$E = -\sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} + \frac{\hbar^2 k^2}{2m_b}} = -\alpha\hbar\omega.$$

This self energy is no longer divergent. The divergence is elmininated by the quantum cut-off occurring at $k = \sqrt{2m_b\omega_{\mathrm{LO}}}/\hbar$.

For the self energy the LLP result is equivalent to the perturbation result. The effective mass however is now given by

$$m^* = m_b \left(1 + \frac{\alpha}{6}\right),$$

a result, which follows if one considers the case $\mathbf{P} \neq 0$ and which is also exact for $\alpha \to 0$. However, the LLP effective mass is different from the perturbation result if α insreases.

The LLP approximation has often been called "intermediate-coupling approximation". However its range of validity is the same as that of perturbation theory to order α. The significance of the LLP approximation consists of the flexibility of the canonical transformations together with the fact that it puts the Fröhlich result on a variational basis.

To order α^2, the analytical expressions for the coefficients are α^2: $2\ln(\sqrt{2}+1) - \frac{3}{2}\ln 2 - \frac{\sqrt{2}}{2} \approx -0.01591962$ for the energy and $\frac{4}{3}\ln(\sqrt{2}+1) - \frac{2}{3}\ln 2 - \frac{5\sqrt{2}}{8} + \frac{7}{36} \approx 0.02362763$ for the polaron mass [30].

At present the following weak-coupling expansions are known: for the energy [31, 32]

$$\frac{E_0}{\hbar\omega_{\mathrm{LO}}} = -\alpha - 0.0159196220\alpha^2 - 0.000806070048\alpha^3 - \ldots, \tag{1.28}$$

and for the polaron mass [30]

$$\frac{m^*}{m_b} = 1 + \frac{\alpha}{6} + 0.02362763\alpha^2 + \ldots \tag{1.29}$$

3. Strong coupling via a canonical transformation [“shift”-operators]

Historically, the strong coupling limit was studied before all other treatments (Landau, Pekar [21, 33]). Although it is only a formal case because the actual crystals seems to have α values smaller than 5, it is very interesting because it contains some indication of the intermediate coupling too: approach the excitations from the strong coupling limit and extrapolate to intermediate coupling is interesting because it is expected that some specific strong coupling properties “survive” at intermediate coupling. In what follows, a treatment, equivalent to that of Pekar, but in second quantization and written with as much analogy to the LLP treatment as possible is given.

We start from the Fröhlich Hamiltonian (1.16). At strong coupling one makes the assumption (a “Produkt-Ansatz”) for the polaron wave-function

$$|\Phi\rangle = |\varphi\rangle|f\rangle \tag{1.30}$$

where $|\varphi\rangle$ is the “electron-component” of the wave function ($\langle\varphi|\varphi\rangle = 1$).The “field-component” of the wave function $|f\rangle$ ($\langle f|f\rangle = 1$) parametrically depends on $|\varphi\rangle$. The Produkt-Ansatz (1.30) — or Born-Oppenheimer approximation — implies that the electron adiabatically follows the motion of the atoms, while the field cannot follow the instantaneous motion of the electron. Fröhlich showed that the approximation (1.30) leads to results, which are only valid for sufficiently large $\alpha \to \infty$, i. e. in the strong-coupling regime. A more systematic analysis of strong-coupling polarons based on canonical transformations applied to the Hamiltonian (1.16) was performed in Refs. [34–36].

The expectation value for the energy is now:

$$\langle H\rangle = \langle\varphi|\,\frac{\mathbf{p}^2}{2m_b}\,|\varphi\rangle + \langle f|\left[\sum_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}a^{+}_{\mathbf{k}}a_{\mathbf{k}} + \sum_{\mathbf{k}}(V_k a_{\mathbf{k}}\rho_{\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}}\rho^*_{\mathbf{k}})\right]|f\rangle$$

with

$$\rho_{\mathbf{k}} = \langle\varphi|\,e^{i\mathbf{k}\cdot\mathbf{r}}\,|\varphi\rangle\,.$$

We wish to minimize $\langle H\rangle$, but also

$$\langle f|\left[\sum_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}a^{+}_{\mathbf{k}}a_{\mathbf{k}} + \sum_{\mathbf{k}}(V_k a_{\mathbf{k}}\rho_{\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}}\rho^*_{\mathbf{k}})\right]|f\rangle$$

has to be minimized. This expression will be minimized if $|f\rangle$ is the ground state wave function of the “shifted” oscullator-type Hamiltonian. As we can diagonalize this Hamiltonian exactly:

$$\begin{aligned}
&\sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^{+}_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} \rho_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}} \rho^*_{\mathbf{k}}) && (1.31)\\
&= \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} \left(a^{\dagger}_{\mathbf{k}} + \frac{V_k \rho_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} \right) \left(a_{\mathbf{k}} + \frac{V_k^* \rho^*_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} \right) - \sum_{\mathbf{k}} \frac{|V_k|^2 |\rho_{\mathbf{k}}|^2}{\hbar\omega_{\mathrm{LO}}}, && (1.32)
\end{aligned}$$

we can apply a canonical transformation similar to (1.21):

$$S = \exp\left[\sum_{\mathbf{k}} \left(\frac{V_k \rho_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} a_{\mathbf{k}} - \frac{V_k^* \rho^*_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} a^{\dagger}_{\mathbf{k}} \right) \right], \tag{1.33}$$

which has the property:

$$S^{-1} a_{\mathbf{k}} S = a_{\mathbf{k}} - \frac{V_k^* \rho_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}}, S^{-1} a^{\dagger}_{\mathbf{k}} S = a^{\dagger}_{\mathbf{k}} - \frac{V_k \rho^*_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}}.$$

The transformed Hamiltonian is now:

$$\begin{aligned}
&S^{-1} \left[\sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^{+}_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} \rho_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}} \rho^*_{\mathbf{k}}) \right] S\\
&= \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}} - \sum_{\mathbf{k}} \frac{|V_k|^2 |\rho_{\mathbf{k}}|^2}{\hbar\omega_{\mathrm{LO}}}.
\end{aligned}$$

The phonon vacuum $|0\rangle$ provides a minimum:

$$\langle 0| S^{-1} \left[\sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} a^{+}_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} \rho_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}} \rho^*_{\mathbf{k}}) \right] S |0\rangle = -\sum_{\mathbf{k}} \frac{|V_k|^2 |\rho_{\mathbf{k}}|^2}{\hbar\omega_{\mathrm{LO}}}.$$

Hence, the Hamiltonian (1.31) is minimized by the ground state wave function

$$S|0\rangle = \exp\left[\sum_{\mathbf{k}} \left(\frac{V_k \rho_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} a_{\mathbf{k}} - \frac{V_k^* \rho^*_{\mathbf{k}}}{\hbar\omega_{\mathrm{LO}}} a^{\dagger}_{\mathbf{k}} \right) \right] |0\rangle. \tag{1.34}$$

It gives the ground state energy

$$E_0 = \langle\varphi| \frac{\mathbf{p}^2}{2m_b} |\varphi\rangle - \sum_{\mathbf{k}} \frac{|V_k|^2 |\rho_{\mathbf{k}}|^2}{\hbar\omega_{\mathrm{LO}}}, \tag{1.35}$$

which is still a functional of $|\varphi\rangle$. The functionals $\rho_{\mathbf{k}}$ are different for differerent excitations.

a. Ground state of strong-coupling polarons For the ground state one considers a Gaussian wave function:

$$|\varphi_{"1s"}\rangle = C\exp\left(-\frac{m_b\Omega_0}{2\hbar}r^2\right)$$

with a variational parameter Ω_0.

$$\langle\varphi_{"1s"}|\varphi_{"1s"}\rangle = C^2\int d^3r\exp\left(-\frac{m_b\Omega_0}{\hbar}r^2\right) = C^2\left[\int_{-\infty}^{\infty}dx\exp\left(-\frac{m_b\Omega_0}{\hbar}x^2\right)\right]^3$$

$$= C^2\left(\frac{\sqrt{\pi}}{\frac{m_b\Omega_0}{\hbar}}\right)^3 = C^2\left(\frac{\pi\hbar}{m_b\Omega_0}\right)^{3/2} = 1 \Rightarrow C^2 = \left(\frac{m_b\Omega_0}{\pi\hbar}\right)^{3/2}$$

$$|\varphi_{"1s"}\rangle = \left(\frac{m_b\Omega_0}{\pi\hbar}\right)^{3/4}\exp\left(-\frac{m_b\Omega_0}{2\hbar}r^2\right).$$

For the further use, we introduce a notation $C_1^2 = \left(\frac{m_b\Omega_0}{\pi\hbar}\right)^{1/2}$. Such a wave function is consistent with the localization of the electron, which we expect for large α. The kinetic energy in (1.35) for this function is calculated using the representation of the operator $\mathbf{p}^2 = -\hbar^2\nabla^2 = -\hbar^2\left(\nabla_x^2+\nabla_x^2+\nabla_x^2\right)$:

$$\langle\varphi_{"1s"}|\frac{\mathbf{p}^2}{2m_b}|\varphi_{"1s"}\rangle = -\frac{\hbar^2}{2m_b}C^2\int d^3r\exp\left(-\frac{m_b\Omega_0}{2\hbar}r^2\right)$$

$$\times\left(\nabla_x^2+\nabla_x^2+\nabla_x^2\right)\exp\left(-\frac{m_b\Omega_0}{2\hbar}r^2\right)$$

$$= -3\frac{\hbar^2}{2m_b}C^2\int_{-\infty}^{\infty}dx\exp\left(-\frac{m_b\Omega_0}{2\hbar}x^2\right)\nabla_x^2\exp\left(-\frac{m_b\Omega_0}{2\hbar}x^2\right)$$

$$\times\int_{-\infty}^{\infty}dy\exp\left(-\frac{m_b\Omega_0}{\hbar}y^2\right)\int_{-\infty}^{\infty}dz\exp\left(-\frac{m_b\Omega_0}{\hbar}z^2\right)$$

$$= -3\frac{\hbar^2}{2m_b}C_1^2\int_{-\infty}^{\infty}dx\exp\left(-\frac{m_b\Omega_0}{2\hbar}x^2\right)$$

$$\times\nabla_x\left[-\frac{m_b\Omega_0}{\hbar}x\exp\left(-\frac{m_b\Omega_0}{2\hbar}x^2\right)\right]$$

$$= 3\frac{\hbar^2}{2m_b}C_1^2\frac{m_b\Omega_0}{\hbar}\int_{-\infty}^{\infty}dx\left(1-\frac{m_b\Omega_0}{\hbar}x^2\right)\exp\left(-\frac{m_b\Omega_0}{\hbar}x^2\right)$$

$$== 3\frac{\hbar^2}{m_b}C_1^2\frac{m_b\Omega_0}{\hbar}\int_0^{\infty}dx\left(1-\frac{m_b\Omega_0}{\hbar}x^2\right)\exp\left(-\frac{m_b\Omega_0}{\hbar}x^2\right)$$

$$= 3\frac{\hbar^2}{m_b}C_1^2\frac{m_b\Omega_0}{\hbar}\left[\frac{\sqrt{\pi}}{2\sqrt{\frac{m_b\Omega_0}{\hbar}}}-\frac{m_b\Omega_0}{\hbar}\frac{\sqrt{\pi}}{4\sqrt{\left(\frac{m_b\Omega_0}{\hbar}\right)^3}}\right]$$

$$= 3\frac{\hbar\Omega_0}{4}C_1^2\sqrt{\frac{\pi\hbar}{m_b\Omega_0}} = 3\frac{\hbar\Omega_0}{4}\sqrt{\frac{m_b\Omega_0}{\pi\hbar}}\sqrt{\frac{\pi\hbar}{m_b\Omega_0}} = \frac{3}{4}\hbar\Omega_0.$$

The functional

$$\begin{aligned}
\rho_{\mathbf{k}\text{"}1s\text{"}} &= \langle\varphi_{\text{"}1s\text{"}}|\, e^{i\mathbf{k}\cdot\mathbf{r}}\, |\varphi_{\text{"}1s\text{"}}\rangle = C^2\int d^3r \exp\left(-\frac{m_b\Omega_0}{\hbar}r^2 + i\mathbf{k}\cdot\mathbf{r}\right)\\
&= C^2\int d^3r\exp\left(-\frac{m_b\Omega_0}{\hbar}\left[r^2 + i\frac{\hbar}{m_b\Omega_0}\mathbf{k}\cdot\mathbf{r} - \frac{\hbar^2k^2}{4m_b^2\Omega_0^2} + \frac{\hbar^2k^2}{4m_b^2\Omega_0^2}\right]\right)\\
&= C^2\exp\left(-\frac{m_b\Omega_0}{\hbar}\frac{\hbar^2k^2}{4m_b^2\Omega_0^2}\right)\int d^3r\exp\left(-\frac{m_b\Omega_0}{\hbar}\left[\mathbf{r} + i\frac{\hbar}{2m_b\Omega_0}\mathbf{k}\right]^2\right) \Rightarrow\\
\rho_{\mathbf{k}\text{"}1s\text{"}} &= \exp\left(-\frac{\hbar k^2}{4m_b\Omega_0}\right). \qquad (1.36)
\end{aligned}$$

The second term in (1.35) is then

$$\begin{aligned}
-\sum_{\mathbf{k}}\frac{|V_k|^2\,|\rho_{\mathbf{k}\text{"}1s\text{"}}|^2}{\hbar\omega_{\text{LO}}} &= -\frac{V}{(2\pi)^3}\int d^3k\frac{\hbar\omega_{\text{LO}}}{k^2}\frac{4\pi\alpha}{V}\left(\frac{\hbar}{2m_b\omega_{\text{LO}}}\right)^{\frac{1}{2}}\exp\left(-\frac{\hbar k^2}{2m_b\Omega_0}\right)\\
&= -\frac{\alpha\hbar\omega_{\text{LO}}}{2\pi^2}.4\pi\left(\frac{\hbar}{2m_b\omega_{\text{LO}}}\right)^{\frac{1}{2}}\int_0^\infty dk\exp\left(-\frac{\hbar k^2}{2m_b\Omega_0}\right)\\
&= -\frac{2\alpha\hbar\omega_{\text{LO}}}{\pi}\left(\frac{\hbar}{2m_b\omega_{\text{LO}}}\right)^{\frac{1}{2}}\frac{\sqrt{\pi}}{2}\sqrt{\frac{2m_b\Omega_0}{\hbar}} = -\frac{\alpha\hbar}{\sqrt{\pi}}\sqrt{\Omega_0\omega_{\text{LO}}}. \qquad (1.37)
\end{aligned}$$

The variational energy (1.35) thus becomes

$$E_0 = \frac{3}{4}\hbar\Omega_0 - \frac{\hbar\omega_{\text{LO}}\alpha}{\sqrt{\pi}}\sqrt{\frac{\Omega_0}{\omega_{\text{LO}}}}. \qquad (1.38)$$

Putting

$$\frac{\partial E_0}{\partial\Omega_0} = 0,$$

one obtains

$$\frac{3}{4} = \frac{\alpha}{2\sqrt{\pi}}\sqrt{\frac{\omega_{\text{LO}}}{\Omega_0}} \Longrightarrow \sqrt{\frac{\Omega_0}{\omega_{\text{LO}}}} = \frac{2\alpha}{3\sqrt{\pi}} \Longrightarrow \frac{\Omega_0}{\omega_{\text{LO}}} = \frac{4}{9}\frac{\alpha^2}{\pi} \Longrightarrow$$

$$\Omega_0 = \frac{4}{9}\frac{\alpha^2}{\pi}\omega_{\text{LO}}. \qquad (1.39)$$

Substituting (1.39) in (1.38), we find the ground state energy of the polaron E_0 (calculated with the energy of the uncoupled electron-phonon system as zero energy):

$$E_0 = \frac{3}{4}\hbar\frac{4}{9}\frac{\alpha^2}{\pi}\omega_{\text{LO}} - \frac{\hbar\omega_{\text{LO}}\alpha}{\sqrt{\pi}}\frac{2\alpha}{3\sqrt{\pi}} = \left(\frac{1}{3} - \frac{2}{3}\right)\frac{\alpha^2}{\pi}\hbar\omega_{\text{LO}} \Longrightarrow$$

$$E_0 = -\frac{1}{3}\frac{\alpha^2}{\pi}\hbar\omega_{\text{LO}} = -0.106\alpha^2\hbar\omega_{\text{LO}}. \qquad (1.40)$$

The strong-coupling mass of the polaron, resulting again from the approximation (1.30), is given [37] as:

$$m_0^* = 0.0200\alpha^4 m_b. \qquad (1.41)$$

More rigorous strong-coupling expansions for E_0 and m^* have been presented in the literature [38]:

$$\frac{E_0}{\hbar\omega_{\rm LO}} = -0.108513\alpha^2 - 2.836, \tag{1.42}$$

$$\frac{m_0^*}{m_b} = 1 + 0.0227019\alpha^4. \tag{1.43}$$

The strong-coupling ground state energy (1.40) is lower than the LLP ground state energy for $\alpha > 10$.

b. The excited states of the polaron: SS, FC, RES In principle, excited states of the polaron exist at all coupling. In the general case, and for simplicity for $\mathbf{P} = 0$, a continuum of states starts at $\hbar\omega_{\rm LO}$ above the ground state of the polaron. This continuum physically corresponds to the scattering of free phonons on the polaron. Those "scattering.states" (SS) were studied in [39] anf for the first time more generally in [37] are not the only excitations of the polaron. There are also *internal* excitation states corresponding to the excitations of the electron in the potential it created itself. By analogy with the excited states of colour centers, the following terminology is used.

(i) The states where the electron is excited in the potential belonging to the ground state configuration of the lattice are called *Franck-Condon* (FC) states

(ii) Excitations of the electron in which the lattice polarization is adapted to the electronic configuration of the excited electron (which itself then adapts its wave function to the new potential, etc. ...leading to a self-consistent final state), are called *relaxed excited state* (RES) [21].

c. Calculation of the lowest FC state The formalism used until now is well adapted to treat the polaron excitations at strong coupling. The field dependence of the wave function is (1.34). For the FC state the $\rho_{\mathbf{k}}$ are the same as for the ground state (1.36). Physically $\rho_{\mathbf{k}}$ tells us, to what electronic distribution the field is adapted. The electronic part of the excited wave function is $2p$-like:

$$|\varphi_{"2p"}\rangle = C_{"2p"} z \exp\left(-\frac{m_b\Omega_p}{2\hbar} r^2\right) \tag{1.44}$$

with a parameter Ω_p, which is equal to Ω_0 :

$$\langle \varphi_{"2p"} | \varphi_{"2p"} \rangle = C^2_{"2p"} \int_{-\infty}^{\infty} dz z^2 \exp\left(-\frac{m_b \Omega_p}{\hbar} z^2\right) \left[\int_{-\infty}^{\infty} dx \exp\left(-\frac{m_b \Omega_p}{\hbar} x^2\right)\right]^2$$

$$= C^2_{"2p"} \left(\frac{\sqrt{\pi}}{\frac{m_b \Omega_p}{\hbar}}\right)^2 \frac{\sqrt{\pi}}{2\left(\frac{m_b \Omega_p}{\hbar}\right)^{3/2}} = C^2_{"2p"} \left(\frac{\pi \hbar}{m_b \Omega_p}\right)^{3/2} \frac{\hbar}{2 m_b \Omega_p} = 1 \Rightarrow$$

$$C^2_{"2p"} = \left(\frac{m_b \Omega_p}{\pi \hbar}\right)^{3/2} \frac{2 m_b \Omega_p}{\hbar}$$

$$|\varphi_{"2p"}\rangle = \left(\frac{m_b \Omega_0}{\pi \hbar}\right)^{3/4} \left(\frac{2 m_b \Omega_p}{\hbar}\right)^{1/2} z \exp\left(-\frac{m_b \Omega_0}{2\hbar} r^2\right).$$

We introduce still a notation

$$C_2^2 = \frac{2}{\sqrt{\pi}} \left(\frac{m_b \Omega_p}{\hbar}\right)^{3/2}$$

The FC state energy is, similarly to (1.35),

$$E_{FC} = \langle \varphi_{"2p"} | \frac{\mathbf{p}^2}{2 m_b} | \varphi_{"2p"} \rangle - \sum_{\mathbf{k}} \frac{|V_k|^2 |\rho_{\mathbf{k}"1s"}|^2}{\hbar \omega_{\mathrm{LO}}}. \tag{1.45}$$

The kinetic energy term is

$$
\begin{aligned}
\left\langle \varphi_{``2p"} \right| \frac{\mathbf{p}^2}{2m_b} \left| \varphi_{``2p"} \right\rangle &= -\frac{\hbar^2}{2m_b} C^2_{``2p"} \int d^3r\, z \exp\left(-\frac{m_b\Omega_p}{2\hbar} r^2\right) \\
&\times \left(\nabla_x^2 + \nabla_x^2 + \nabla_x^2\right) \left[z \exp\left(-\frac{m_b\Omega_p}{2\hbar} r^2\right)\right] \\
&= -\frac{\hbar^2}{2m_b} C^2_{``2p"} \int_{-\infty}^{\infty} dz z \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right) \nabla_z^2 \left[z \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right)\right] \\
&\times \int_{-\infty}^{\infty} dy \exp\left(-\frac{m_b\Omega_p}{\hbar} y^2\right) \int_{-\infty}^{\infty} dz \exp\left(-\frac{m_b\Omega_p}{\hbar} z^2\right) + \frac{2}{4}\hbar\Omega_p \\
&= -\frac{\hbar^2}{2m_b} C_2^2 \int_{-\infty}^{\infty} dz z \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right) \\
&\times \nabla_z \left[\left(1 - \frac{m_b\Omega_p}{\hbar} z^2\right) \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right)\right] + \frac{1}{2}\hbar\Omega_p \\
&= \frac{\hbar^2}{2m_b} C_2^2 \int_{-\infty}^{\infty} dz z \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right) \\
&\times \left[\frac{2m_b\Omega_p}{\hbar} z + \left(1 - \frac{m_b\Omega_p}{\hbar} z^2\right) \frac{m_b\Omega_p}{\hbar} z \right] \exp\left(-\frac{m_b\Omega_p}{2\hbar} z^2\right) \\
+\frac{1}{2}\hbar\Omega_p &= \frac{\hbar^2}{2m_b} C_2^2 \int_{-\infty}^{\infty} dz \left(3 - \frac{m_b\Omega_p}{\hbar} z^2\right) \frac{m_b\Omega_p}{\hbar} z^2 \exp\left(-\frac{m_b\Omega_p}{\hbar} z^2\right) \\
+\frac{1}{2}\hbar\Omega_p &= \frac{\hbar\Omega_p}{2} C_2^2 \left[\frac{3\sqrt{\pi}}{2\left(\frac{m_b\Omega_p}{\hbar}\right)^{3/2}} - \frac{m_b\Omega_p}{\hbar} \frac{3\sqrt{\pi}}{4\left(\frac{m_b\Omega_p}{\hbar}\right)^{5/2}} \right] + \frac{1}{2}\hbar\Omega_p \\
&= \frac{\hbar\Omega_p}{2} C_2^2 \frac{3\sqrt{\pi}}{4\left(\frac{m_b\Omega_p}{\hbar}\right)^{3/2}} + \frac{1}{2}\hbar\Omega_p \\
&= \frac{\hbar\Omega_p}{2} \frac{2}{\sqrt{\pi}} \left(\frac{m_b\Omega_p}{\hbar}\right)^{3/2} \frac{3\sqrt{\pi}}{4\left(\frac{m_b\Omega_p}{\hbar}\right)^{3/2}} + \frac{1}{2}\hbar\Omega_p \\
&= \frac{3}{4}\hbar\Omega_p + \frac{1}{2}\hbar\Omega_p = \frac{5}{4}\hbar\Omega_p. \qquad (1.46)
\end{aligned}
$$

For the FC state, $\Omega_p = \Omega_0$. The second term in (1.45) is precisely (1.37),

$$
-\sum_{\mathbf{k}} \frac{\left|V_k\right|^2 \left|\rho_{\mathbf{k}``1s"}\right|^2}{\hbar\omega_{\mathrm{LO}}} = -\frac{\hbar}{\sqrt{\pi}} \sqrt{\Omega_0 \omega_{\mathrm{LO}}},
$$

and the FC energy (1.45) becomes

$$
\begin{aligned}
E_{FC} &= \frac{5}{4}\hbar\Omega_0 - \frac{\hbar\omega_{\mathrm{LO}}\alpha}{\sqrt{\pi}} \sqrt{\frac{\Omega_0}{\omega_{\mathrm{LO}}}} \\
&= \frac{5}{4}\hbar\frac{4}{9}\frac{\alpha^2}{\pi}\omega_{\mathrm{LO}} - \frac{2}{3}\frac{\alpha^2}{\pi}\hbar\omega_{\mathrm{LO}} = \left(\frac{5}{9} - \frac{2}{3}\right)\frac{\alpha^2}{\pi}\hbar\omega_{\mathrm{LO}} \Longrightarrow
\end{aligned}
$$

The energy of the lowest FC state is, within the Produkt-Ansatz [40]:

$$E_{\mathrm{FC}} = \frac{\alpha^2}{9\pi}\hbar\omega_{\mathrm{LO}} = 0.0354\alpha^2\hbar\omega_{\mathrm{LO}}. \tag{1.47}$$

The fact that this energy is positive, is presumably due to the choice of a harmonic potential. The real potential the electron sees is anharmonic, and a bound state may be expected.

d. Calculation of RES The electronic part of the excited wave function is (1.44) with a variational parameter Ω_p,which is determined below. The variational RES energy is, similarly to (1.35),

$$E_{RES} = \langle\varphi_{"2p"}|\,\frac{\mathbf{p}^2}{2m_b}\,|\varphi_{"2p"}\rangle - \sum_{\mathbf{k}} \frac{|V_k|^2\,|\rho_{\mathbf{k}"2p"}|^2}{\hbar\omega_{\mathrm{LO}}}. \tag{1.48}$$

Here the kinetic energy term is given by Eq. (1.46). The functional, which is now needed, is

$$\begin{aligned}
\rho_{\mathbf{k}"2p"} &= \langle\varphi_{"2p"}|\,e^{i\mathbf{k}\cdot\mathbf{r}}\,|\varphi_{"2p"}\rangle = C^2_{"2p"}\int d^3r z^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}r^2 + i\mathbf{k}\cdot\mathbf{r}\right)\\
&= C^2_{"2p"}\int d^3r z^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}\left[r^2 + i\frac{\hbar}{m_b\Omega_p}\mathbf{k}\cdot\mathbf{r} - \frac{\hbar^2k^2}{4m_b^2\Omega_p^2} + \frac{\hbar^2k^2}{4m_b^2\Omega_p^2}\right]\right)\\
&= C^2_{"2p"}\ exp\left(-\frac{m_b\Omega_p}{\hbar}\frac{\hbar^2k^2}{4m_b^2\Omega_p^2}\right)\int d^3r z^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}\left[\mathbf{r} + i\frac{\hbar}{2m_b\Omega_p}\mathbf{k}\right]^2\right) \Rightarrow
\end{aligned}$$

$$\begin{aligned}
\rho_{\mathbf{k}"2p"} &= \exp\left(-\frac{\hbar k^2}{4m_b\Omega_p}\right) C_2^2 \int_{-\infty}^{\infty} dz z^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}\left[z + i\frac{\hbar}{2m_b\Omega_p}k_z\right]^2\right)\\
&= .\exp\left(-\frac{\hbar k^2}{4m_b\Omega_p}\right) C_2^2 \int_{-\infty}^{\infty} dz \left(z - i\frac{\hbar}{2m_b\Omega_p}k_z\right)^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}z^2\right)\\
&= \exp\left(-\frac{\hbar k^2}{4m_b\Omega_p}\right) C_2^2 \int_{-\infty}^{\infty} dz \left[z^2 - \left(\frac{\hbar}{2m_b\Omega_p}k_z\right)^2\right]^2 \exp\left(-\frac{m_b\Omega_p}{\hbar}z^2\right)\\
&= \exp\left(-\frac{\hbar k^2}{4m_b\Omega_p}\right) C_2^2 \left\{\frac{\sqrt{\pi}}{2\left(\frac{m_b\Omega_p}{\hbar}\right)^{3/2}} - \left(\frac{\hbar}{2m_b\Omega_p}k_z\right)^2 \frac{\sqrt{\pi}}{\left(\frac{m_b\Omega_p}{\hbar}\right)^{1/2}}\right\}\\
&= \left(1 - \frac{\hbar k_z^2}{2m_b\Omega_p}\right)\exp\left(-\frac{\hbar k^2}{4m_b\Omega_p}\right).
\end{aligned} \tag{1.49}$$

Further, we substitute (1.49) in the second term in the r.h.s. of Eq. (1.48):

$$
\begin{aligned}
-\sum_{\mathbf{k}} \frac{|V_k|^2 \left|\rho_{\mathbf{k}\text{"}1s\text{"}}\right|^2}{\hbar\omega_{\mathrm{LO}}} &= -\frac{V}{(2\pi)^3}\int d^3k \frac{\hbar\omega_{\mathrm{LO}}}{k^2}\frac{4\pi\alpha}{V}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}} \\
&\times \left(1-\frac{\hbar k_z^2}{2m_b\Omega_p}\right)^2 \exp\left(-\frac{\hbar k^2}{2m_b\Omega_p}\right) \\
&= -\frac{\alpha\hbar\omega_{\mathrm{LO}}}{2\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}} 2\pi\int_{-1}^{1} dx \\
&\times \int_0^\infty dk \left(1-\frac{\hbar k^2x^2}{m_b\Omega_p}+\frac{\hbar^2k^4x^4}{4m_b^2\Omega_p^2}\right)\exp\left(-\frac{\hbar k^2}{2m_b\Omega_p}\right) \\
&= -\frac{\alpha\hbar\omega_{\mathrm{LO}}}{\pi}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}} \\
&\times \int_0^\infty dk \left(2-\frac{2}{3}\frac{\hbar k^2}{m_b\Omega_p}+\frac{2}{5}\frac{\hbar^2k^4}{4m_b^2\Omega_p^2}\right)\exp\left(-\frac{\hbar k^2}{2m_b\Omega_p}\right) \\
&= -\frac{\alpha\hbar\omega_{\mathrm{LO}}}{\pi}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}} \\
&\times \left[\frac{\sqrt{\pi}}{\left(\frac{\hbar}{2m_b\Omega_p}\right)^{1/2}}-\frac{1}{3}\frac{\hbar}{m_b\Omega_p}\frac{\sqrt{\pi}}{2\left(\frac{\hbar}{2m_b\Omega_p}\right)^{3/2}}+\frac{1}{5}\frac{\hbar^2}{4m_b^2\Omega_p^2}\frac{3\sqrt{\pi}}{4\left(\frac{\hbar}{2m_b\Omega_p}\right)^{5/2}}\right]
\end{aligned}
$$

$$
\begin{aligned}
&= -\frac{\alpha\hbar}{\sqrt{\pi}}\sqrt{\omega_{\mathrm{LO}}\Omega_p}\left[1-\frac{1}{3}+\frac{3}{20}\right] \\
&= -\frac{\alpha\hbar}{\sqrt{\pi}}\sqrt{\omega_{\mathrm{LO}}\Omega_p}\frac{60-20+9}{60} = -\frac{49}{60}\frac{\alpha\hbar}{\sqrt{\pi}}\sqrt{\omega_{\mathrm{LO}}\Omega_p}
\end{aligned}
$$

The variational energy (1.48) becomes

$$E_{RES} = \frac{5}{4}\hbar\Omega_p - \frac{49}{60}\frac{\alpha\hbar}{\sqrt{\pi}}\sqrt{\omega_{\mathrm{LO}}\Omega_p}.$$

Putting

$$\frac{\partial E_{RES}}{\partial\Omega_p} = 0,$$

one obtains

$$\frac{5}{4} = \frac{49\alpha}{120\sqrt{\pi}}\sqrt{\frac{\omega_{\mathrm{LO}}}{\Omega_p}} \Longrightarrow \sqrt{\frac{\Omega_p}{\omega_{\mathrm{LO}}}} = \frac{49\alpha}{150\sqrt{\pi}} \Longrightarrow \frac{\Omega_p}{\omega_{\mathrm{LO}}} = \left(\frac{49}{150}\right)^2\frac{\alpha^2}{\pi} \Longrightarrow$$

$$
\begin{aligned}
E_{RES} &= \frac{5}{4}\hbar\left(\frac{49}{150}\right)^2\frac{\alpha^2}{\pi}\omega_{\mathrm{LO}} - \frac{\hbar\omega_{\mathrm{LO}}\alpha}{\sqrt{\pi}}\frac{49}{60}\frac{49\alpha}{150\sqrt{\pi}} = \left(\frac{5}{4}-\frac{5}{2}\right)\left(\frac{49}{150}\right)^2\frac{\alpha^2}{\pi}\hbar\omega_{\mathrm{LO}} \\
&= -\frac{5}{4}\left(\frac{49}{150}\right)^2\frac{\alpha^2}{\pi}\hbar\omega_{\mathrm{LO}} = -\frac{49^2}{120\times 150}\frac{\alpha^2}{\pi}\hbar\omega_{\mathrm{LO}} \Longrightarrow
\end{aligned}
$$

The energy of the RES is (see Refs. [37, 39]):

$$E_{\mathrm{RES}} = -0.042\alpha^2\hbar\omega_{\mathrm{LO}}. \tag{1.50}$$

The effective mass of the polaron in the RES is given [37] as:

$$m^*_{RES} = 0.621\frac{\alpha^4}{81\pi^2}m_b = 0.0200\alpha^4 m_b. \tag{1.51}$$

The structure of the energy spectrum of the strong-coupling polaron is shown in Fig. 2.



FIG. 2: Structure of the energy spectrum of a polaron at strong coupling: E_0 — the ground state, E_{RES} — the (first) relaxed excited state; the Franck-Condon states (E_{FC}). In fact, both the Franck-Condon states and the relaxed excited states lie in the continuum and, strictly speaking, are resonances.

The significance of the strong-coupling large polaron theory is formal only: it allows to test "all-coupling" theories in the limit $\alpha \to \infty$. Remarkably, the effective electron-phonon coupling strength significantly increases in systems of low dimension and low dimensionality.

4. All-coupling theory. The Feynman path integral

Feynman developed a superior all-coupling polaron theory using his path-integral formalism [41]. He studied first the self-energy E_0 and the effective mass m^* of polarons [41].

Feynman got the idea to formulate the polaron problem into the Lagrangian form of quantum mechanics and then eliminate the field oscillators, "... in exact analogy to Q. E. D. ... (resulting in) ... a sum over all trajectories ... ". The resulting path integral (here limited to the ground-state properties) is of the form (Ref. [41]):

$$\langle 0,\beta|0,0\rangle = \int \mathcal{D}\mathbf{r}(\tau)\exp\left[-\frac{1}{2}\int_0^\beta \dot{\mathbf{r}}^2 d\tau + \frac{\alpha}{2^{\frac{3}{2}}}\int_0^\beta\int_0^\beta \frac{e^{-|\tau-\sigma|}}{|\mathbf{r}(\tau)-\mathbf{r}(\sigma)|}d\tau d\sigma\right], \tag{1.52}$$

where $\beta = 1/(k_B T)$. (1.52) gives the amplitude that an electron found at a point in space at time zero will appear at the same point at the (imaginary) time β. This path integral (1.52) has a great intuitive appeal: it shows the polaron problem as an equivalent one-particle problem in which the interaction, non-local in time or "retarded", occurs between the electron and itself. Subsequently Feynman showed how the variational principle of quantum mechanics could be adapted to the path-integral formalism and he introduced a quadratic trial action (non-local in time) to simulate (1.52).

Applying the variational principle for path integrals then results in an upper bound for the polaron self-energy at all α, which at weak and strong coupling gives accurate expressions. Feynman obtained smooth interpolation between a weak and strong coupling (for the ground state energy). The weak-coupling expansions of Feynman for the ground-state energy and the effective mass of the polaron are:

$$\frac{E_0}{\hbar\omega_{\mathrm{LO}}} = -\alpha - 0.0123\alpha^2 - 0.00064\alpha^3 - \ldots\ (\alpha \to 0), \tag{1.53}$$

$$\frac{m^*}{m_b} = 1 + \frac{\alpha}{6} + 0.025\alpha^2 + \ldots\ (\alpha \to 0). \tag{1.54}$$

In the strong-coupling limit Feynman found for the ground-state energy energy:

$$\frac{E_0}{\hbar\omega_{\mathrm{LO}}} \equiv \frac{E_{3D}(\alpha)}{\hbar\omega_{\mathrm{LO}}} = -0.106\alpha^2 - 2.83 - \ldots \quad (\alpha \to \infty) \tag{1.55}$$

and for the polaron mass:

$$\frac{m^*}{m_b} \equiv \frac{m^*_{3D}(\alpha)}{m_b} = 0.0202\alpha^4 + \ldots \quad (\alpha \to \infty). \tag{1.56}$$

Over the years the Feynman model for the polaron has remained the most successful approach to this problem. The analysis of an exactly solvable ("symmetrical") 1D-polaron model [39, 42], Monte Carlo schemes [28, 43] and other numerical schemes [44] demonstrate the remarkable accuracy of Feynman's path-integral approach to the polaron ground-state energy. Experimentally more directly accessible properties of the polaron, such as its mobility and optical absorption, have been investigated subsequently. Within the path-integral approach, Feynman et al. studied later the mobility of polarons [45, 46]. Subsequently the path-integral approach to the polaron problem was generalized and developed to become a powerful tool to study optical absorption, magnetophonon resonance and cyclotron resonance [47–51].

In Ref. [52], a self-consistent treatment for the polaron problem at all α was presented, which is based on the Heisenberg equations of motion starting from a trial expression for the electron position. It was used to derive the effective mass and the optical properties of the polaron at arbitrary coupling. A variational justification of the approximation used in Ref. [52] (through a Stiltjes continuous fraction) is reproduced in Appendix 2.

5. *On Monte Carlo calculations of the polaron free energy*

In Ref. [53], using a Monte Carlo calculation, the ground-state energy of a polaron was derived as $E_0 = \lim_{\beta\to\infty} \Delta F$, where $\Delta F = F_\beta - F_\beta^0$ with F_β the free energy per polaron and $F_\beta^0 = [3/(2\beta)]\ln(2\pi\beta)$ the free energy per electron. The value $\beta\hbar\omega_{\mathrm{LO}} = 25$, used for the actual computation in Ref. [53], corresponds to $T/T_D = 0.04$ ($T_D = \hbar\omega_{\mathrm{LO}}/k_B$; $\hbar\omega_{\mathrm{LO}}$ is the LO phonon energy). So, as pointed out in Ref. [54], the authors of Ref. [53] actually calculated the *free energy* ΔF, rather than the polaron *ground-state energy.*

To investigate the importance of temperature effects on ΔF, the authors of Ref. [54] considered the polaron energy as obtained by Osaka [55], who generalized the Feynman [41] polaron theory to nonzero temperatures:

$$\begin{aligned}\frac{\Delta F}{\hbar\omega} = \frac{3}{\beta}\ln\left(\frac{w}{v}\frac{\sinh\frac{\beta_0 v}{2}}{\sinh\frac{\beta_0 w}{2}}\right) - \frac{3}{4}\frac{v^2-w^2}{v}\left(\coth\frac{\beta_0 v}{2} - \frac{2}{\beta_0 v}\right)\\ - \frac{\alpha}{\sqrt{2\pi}}\left[1+n\left(\omega_{\mathrm{LO}}\right)\right]\int_0^{\beta_0} du\frac{e^{-u}}{\sqrt{D\left(u\right)}},\end{aligned} \tag{1.57}$$

where $\beta_0 = \beta\hbar\omega_{\mathrm{LO}}$, $n(\omega) = 1/\left(e^{\beta\hbar\omega}-1\right)$, and

$$D\left(u\right) = \frac{w^2}{v^2}\frac{u}{2}\left(1-\frac{u}{\beta_0}\right) + \frac{v^2-w^2}{2v^3}\left(1-e^{-vu}-4n\left(v\right)\sinh^2\frac{vu}{2}\right). \tag{1.58}$$

This result is variational, with variational parameters v and w, and gives an upper bound to the exact polaron free energy.

The results of a numerical-variational calculation of Eq. (1.57) are shown in Fig. 3, where the free energy $-\Delta F$ is plotted (in units of $\hbar\omega_{\rm LO}$) as a function of α for different values of the lattice temperature. As seen from Fig. 3, (i) $-\Delta F$ increases with increasing temperature and (ii) the effect of temperature on ΔF increases with increasing α.



FIG. 3: Contribution of the electron-phonon interaction to the free energy of the Feynman polaron as a function of the electron-phonon coupling constant α for different values of the lattice temperature. Inset: temperature dependence of the free energy for $\alpha = 3$. (From Ref. [54].)

In Table II, the Monte Carlo results [53], $(\Delta F)_{\rm MC}$, are compared with the free energy of the Feynman polaron, $(\Delta F)_{\rm F}$, calculated in [54]. The values for the free energy obtained from the Feynman polaron model are *lower* than the MC results for $\alpha \lesssim 2$ and $\alpha \geq 4$ (but lie within the 1% error of the Monte Carlo results). Since the Feynman result for the polaron free energy is an upper bound to the exact result, we conclude that for $\alpha \lesssim 2$ and $\alpha \geq 4$ the results of the Feynman model are closer to the exact result than the MC results of [53].

TABLE II: Comparison between the free energy of the Feynman polaron theory, $-(\Delta F)_{\rm F}$, and the Monte Carlo results of Ref. [53], $-(\Delta F)_{\rm MC}$, for $T/T_D = 0.04$. The relative difference is defined as $\Delta = 100 \times [(\Delta F)_{\rm F} - (\Delta F)_{\rm MC}]/(\Delta F)_{\rm MC}$. (From Ref. [54])

α	$-(\Delta F)_{\rm F}$	$-(\Delta F)_{\rm MC}$	Δ (%)
0.5	0.50860	0.505	0.71
1.0	1.02429	1.020	0.42
1.5	1.54776	1.545	0.18
2.0	2.07979	2.080	−0.010
2.5	2.62137	2.627	−0.21
3.0	3.17365	3.184	−0.32
3.5	3.73814	3.747	−0.24
4.0	4.31670	4.314	0.063

6. *On the contributions of the N-phonon states to the polaron ground state*

The analysis of an exactly solvable ("symmetric") 1D-polaron model was performed in Refs. [39, 56, 57]. The model consists of an electron interacting with two oscillators possessing the opposite wave vectors: **k** and -**k**.The parity operator, which changes $a_{\mathbf{k}}$ and $a_{-\mathbf{k}}$ (and also $a_{\mathbf{k}}^{\dagger}$ and $a_{-\mathbf{k}}^{\dagger}$), commutes with the Hamiltonian of the system. Hence, the polaron states are classified into the even and odd ones with the eigenvalues of the parity operator +1 and −1, respectively. For the lowest even and odd states, the phonon distribution functions W_N are plotted in Fig. 1, upper panel, at some values of the effective coupling constant λ of the "symmetric" model. The value of the parameter

$$\varkappa = \sqrt{\frac{(\hbar k)^2}{m_b \hbar \omega_{\rm LO}}}$$

for these graphs was taken 1, while the total polaron momentum $\mathbf{P} = 0$. In the weak-coupling case ($\lambda \approx 0.6$) W_N is a decaying function of N. When increasing λ, W_N acquires a maximum, e.g. at $N = 8$ for the lowest even state at $\lambda \approx 5.1$. The phonon distribution function W_N has the same character for the lowest even and the lowest odd states at all values of the number of the virtual phonons in the ground state. (as distinct from the higher states). This led to the conclusion that the lowest odd state is an internal excited state of the polaron.

In Ref [28], the structure of the polaron cloud was investigated using the diagrammatic quantum Monte Carlo (DQMC) method. In particular, partial contributions of N-phonon states to the polaron ground state were found as a function of N for a few values of the coupling constant α, see Fig. 1, lower panel. It was shown to gradually evolve from the weak-coupling case ($\alpha = 1$) into the strong-coupling regime ($\alpha = 17$). Comparion of the lower panel to the upper panel in Fig. 4 clearly shows that the evolution of the shape and the scale of the distribution of the N-phonon states with increasing α as derived for a large polaron within DQMC method [28] is *in remarkable agreement* with the results obtained within the "symmetric" 1D polaron model [39, 56, 57].





FIG. 4: *Upper panel*: The phonon distribution functions W_N in the "symmetric" polaron model for various values of the effective coupling constant λ at $\kappa = 1, \mathbf{P} = 0$ (from [56], Fig. 23). *Lower panel*: Distribution of multiphonon states in the polaron cloud within DQMC method for various values of α (from [28], Fig. 7).

F. Polaron mobility

The mobility of large polarons was studied within various theoretical approaches(see Ref. [58] for the detailed references). Fröhlich [59] pointed out the typical behavior of the large-polaron mobility

$$\mu \propto \exp(\hbar\omega_{\mathrm{LO}}\beta), \tag{1.59}$$

which is characteristic for weak coupling. Here, $\beta = 1/k_B T$, T is the temperature. Within the weak-coupling regime, the mobility of the polaron was then derived, e. g., using the Boltzmann equation in Refs. [60, 61] and starting from the LLP-transformation in Ref. [62].

A nonperturbative analysis was embodied in the Feynman polaron theory, where the mobility μ of the polaron using the path-integral formalism was derived by Feynman et al. (usually referred to as FHIP) as a static limit starting from a frequency-dependent impedance function. For sufficiently low temperature T the mobility then takes the form [45]

$$\mu = \left(\frac{w}{v}\right)^3 \frac{3e}{4m_b\hbar\omega_{\mathrm{LO}}^2\alpha\beta} \mathrm{e}^{\hbar\omega_{\mathrm{LO}}\beta} \exp\{(v^2 - w^2)/w^2 v\}\ , \tag{1.60}$$

where v and w are (variational) functions of α obtained from the Feynman polaron model.

Using the Boltzmann equation for the Feynman polaron model, Kadanoff [63] found the mobility, which for low temperatures can be represented as follows:

$$\mu = \left(\frac{w}{v}\right)^3 \frac{e}{2m_b\omega_{\mathrm{LO}}\alpha} \mathrm{e}^{\hbar\omega_{\mathrm{LO}}\beta} \exp\{(v^2 - w^2)/w^2 v\}\ , \tag{1.61}$$

The weak-coupling perturbation expansion of the low-temperature polaron mobility as found using the Green's function technique [64] has confirmed that the mobility derived from the Boltzmann equation is exceedingly exact for weak coupling ($\alpha/6 \ll 1$) and at low temperatures ($k_B T \ll \hbar\omega_{\mathrm{LO}}$). As shown in Ref. [63], the mobility of Eq. (1.60) differs by the factor of $3/(2\beta\hbar\omega_{\mathrm{LO}})$ from that derived using the polaron Boltzmann equation as given by Eq. (1.61).

In the limit of weak electron-phonon coupling and low temperature, the FHIP polaron mobility of Eq. (1.60) differs by the factor of $3/(2\beta\hbar\omega_{\mathrm{LO}})$ from the previous result [60–62], which, as pointed out in Ref. [45] and in later publications (see, e.g., Refs.[58, 63, 64]), is correct for $\beta \gg 1$. As follows from this comparison, the result of Ref. [45] is not valid when $T \to 0$. As argued in Ref. [45] and later confirmed, in particular, in Ref. [65] the above discrepancy can be attributed to an interchange of two limits in calculating the impedance. In FHIP, for weak electron-phonon coupling, one takes $\lim_{\Omega\to 0}\lim_{\alpha\to 0}$, whereas the correct order is $\lim_{\alpha\to 0}\lim_{\Omega\to 0}$ (Ω is the frequency of the applied electric field). It turns out that for the correct result the mobility at low temperatures is predominantly limited by the absorption of phonons, while in the theory of FHIP it is the emission of phonons which gives the dominant contribution as T goes to zero [65].

The analysis based on the Boltzmann equation takes into account the phonon emission processes whenever the energy of the polaron is above the emission threshold. The independent-collision model, which underlies the Boltzmann-equation approach, however, fails in the "strong coupling regime" of the large polaron, when the thermal mean free path becomes less than the de Broglie wavelength; in this case, the Boltzmann equation cannot be expected to be adequate [45, 66].

In fact, the expression (1.60) for the polaron mobility was reported to adequately describe the experimental data in several polar materials (see, e.g., Refs. [66–68]). Experimental work on alkali halides and silver halides indicates that the mobility obtained from Eq. (1.60) describes the experimental results quite accurately [67]. Measurements of mobility as a function of temperature for photoexcited electrons in cubic n-type $Bi_{12}SiO_{20}$ are explained well in terms of large polarons within the Feynman approach [66]. The experimental findings on electron transport in crystalline TiO_2 (rutile phase) probed by THz time-domain spectroscopy are quantitatively interpreted within the Feynman model [68]. One of the reasons for the agreement between theory based on Eq. (1.60) and experiment is that in the path-integral approximation to the polaron mobility, a Maxwellian distribution for the electron velocities is assumed, when applying the adiabatic switching on of the Fröhlich interaction. Although such a distribution is not inherent in the Fröhlich interaction, its incorporation tends to favor agreement with experiment because other mechanisms (interaction with acoustic phonons etc.) cause a Gaussian distribution.

II. OPTICAL ABSORPTION. WEAK COUPLING

A. Optical absorption at weak coupling [within the perturbation theory]

At zero temperature and in the weak-coupling limit, the optical absorption is due to the elementary polaron scattering process, schematically shown in Fig. 5.



FIG. 5: Elementary polaron scattering process describing the absorption of an incoming photon and the generation of an outgoing phonon.

In the weak-coupling limit ($\alpha \ll 1$) the polaron absorption coefficient was first obtained by V. Gurevich, I. Lang and Yu. Firsov [69], who started from the Kubo formula. Their optical-absorption coefficient is equivalent to a particular case of the result of J. Tempere and J. T. Devreese (Ref. [51]), with the dynamic srtucture factor $S(\mathbf{q},\omega)$ corresponding to the Hartree-Fock approximation (see also [70], p. 585). At zero temperature, the absorption coefficient for absorption of light with frequency Ω can be expressed in terms of elementary functions in two limiting cases: in the region of comparatively high polaron densities ($\hbar(\Omega-\omega_{\text{LO}})/\zeta \ll 1$)

$$\Gamma(\omega)=\frac{1}{\epsilon_0 nc}\frac{2^{1/2}N^{2/3}\alpha}{(3\pi^2)^{1/3}}\frac{e^2}{(\hbar m_b\omega_{\text{LO}})^{1/2}}\frac{\omega-1}{\omega^3}\Theta(\omega-1), \tag{2.1}$$

and in the low-concentration region ($\hbar(\Omega-\omega_{\text{LO}})/\zeta \gg 1$)

$$\Gamma(\omega)=\frac{1}{\epsilon_0 nc}\frac{2Ne^2\alpha}{3m_b\omega_{\text{LO}}}\frac{(\omega-1)^{1/2}}{\omega^3}\Theta(\omega-1), \tag{2.2}$$

where $\omega=\Omega/\omega_{\text{LO}}$, ϵ_0 is the dielectric permittivity of the vacuum, n is the refractive index of the medium, N is the concentration of polarons and ζ is the Fermi level for the electrons. A step function

$$\Theta(\omega-1)=\begin{cases}1 \text{ if } \omega>1,\\ 0 \text{ if } \omega<1\end{cases}$$

reflects the fact that at zero temperature the absorption of light accompanied by the emission of a phonon can occur only if the energy of the incident photon is larger than that of a phonon ($\omega > 1$). In the weak-coupling limit, according to Eqs. (2.1), (2.2), the absorption spectrum consists of a "one-phonon line".

At nonzero temperature, the absorption of a photon can be accompanied not only by emission, but also by absorption of one or more phonons.

A simple derivation in Ref. [71] using a canonical transformation method gives the absorption coefficient of free polarons, which coincides with the result (2.2) of Ref. [69].

B. Optical absorption at weak coupling [within the canonical-transformation method [71] (DHL)]

The optical absorption of large polarons as a function of the frequency of the incident light is calculated using the canonical-transformation formalism by Devreese, Huybrechts and Lemmens (DHL) Ref. [71]. A simple calculation, which is developed below in full detail, gives a result for the absorption coefficient, which is exact to order α.

We start from the Hamiltonian of the electron-phonon system interacting with light is written down using the vector potential of an electromagnetic field $\mathbf{A}(t)$:

$$H = \frac{1}{2m_b}\left(\mathbf{p} + \frac{e}{c}\mathbf{A}(t)\right)^2 + \sum_{\mathbf{k}} \omega_{\mathrm{LO}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} + \sum_{\mathbf{k}} \left(V_{\mathbf{k}} a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_{\mathbf{k}}^{*} a_{\mathbf{k}}^{+} e^{-i\mathbf{k}\cdot\mathbf{r}}\right). \tag{2.3}$$

The electric field is related to the vector potential as

$$\mathbf{E}(t) = -\frac{1}{c}\frac{\partial \mathbf{A}(t)}{\partial t}. \tag{2.4}$$

Within the electric dipole interaction the electric field with frequency Ω is

$$\mathbf{E}(t) = \mathbf{E}\cos(\Omega t) \Rightarrow \tag{2.5}$$

$$\mathbf{A} = -\frac{c}{\Omega}\mathbf{E}\sin(\Omega t). \tag{2.6}$$

When expanding $\frac{1}{2m_b}\left(\mathbf{p} + \frac{e}{c}\mathbf{A}(t)\right)^2$ in the Hamiltonian, we find

$$\frac{1}{2m_b}\left(\mathbf{p} + \frac{e}{c}\mathbf{A}(t)\right)^2 = \frac{\mathbf{p}^2}{2m_b} + \frac{e}{m_b c}\mathbf{A}(t)\cdot\mathbf{p} + \frac{e^2}{2m_b c^2}A^2(t) \tag{2.7}$$

where the first term is the kinetic energy of the electron, and the second term describes the interaction of the electron-phonon system with light

$$V_t = \frac{e}{m_b c}\mathbf{A}\left(t\right)\cdot\mathbf{p} = -\frac{e}{m_b\Omega}\mathbf{E}\cdot\mathbf{p}\sin\Omega t \tag{2.8}$$

$$V_t \equiv V\sin\Omega t, \tag{2.9}$$

$$V \equiv -\frac{e}{m_b\Omega}\mathbf{E}\cdot\mathbf{p}. \tag{2.10}$$

Since $\mathbf{A}\left(t\right)$ does not depend on the electron coordinates, the term $\frac{e^2}{2m_b c^2}A^2\left(t\right)$ in (2.7) does not play a role in our description of the optical absorption. The total Hamiltonian for the system of a continuum polaron interacting with light is thus

$$H_{tot} = H_{pol} + V_t,$$

where H_{pol} is Fröhlich's Hamiltonian (1.16).

The absorption coefficient for absorption of light with frequency Ω by free polarons is proportional to the probability $P(\Omega)$ that a photon is absorbed by these polarons in their ground state,

$$\Gamma_p(\Omega) = \frac{N}{\varepsilon_0 cn2E^2}\Omega P(\Omega). \tag{2.11}$$

Here N is number of polarons, which are considered as independent from each other, ε_0 is the permittivity of vacuum, c is the velocity of light, n is the refractive index of the medium in which the polarons move, E is the modulus of the electric field vector of the incident photon. If the incident light can be treated as a perturbation, the transition probability $P(\Omega)$ is given by the Golden Rule of Fermi:

$$P(\Omega) = 2\pi\sum\nolimits_f \left\langle\Phi_0\left|V\right|f\right\rangle\left\langle f\left|V\right|\Phi_0\right\rangle\delta(E_0+\Omega-E_f). \tag{2.12}$$

V is the amplitude of the time-dependent perturbation given by (2.10). The ground state wave function of a free polaron is $\left|\Phi_0\right\rangle$ and its energy is E_0. The wave functions of all possible final states are $\left|f\right\rangle$ and the corresponding energies are E_f. The possible final states are all the excited states of the polaron. The main idea of the present calculation is to avoid the explicit summation over the final polaron states, which are poorly known, by eliminating all the excited state wave functions $\left|f\right\rangle$ from the expression (2.12).

With this aim, the representation of the δ-function is used:

$$\delta(x) = \frac{1}{\pi}\,\mathrm{Re}\int_{-\infty}^{0} dt \exp\left[-i\left(x+i\varepsilon\right)t\right].$$

This leads to:

$$\begin{aligned}P(\Omega) &= 2\,\mathrm{Re}\sum\nolimits_f \int_{-\infty}^{0} dt \left\langle \Phi_0 \left| V \right| f\right\rangle \left\langle f \left| V \right| \Phi_0\right\rangle \exp\left[-i(\Omega+i\varepsilon+E_0-E_f)t\right] \\ &= 2\,\mathrm{Re}\sum\nolimits_f \int_{-\infty}^{0} dt \exp\left[-i(\Omega+i\varepsilon)t\right] \left\langle \Phi_0 \left| V \right| f\right\rangle \left\langle f \left| e^{iHt} V e^{-iHt} \right| \Phi_0\right\rangle.\end{aligned}$$

Using the fact that

$$\sum\nolimits_f \left| f \right\rangle \left\langle f \right| = 1$$

and the notation

$$\begin{aligned} e^{iHt}V(0)e^{-iHt} &= V(t), \\ \frac{dV(t)}{dt} &= i\left[H, V(t)\right]\end{aligned}$$

we find

$$P(\Omega) = 2\,\mathrm{Re}\int_{-\infty}^{0} dt \exp\left[-i(\Omega+i\varepsilon)t\right] \left\langle \Phi_0 \left| V(0)V(t) \right| \Phi_0\right\rangle. \tag{2.13}$$

Defining

$$R(\Omega) = \int_{-\infty}^{0} dt \exp\left[-i(\Omega+i\varepsilon)t\right] \left\langle \Phi_0 \left| V(0)V(t) \right| \Phi_0\right\rangle, \tag{2.14}$$

one has

$$P(\Omega) = 2\,\mathrm{Re}\,R(\Omega). \tag{2.15}$$

Substituting (2.10) to (2.14), we find that

$$R\left(\Omega\right) = \left(\frac{e}{m_b\Omega}\right)^2 \int_{-\infty}^{0} dt e^{-i(\Omega+i\varepsilon)t} \left\langle \Phi_0 \left| \left(\mathbf{E}\cdot\mathbf{p}\left(0\right)\right)\left(\mathbf{E}\cdot\mathbf{p}\left(t\right)\right) \right| \Phi_0\right\rangle \tag{2.16}$$

and hence

$$P\left(\Omega\right) = 2\left(\frac{e}{m_b\Omega}\right)^2 \mathrm{Re}\int_{-\infty}^{0} dt e^{-i(\Omega+i\varepsilon)t} \left\langle \Phi_0 \left| \left(\mathbf{E}\cdot\mathbf{p}\left(0\right)\right)\left(\mathbf{E}\cdot\mathbf{p}\left(t\right)\right) \right| \Phi_0\right\rangle. \tag{2.17}$$

It is convenient to apply the first LLP transformation S_1(4.187), which eliminates the electron operators from the polaron Hamiltonian:

$$\begin{aligned} &H \longrightarrow \mathcal{H} = S_1^{-1} H_{pol} S_1 = \mathcal{H}_0 + \mathcal{H}_1 : \\ &\mathcal{H}_0 = \frac{\mathbf{P}^2}{2m_b} + \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b} - \frac{\mathbf{k}\cdot\mathbf{P}}{m_b}\right) a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} + V_k^* a_{\mathbf{k}}^{\dagger}), \\ &\mathcal{H}_1 = \frac{1}{2m_b}\sum_{\mathbf{k}} \mathbf{k}\cdot\mathbf{k}' a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}'}^{\dagger} a_{\mathbf{k}} a_{\mathbf{k}'},\end{aligned}$$

where $\mathcal{H}_0$ can be diagonalized exactly and gives rise to the self-energy $E = -\alpha\omega_{\mathrm{LO}}$,and $\mathcal{H}_1$ contains the correlation effects between the phonons. The optical absorption will be calculated here for the total momentum of the system $\mathbf{P} = \mathbf{0}$.

In the LLP approximation the explicit form of the matrix element in (2.16) is

$$\left\langle \Phi_0 \left| \left(\mathbf{E} \cdot \mathbf{p}\left(0\right) \right) \left(\mathbf{E} \cdot \mathbf{p}\left(t\right) \right) \right| \Phi_0 \right\rangle = \left\langle 0 \left| S_2^{-1} S_1^{-1} \mathbf{E} \cdot \mathbf{p} S_1 S_1^{-1} \mathbf{E} \cdot \mathbf{p}(t) S_1 S_2 \right| 0 \right\rangle, \tag{2.18}$$

where $S_1$and S_2 are the first (4.187) and the second (4.196) LLP transformations. The application of S_1 gives:

$$S_1^{-1} \mathbf{p}(t) S_1 = S_1^{-1} e^{iHt} \mathbf{p} e^{-iHt} S_1 = S_1^{-1} e^{iHt} S_1 S_1^{-1} \mathbf{p} S_1 S_1^{-1} e^{-iHt} S_1.$$

Using $\mathcal{H} = S_1^{-1} H S_1$, we arrive at $S_1^{-1} e^{iHt} S_1 = e^{i\mathcal{H}t}$.Further we recall $S_1^{-1} \mathbf{p} S_1 = \mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \mathbf{p}$, where $\mathbf{P} = \mathbf{0}$ and $\mathbf{p}$ is set 0 (see Appendix 1). This results in

$$S_1^{-1} \mathbf{p}(t) S_1 = e^{i\mathcal{H}t} \mathbf{p} e^{-i\mathcal{H}t} = -\sum_{\mathbf{k}} \hbar \mathbf{k} e^{i\mathcal{H}t} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} e^{-i\mathcal{H}t} = -\sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger}(t) a_{\mathbf{k}}(t).$$

Then (2.18) takes the form

$$\left\langle \Phi_0 \left| \left(\mathbf{E} \cdot \mathbf{p}\left(0\right) \right) \left(\mathbf{E} \cdot \mathbf{p}\left(t\right) \right) \right| \Phi_0 \right\rangle = \left\langle 0 \left| S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger}(t) a_{\mathbf{k}}(t) \right) S_2 \right| 0 \right\rangle. \tag{2.19}$$

Here the second LLP transformation is given by (4.196) with

$$f_k = -\frac{V_k^*}{\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}} \tag{2.20}$$

and the vacuum is defined by $a_{\mathbf{k}} \left| 0 \right\rangle = 0$. The calculation of the matrix element (2.19) proceeds as follows:

$$\begin{aligned} & \left\langle 0 \left| S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger}(t) a_{\mathbf{k}}(t) \right) S_2 \right| 0 \right\rangle \\ & = \left\langle 0 \left| S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) S_2 S_2^{-1} e^{i\mathcal{H}t} S_2 S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) S_2 S_2^{-1} e^{-i\mathcal{H}t} S_2 \right| 0 \right\rangle \\ & = \left\langle 0 \left| S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) S_2 e^{i S_2^{-1} \mathcal{H} S_2 t} S_2^{-1} \left(\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right) S_2 e^{-i S_2^{-1} \mathcal{H} S_2 t} \right| 0 \right\rangle. \end{aligned} \tag{2.21}$$

Further on, we calculate

$$S_2^{-1} \mathcal{H} S_2 = H_0 + H_1$$

where

$$H_0 = S_2^{-1}\mathcal{H}_0 S_2 = S_2^{-1}\left[\sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \sum_{\mathbf{k}}(V_k a_{\mathbf{k}} + V_k^* a_{\mathbf{k}}^{\dagger})\right] S_2$$

Further we use $S_2^{-1}a_{\mathbf{k}}S_2 = a_{\mathbf{k}} + f_{\mathbf{k}}$:

$$\begin{aligned}
H_0 &= \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)|f_{\mathbf{k}}|^2 \\
&\quad + \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)\left(a_{\mathbf{k}}^{\dagger}f_{\mathbf{k}} + a_{\mathbf{k}}f_{\mathbf{k}}^*\right) + \sum_{\mathbf{k}}\left[V_k\left(a_{\mathbf{k}} + f_{\mathbf{k}}\right) + V_k^*\left(a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^*\right)\right] \\
&= \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \sum_{\mathbf{k}}\frac{|V_k|^2}{\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)} - 2\sum_{\mathbf{k}}\frac{|V_k|^2}{\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)} \\
&= \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} - \sum_{\mathbf{k}}\frac{|V_k|^2}{\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)}.
\end{aligned}$$

The last term can be calculated analytically:

$$\begin{aligned}
\sum_{\mathbf{k}}\frac{|V_k|^2}{\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}} &= \frac{V}{(2\pi)^3}\int d^3k\left(\frac{\omega_{\mathrm{LO}}}{k}\right)^2\frac{4\pi\alpha}{V}\left(\frac{1}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\cdot\frac{1}{\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}} \\
&= \frac{\alpha\omega_{\mathrm{LO}}}{2\pi^2}4\pi\int_0^{\infty}dk\left(\frac{1}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\frac{1}{1 + \frac{k^2}{2m_b\omega_{\mathrm{LO}}}} \\
&= \frac{2\alpha\omega_{\mathrm{LO}}}{\pi}\int_0^{\infty}d\kappa\frac{1}{1+\varkappa^2} = \frac{2\alpha\omega_{\mathrm{LO}}}{\pi}\arctan\varkappa\Big|_0^{\infty} = \alpha\omega_{\mathrm{LO}},
\end{aligned}$$

$$H_0 = -\alpha\omega_{\mathrm{LO}} + \sum_{\mathbf{k}}\left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b}\right)a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}.$$

The term

$$H_1 = S_2^{-1}\mathcal{H}_1 S_2$$

will be neglected:

$$e^{iS_2^{-1}\mathcal{H}S_2 t} \approx e^{iH_0 t}.$$

Neglecting H_1, consistent with the LLP description, introduces no error in order α. Therefore (2.21) becomes

$$\left\langle 0 \left| \begin{array}{c} \sum_{\mathbf{k}}\mathbf{E}\cdot\mathbf{k}\left(a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + f_{\mathbf{k}}a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^*a_{\mathbf{k}} + f_{\mathbf{k}}f_{\mathbf{k}}^*\right)e^{iH_0t} \\ \times\sum_{\mathbf{k}}\mathbf{E}\cdot\mathbf{k}\left(a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + f_{\mathbf{k}}a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^*a_{\mathbf{k}} + f_{\mathbf{k}}f_{\mathbf{k}}^*\right)e^{-iH_0t} \end{array} \right| 0 \right\rangle. \qquad (2.22)$$

For $\mathbf{P} = 0$ there is no privileged direction and $\sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} f_{\mathbf{k}} f_{\mathbf{k}}^* = 0$, (2.22) reduces to:

$$\left\langle 0 \left| \sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} f_{\mathbf{k}}^* a_{\mathbf{k}} e^{iH_0 t} \sum_{\mathbf{k}} \mathbf{E} \cdot \mathbf{k} f_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} e^{-iH_0 t} \right| 0 \right\rangle .$$

From the equation of motion for $a_{\mathbf{k}}^{\dagger}$:

$$\frac{da_{\mathbf{k}}^{\dagger}(t)}{dt} = i \left[H_0, a_{\mathbf{k}}^{\dagger} \right] = i \left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b} \right) a_{\mathbf{k}}^{\dagger},$$

it is easy now to calculate

$$e^{iH_0 t} a_{\mathbf{k}}^{\dagger} e^{-iH_0 t} = a_{\mathbf{k}}^{\dagger} \exp \left[i \left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b} \right) t \right] .$$

The matrix element (2.19) now becomes

$$\langle \Phi_0 \left| (\mathbf{E} \cdot \mathbf{p}(0)) (\mathbf{E} \cdot \mathbf{p}(t)) \right| \Phi_0 \rangle = \sum_{\mathbf{k}} (\mathbf{E} \cdot \mathbf{k})^2 f_{\mathbf{k}}^* f_{\mathbf{k}} \exp \left[i \left(\omega_{\mathrm{LO}} + \frac{k^2}{2m_b} \right) t \right] + O(\alpha^2).$$

The transition probability (2.13) is then given by the expression

$$\begin{aligned} P(\Omega) &= 2 \operatorname{Re} \frac{e^2}{m_b^2 \Omega^2} \sum_{\mathbf{k}} (\mathbf{E} \cdot \mathbf{k})^2 f_{\mathbf{k}}^* f_{\mathbf{k}} \int_{-\infty}^{0} dt \exp \left[-i \left(\Omega + i\varepsilon - \omega_{\mathrm{LO}} - \frac{k^2}{2m_b} \right) t \right] \\ &= 2\pi \frac{e^2}{m_b^2 \Omega^2} \sum_{\mathbf{k}} (\mathbf{E} \cdot \mathbf{k})^2 \left| f_{\mathbf{k}} \right|^2 \delta \left(\Omega - \omega_{\mathrm{LO}} - \frac{k^2}{2m_b} \right) . \end{aligned}$$

Using (2.20), we obtain

$$
\begin{aligned}
P(\Omega) &= \frac{2\pi e^2}{m_b^2\Omega^2}\sum_{\mathbf{k}}\frac{(\mathbf{E}\cdot\mathbf{k})^2}{\left(\omega_{\mathrm{LO}}+\frac{k^2}{2m_b}\right)^2}\left|V_k\right|^2\delta\left(\Omega-\omega_{\mathrm{LO}}-\frac{k^2}{2m_b}\right)\\
&= \frac{2\pi e^2}{m_b^2\Omega^2}\frac{V}{(2\pi)^3}\int d^3k\left(\frac{\omega_{\mathrm{LO}}}{k}\right)^2\frac{4\pi\alpha}{V}\left(\frac{1}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\\
&\times\frac{(\mathbf{E}\cdot\mathbf{k})^2}{\left(\omega_{\mathrm{LO}}+\frac{k^2}{2m_b}\right)^2}\delta\left(\Omega-\omega_{\mathrm{LO}}-\frac{k^2}{2m_b}\right)\\
&= \frac{e^2\alpha E^2}{m_b^2\Omega^2\pi}2\pi\int_{-1}^{1}dx x^2\int_0^\infty dk\left(\frac{1}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\\
&\times\frac{k^2}{\left(1+\frac{k^2}{2m_b\omega_{\mathrm{LO}}}\right)^2}\frac{1}{\omega_{\mathrm{LO}}}\delta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1-\frac{k^2}{2m_b\omega_{\mathrm{LO}}}\right)\\
&= \frac{8e^2\alpha E^2}{3m_b\Omega^2}\int_0^\infty d\kappa\frac{\varkappa^2}{(1+\varkappa^2)^2}\delta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1-\varkappa^2\right)\\
&= \frac{4e^2\alpha E^2}{3m_b\Omega^2}\int_0^\infty d\zeta\frac{\sqrt{\zeta}}{(1+\zeta)^2}\delta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1-\zeta\right)\\
&= \frac{4e^2\alpha E^2}{3m_b\Omega^2}\Theta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1\right)\frac{\sqrt{\frac{\Omega}{\omega_{\mathrm{LO}}}-1}}{\left(\frac{\Omega}{\omega_{\mathrm{LO}}}\right)^2}=\frac{4e^2\alpha E^2\omega_{\mathrm{LO}}^2}{3m_b\Omega^4}\sqrt{\frac{\Omega}{\omega_{\mathrm{LO}}}-1}\quad\Theta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1\right),
\end{aligned}
$$

where

$$
\Theta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1\right)=\begin{cases}1 \text{ if} & \frac{\Omega}{\omega_{\mathrm{LO}}}>1\\ 0 \text{ if} & \frac{\Omega}{\omega_{\mathrm{LO}}}<1\end{cases}.
$$

The absorption coefficient (2.11) for absorption by free polarons for $\alpha\longrightarrow 0$ finally takes the form

$$
\Gamma_p(\Omega)=\frac{1}{\varepsilon_0 cn}\frac{2Ne^2\alpha\omega_{\mathrm{LO}}^2}{3m_b\Omega^3}\sqrt{\frac{\Omega}{\omega_{\mathrm{LO}}}-1}\quad\Theta\left(\frac{\Omega}{\omega_{\mathrm{LO}}}-1\right). \qquad (2.23)
$$

The behaviuor of $\Gamma_p(\Omega)$ (2.23) as a function of Ω is as follows. For $\Omega<\omega_{\mathrm{LO}}$ there is no absorption. The threshold for absorption is at $\Omega=\omega_{\mathrm{LO}}$.From $\Omega=\omega_{\mathrm{LO}}$ up to $\Omega=\frac{6}{5}\omega_{\mathrm{LO}},\Gamma_p$ increases to a maximum and for $\Omega>\frac{6}{5}\omega_{\mathrm{LO}}$ the absorption coefficient decreases slowly with increasing Ω.

Experimentally, this one-phonon line has been observed for free polarons in the infrared absorption spectra of CdO-films, see Fig. 6. In CdO, which is a weakly polar material with $\alpha\approx 0.74$, the polaron absorption band is observed in the spectral region between 6 and 20 μm (above the LO phonon frequency). The difference between theory and experiment in the wavelength region where polaron absorption dominates the spectrum is due to many-polaron effects.



FIG. 6: Optical absorption spectrum of a CdO-film with the carrier concentration $n = 5.9 \times 10^{19}$ cm^{-3} at $T = 300$ K. The experimental data (solid dots) of Ref. [72] are compared to different theoretical results: with (solid curve) and without (dashed line) the one-polaron contribution of Ref. [69] and for many polarons (dash-dotted curve) of Ref. [51].

III. OPTICAL ABSORPTION. STRONG COUPLING

The absorption of light by free large polarons was treated in Ref. [47] using the polaron states obtained wihtin the adiabatic strong-coupling approximation, which was considered above in subsection I E 3.

It was argued in Ref. [47], that for sufficiently large α ($\alpha > 3$), the (first) RES of a polaron is a relatively stable state, which can participate in optical absorption transitions. This idea was necessary to understand the polaron optical absorption spectrum in the strong-coupling regime. The following scenario of a transition, which leads to a *"zero-phonon" peak* in the absorption by a strong-coupling polaron, can then be suggested. If the frequency of the incoming photon is equal to

$$\Omega_{\text{RES}} \equiv \frac{\text{E}_{\text{RES}}\text{-}E_0}{\hbar} = 0.065\alpha^2\omega_{\text{LO}},$$

then the electron jumps from the ground state (which, at large coupling, is well-characterized by "s"-symmetry for the electron) to an excited state ("$2p$"), while the lattice polarization in the final state is adapted to the "$2p$" electronic state of the polaron. In Ref. [47] considering the decay of the RES with emission of one real phonon it is demonstrated, that the "zero-phonon" peak can be described using the Wigner-Weisskopf formula valid when the linewidth of that peak is much smaller than $\hbar\omega_{\mathrm{LO}}$.

For photon energies larger than

$$\Omega_{\mathrm{RES}} + \omega_{\mathrm{LO}},$$

a transition of the polaron towards the first scattering state, belonging to the RES, becomes possible. The final state of the optical absorption process then consists of a polaron in its lowest RES plus a free phonon. A "one-phonon sideband" then appears in the polaron absorption spectrum. This process is called *one-phonon sideband absorption.*

The one-, two-, ... K-, ... phonon sidebands of the zero-phonon peak give rise to a broad structure in the absorption spectrum. It turns that the *first moment* of the phonon sidebands corresponds to the FC frequency Ω_{FC}:

$$\Omega_{\mathrm{FC}} \equiv \frac{E_{\mathrm{FC}} - E_0}{\hbar} = 0.141\alpha^2\omega_{\mathrm{LO}}.$$

To summarize, the polaron optical absorption spectrum at strong coupling is characterized by the following features ($T = 0$):

a) An intense absorption peak ("zero-phonon line") appears, which corresponds to a transition from the ground state to the first RES at Ω_{RES}.

b) For $\Omega > \Omega_{\mathrm{RES}} + \omega_{\mathrm{LO}}$, a phonon sideband structure arises. This sideband structure peaks around Ω_{FC}.

The qualitative behaviour predicted in Ref. [47], namely, an intense zero-phonon (RES) line with a broader sideband at the high-frequency side, was confirmed after an all-coupling expression for the polaron optical absorption coefficient at $\alpha = 5, 6, 7$ had been studied [48].

In what precedes, the low-frequency end of the polaron absorption spectrum was discussed; at higher frequencies, transitions to higher RES and their scattering states can appear. The two-phonon sidebands in the optical absorption of free polarons in the strong-coupling limit were numerically studied in Ref. [73].

The study of the optical absorption of polarons at large coupling is mainly of formal interest because all reported coupling constants of polar semiconductors and ionic crystals are smaller than 5 (see Table 1).

IV. ARBITRARY COUPLING

A. Impedance function of large polarons: An alternative derivation of FHIP [74]

a. Definitions We derive here the linear response of the Fröhlich polaron, described by the Hamiltonian

$$H = \frac{\mathbf{p}^2}{2m_b} + \sum_{\mathbf{k}} \hbar\omega_{\mathbf{k}} a^+_{\mathbf{k}} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^\dagger_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}}), \tag{4.1}$$

to a spatially uniform, time-varying electric field

$$\mathbf{E}_\Omega(t) = E_0 \exp(i\Omega t)\, \mathbf{e}_x. \tag{4.2}$$

This field induces a current in the x-direction

$$j_\Omega(t) = \frac{1}{Z(\Omega)} E_\Omega(t). \tag{4.3}$$

The complex function $Z(\Omega)$ is called the impedance function. The frequency-dependent mobility is defined by

$$\mu(\Omega) = \mathrm{Re}\,\frac{1}{Z(\Omega)}. \tag{4.4}$$

For nonzero frequencies (in the case of polarons the frequencies of interest are in the infrared) one defines the absorption coefficient [48]

$$\Gamma(\Omega) = \frac{1}{n\epsilon_0 c}\,\mathrm{Re}\,\frac{1}{Z(\Omega)}, \tag{4.5}$$

where ϵ_0 is the dielectric constant of the vacuum, n the refractive index of the crystal, and c the velocity of light. In the following the amplitude of the electric field E_0 is taken sufficiently small so that linear-response theory can be applied.

The impedance function can be expressed via a frequency-dependent conductivity of a single polaron in a unit volume

$$\frac{1}{Z(\Omega)} = \sigma(\Omega) \tag{4.6}$$

using *the standard Kubo formula* (cf. Eq. (3.8.8) from Ref. [70]):

$$\sigma(\Omega) = i\frac{e^2}{Vm_b\Omega} + \frac{1}{V\hbar\Omega}\int_0^\infty e^{i\Omega t}\left\langle [j_x(t), j_x]\right\rangle dt. \tag{4.7}$$

In order to introduce a convenient representation of the impedance function, we give in the next subsection a definition and discuss properties of a scalar product of two operators [cf. [75], Chapter 5].

b. Definition and properties of the scalar product For two operators A and B (i.e., elements of the Hilbert space of operators) the scalar product is defined as

$$(A, B) = \int_0^\beta d\lambda \left\langle \left(e^{\lambda\hbar L} A^\dagger\right) B\right\rangle. \tag{4.8}$$

The notation $\left(e^{\lambda\hbar L}A^\dagger\right)$ is used in order to indicate that the operator $e^{\lambda\hbar L}$ acts on the operator $A^\dagger$. The time evolution of the operator A is determined by the Liouville operator L:

$$-i\frac{\partial A}{\partial t} = LA \equiv \frac{1}{\hbar}\left[H, A\right] \tag{4.9}$$

with a commutator $[H, A]$, wherefrom

$$A(t) = e^{iLt}A(0) \equiv e^{iHt/\hbar}A(0)e^{-iHt/\hbar}. \tag{4.10}$$

The expectation value in (4.8) is taken over the Gibbs' ensemble:

$$\langle A\rangle = \mathrm{Tr}(\rho_0(H)A) \tag{4.11}$$

with the equilibtium density matrix when the electric field is absent

$$\rho_0(H) = e^{-\beta H}/\mathrm{Tr}(e^{-\beta H}). \tag{4.12}$$

One can show that (4.8) defines a positive definite scalar product with the following properties

$$\text{(i)} \qquad (A, B) = (B^\dagger, A^\dagger), \tag{4.13}$$

$$\text{(ii)} \qquad (A, LB) = (LA, B), \tag{4.14}$$

$$\text{(iii)} \qquad (A, LB) = \frac{1}{\hbar}\left\langle \left[A^\dagger, B\right]\right\rangle, \tag{4.15}$$

and [cf. Eq. (5.11) of [75]]

$$\text{(iv)} \qquad (A, B)^* = (B, A) \tag{4.16}$$

Demonstration of the property (4.13). Starting from the definition (4.8) and using (4.10), we obtain

$$(A,B) = \int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} B \right\rangle . \tag{4.17}$$

Substituting here (4.11) with (4.12), one finds

$$(A,B) = \int_0^\beta d\lambda \mathrm{Tr}\left[e^{-(\beta-\lambda)H} A^\dagger e^{-\lambda H} B \right] / \mathrm{Tr}(e^{-\beta H}).$$

Change of the variable $\lambda' = \beta - \lambda$ allows us to represnt this integral as

$$(A,B) = \int_0^\beta d\lambda' \mathrm{Tr}\left[e^{-\lambda' H} A^\dagger e^{-(\beta-\lambda')H} B \right] / \mathrm{Tr}(e^{-\beta H}).$$

Further, a cyclic permutation of the operators under the trace Tr sign gives

$$\begin{aligned}(A,B) &= \int_0^\beta d\lambda \mathrm{Tr}\left[e^{-(\beta-\lambda)H} B e^{-\lambda H} A^\dagger \right] / \mathrm{Tr}(e^{-\beta H}) \\ &= \int_0^\beta d\lambda \left\langle e^{\lambda H} B e^{-\lambda H} A^\dagger \right\rangle = \int_0^\beta d\lambda \left\langle \left(e^{\lambda \hbar L} B \right) A^\dagger \right\rangle \\ &= \int_0^\beta d\lambda \left\langle \left[e^{\lambda \hbar L} \left(B^\dagger \right)^\dagger \right] A^\dagger \right\rangle .\end{aligned}$$

According to the definition (4.8), this finalizes the demonstration of (4.13).

Demonstration of the property (4.14). Starting from (4.17) and using (4.9), we obtain

$$\begin{aligned}(A, LB) &= \int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} LB \right\rangle \\ &= \frac{1}{\hbar}\int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} \left(HB - BH \right) \right\rangle .\end{aligned}$$

A cyclic permutation of the operators under the average $\langle \bullet \rangle$ sign gives

$$(A, LB) = \frac{1}{\hbar}\int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} HB - H e^{\lambda H} A^\dagger e^{-\lambda H} B \right\rangle . \tag{4.18}$$

Using the commutation of H and $e^{\pm \lambda H}$, one finds

$$(A, LB) = \frac{1}{\hbar}\int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger H e^{-\lambda H} B - e^{\lambda H} H A^\dagger e^{-\lambda H} B \right\rangle \tag{4.19}$$

$$= \frac{1}{\hbar}\int_0^\beta d\lambda \left\langle e^{\lambda H} \left(A^\dagger H - H A^\dagger \right) e^{-\lambda H} B \right\rangle \tag{4.20}$$

$$= \frac{1}{\hbar}\int_0^\beta d\lambda \left\langle e^{\lambda H} \left(HA - AH \right)^\dagger e^{-\lambda H} B \right\rangle \tag{4.21}$$

$$= \int_0^\beta d\lambda \left\langle e^{\lambda H} \left(\frac{1}{\hbar}[H, A] \right)^\dagger e^{-\lambda H} B \right\rangle . \tag{4.22}$$

With the definition (4.9), this gives

$$(A, LB) = \int_0^\beta d\lambda \left\langle e^{\lambda H} (LA)^\dagger e^{-\lambda H} B \right\rangle = . \int_0^\beta d\lambda \left\langle \left[e^{\lambda\hbar L} (LA)^\dagger \right] B \right\rangle .$$

According to the definition (4.8), this finalizes the demonstration of (4.14).

Demonstration of the property (4.15). Starting from (4.20) and performing a cyclic permutation of the operators under the average $\langle \bullet \rangle$, we find

$$(A, LB) = \frac{1}{\hbar} \int_0^\beta d\lambda \left\langle e^{\lambda H} \left(A^\dagger H - H A^\dagger \right) e^{-\lambda H} B \right\rangle$$

Further we notice that

$$e^{\lambda H} \left(A^\dagger H - H A^\dagger \right) e^{-\lambda H} = - \frac{d \left(e^{\lambda H} A^\dagger e^{-\lambda H} \right)}{d\lambda},$$

consequently,

$$\begin{aligned}
(A, LB) &= -\frac{1}{\hbar} \int_0^\beta d\lambda \left\langle \frac{d \left(e^{\lambda H} A^\dagger e^{-\lambda H} \right)}{d\lambda} B \right\rangle \\
&= -\frac{1}{\hbar} \left\langle \int_0^\beta d\lambda \frac{d \left(e^{\lambda H} A^\dagger e^{-\lambda H} \right)}{d\lambda} B \right\rangle \\
&= -\frac{1}{\hbar} \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} B \Big|_0^\beta \right\rangle = \frac{1}{\hbar} \left\langle \left(A^\dagger B - e^{\beta H} A^\dagger e^{-\beta H} B \right) \right\rangle \\
&= \frac{1}{\hbar} \mathrm{Tr} \left[e^{-\beta H} \left(A^\dagger B - e^{\beta H} A^\dagger e^{-\beta H} B \right) \right] / \mathrm{Tr}(e^{-\beta H}) \\
&= \frac{1}{\hbar} \mathrm{Tr} \left[e^{-\beta H} A^\dagger B - A^\dagger e^{-\beta H} B \right] / \mathrm{Tr}(e^{-\beta H}).
\end{aligned} \tag{4.23}$$

Further, a cyclic permutation of the operators in the second term under the trace Tr sign gives

$$\begin{aligned}
(A, LB) &= \frac{1}{\hbar} \mathrm{Tr} \left[e^{-\beta H} A^\dagger B - e^{-\beta H} B A^\dagger \right] / \mathrm{Tr}(e^{-\beta H}) \\
&= \frac{1}{\hbar} \mathrm{Tr} \left[e^{-\beta H} \left(A^\dagger B - B A^\dagger \right) \right] / \mathrm{Tr}(e^{-\beta H}) \\
&= \frac{1}{\hbar} \left\langle A^\dagger B - B A^\dagger \right\rangle = \frac{1}{\hbar} \left\langle \left[A^\dagger, B \right] \right\rangle .
\end{aligned}$$

Thus, the property (4.15) has been demonstrated.

Demonstration of the property (4.16). We start from the represntation of the scalar product (4.17) and take a complex conjugate:

$$(A,B)^* = \int_0^\beta d\lambda \left\langle e^{\lambda H} A^\dagger e^{-\lambda H} B\right\rangle^* = \int_0^\beta d\lambda \left\langle B^\dagger e^{-\lambda H} A e^{\lambda H}\right\rangle .$$

A cyclic permutation of the operators under the average $\langle\bullet\rangle$ sign gives then

$$\begin{aligned}(A,B)^* &= \int_0^\beta d\lambda \left\langle e^{\lambda H} B^\dagger e^{-\lambda H} A\right\rangle = \int_0^\beta d\lambda \left\langle e^{\lambda H} B^\dagger e^{-\lambda H} A\right\rangle \\ &= \int_0^\beta d\lambda \left\langle \left(e^{\lambda\hbar L} B^\dagger\right) A\right\rangle . \qquad (4.24)\end{aligned}$$

According to the definition (4.8), this finalizes the demonstration of (4.16).

The above scalar product allows one to represent different dynamical quantities in a rather simple way. For example, let us consider a scalar product

$$\begin{aligned}\Phi_{AB}(z) &= \left(A, \frac{1}{z-L}B\right) \qquad (4.25)\\ &= \int_0^\beta d\lambda \left\langle \left(e^{\lambda\hbar L}A^\dagger\right)\frac{1}{z-L}B\right\rangle \\ &= -i\int_0^\beta d\lambda \left\langle e^{\lambda\hbar L}A^\dagger \left[\int_0^\infty dt e^{i(z-L)t}\right] B\right\rangle \qquad (4.26)\\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle e^{\lambda\hbar L}A^\dagger e^{-iLt}B\right\rangle \\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle e^{\lambda H}A^\dagger e^{-\lambda H} e^{-iHt/\hbar} B e^{iHt/\hbar}\right\rangle \\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle e^{iHt/\hbar+\lambda H}A^\dagger e^{-iHt/\hbar-\lambda H} B\right\rangle \\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle e^{iH(t-i\lambda\hbar)/\hbar}A^\dagger e^{-iH(t-i\lambda\hbar)/\hbar} B\right\rangle \\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle e^{iL(t-i\lambda\hbar)}A^\dagger B\right\rangle \qquad (4.27)\\ &= -i\int_0^\infty dt e^{izt}\int_0^\beta d\lambda \left\langle A^\dagger(t-i\hbar\lambda)B(0)\right\rangle . \qquad (4.28)\end{aligned}$$

c. Representation of the impedance function in terms of the relaxation function The impedance function is related to the relaxation function

$$\Phi(z) \equiv \Phi_{\dot{x}\dot{x}}(z) = \left(\dot{x}, \frac{1}{z-L}\dot{x}\right), \qquad (4.29)$$

where $\dot{x}$ is the velocity operator, by the following expression:

$$\frac{1}{Z(\Omega)} = ie^2 \lim_{\epsilon\to 0}\Phi(\Omega+i\epsilon) \qquad (4.30)$$

$(z=\Omega+i\epsilon, \epsilon>0)$.

Demonstration of the representation (4.29). Apply (4.27) to the relaxation function entering (4.30):

$$\Phi(z) = -i \int_0^\beta d\lambda \int_0^\infty dt e^{izt} \left\langle \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle$$

and perform the integration by parts using the formula

$$\begin{aligned} \int_0^\infty e^{izt} f(t)\, dt &= -\frac{i}{z} f(t) e^{izt} \bigg|_{t=0}^{\infty} + \frac{i}{z} \int_0^\infty \frac{\partial f(t)}{\partial t} e^{izt} dt \\ &= \frac{i}{z} f(0) + \frac{i}{z} \int_0^\infty \frac{\partial f(t)}{\partial t} e^{izt} dt : \end{aligned}$$

$$\begin{aligned} \int_0^\infty e^{izt} \left\langle \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt &= \frac{i}{z} \left\langle \left(e^{\hbar\lambda L} \dot{x} \right) \dot{x} \right\rangle + \frac{i}{z} \int_0^\infty e^{izt} \frac{\partial}{\partial t} \left\langle \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt \\ &= \frac{i}{z} \left\langle \left(e^{\hbar\lambda L} \dot{x} \right) \dot{x} \right\rangle - \frac{1}{z} \int_0^\infty e^{izt} \left\langle L \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt. \end{aligned}$$

This allows us to represent the relaxation function in the form

$$\begin{aligned} \Phi(z) &= -i \int_0^\beta d\lambda \left(\frac{i}{z} \left\langle \left(e^{\hbar\lambda L} \dot{x} \right) \dot{x} \right\rangle - \frac{1}{z} \int_0^\infty e^{izt} \left\langle L \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt \right) \\ &= \frac{1}{z} \int_0^\beta d\lambda \left\langle \left(e^{\hbar\lambda L} \dot{x} \right) \dot{x} \right\rangle + \frac{i}{z} \int_0^\beta d\lambda \int_0^\infty e^{izt} \left\langle L \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt \\ &= \frac{1}{m_b z} + \frac{i}{z} \int_0^\beta d\lambda \int_0^\infty e^{izt} \left\langle L \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle dt, \end{aligned}$$

where the expression (4.55) is inserted in the first term. Further on, the integral over λ is taken as follows:

$$\begin{aligned} \int_0^\beta d\lambda \left\langle L \left(e^{i(t-i\hbar\lambda)L} \dot{x} \right) \dot{x} \right\rangle &= \left\langle \left(L \int_0^\beta e^{i(t-i\hbar\lambda)L} d\lambda \dot{x} \right) \dot{x} \right\rangle \\ &= \frac{1}{\hbar} \left\langle \left(e^{iLt} \left(e^{\hbar\beta L} - 1 \right) \dot{x} \right) \dot{x} \right\rangle \\ &= \frac{1}{\hbar} \left\langle e^{iLt} \left(e^{\hbar\beta L} \dot{x} - \dot{x} \right) \dot{x} \right\rangle \\ &= \frac{1}{\hbar} \left\langle \left(e^{\hbar\beta L} \dot{x}(t) - \dot{x}(t) \right) \dot{x} \right\rangle \\ &= \frac{1}{\hbar} \left\langle \left(e^{\beta H} \dot{x}(t) e^{-\beta H} - \dot{x}(t) \right) \dot{x} \right\rangle \\ &= \frac{1}{\hbar \mathrm{Tr} e^{-\beta H}} \mathrm{Tr} \left[e^{-\beta H} \left(e^{\beta H} \dot{x}(t) e^{-\beta H} - \dot{x}(t) \right) \dot{x} \right] \\ &= \frac{1}{\hbar \mathrm{Tr} e^{-\beta H}} \mathrm{Tr} \left[\left(\dot{x}(t) e^{-\beta H} - e^{-\beta H} \dot{x}(t) \right) \dot{x} \right] \\ &= \frac{1}{\hbar \mathrm{Tr} e^{-\beta H}} \mathrm{Tr} \left[e^{-\beta H} \left(\dot{x}\dot{x}(t) - \dot{x}(t) \right) \dot{x} \right] \\ &= \frac{1}{\hbar} \left\langle \dot{x}\dot{x}(t) - \dot{x}(t)\dot{x} \right\rangle = -\frac{1}{\hbar} \left\langle \dot{x}(t), \dot{x} \right\rangle. \end{aligned}$$

Hence,

$$\Phi\left(z\right)=\frac{1}{m_bz}-\frac{i}{\hbar z}\int_0^\infty dte^{izt}\left\langle\left[\dot{x}\left(t\right),\dot{x}\right]\right\rangle. \tag{4.31}$$

When setting $z=\Omega+i\varepsilon$ with $\varepsilon\to+0$, we have

$$\Phi\left(\Omega+i\varepsilon\right)=\frac{1}{m_b}\frac{1}{\Omega+i\varepsilon}-\frac{i}{\hbar\left(\Omega+i\varepsilon\right)}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[\dot{x}\left(t\right),\dot{x}\right]\right\rangle dt. \tag{4.32}$$

For $\Omega\neq 0$, we can set

$$\Phi\left(\Omega+i\varepsilon\right)=\frac{1}{m_b}\frac{1}{\Omega}-\frac{i}{\hbar\Omega}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[\dot{x}\left(t\right),\dot{x}\right]\right\rangle dt. \tag{4.33}$$

Multiplying $\Phi\left(\Omega+i\varepsilon\right)$ by ie^2, we find that

$$\begin{aligned}ie^2\Phi\left(\Omega+i\varepsilon\right)&=ie^2\left(\frac{1}{m_b}\frac{1}{\Omega}-\frac{i}{\hbar\Omega}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[\dot{x}\left(t\right),\dot{x}\right]\right\rangle dt\right)\\&=i\frac{e^2}{m_b\Omega}+\frac{e^2}{\hbar\Omega}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[\dot{x}\left(t\right),\dot{x}\right]\right\rangle dt\\&=i\frac{e^2}{m_b\Omega}+\frac{1}{\hbar\Omega}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[j_x\left(t\right),j_x\right]\right\rangle dt,\end{aligned} \tag{4.34}$$

where the electric current density is

$$j_x=-e\dot{x}.$$

Substituting further (4.34) in (4.30), we arrive at

$$\frac{1}{Z(\Omega)}=i\frac{e^2}{m_b\Omega}+\lim_{\epsilon\to0}\frac{1}{\hbar\Omega}\int_0^\infty e^{i(\Omega+i\varepsilon)t}\left\langle\left[j_x\left(t\right),j_x\right]\right\rangle dt,$$

what coincides with the expression of the impedance function (4.6) through a frequency-dependent conductivity given by the Kubo formula (4.7), q.e.d.

d. Application of the projection operator technique Using the Mori-Zwanzig projection operator technique (cf. [75], Chapter 5), the relaxation function (4.29)

$$\Phi\left(z\right)=\left(\dot{x},\frac{1}{z-L}\dot{x}\right)$$

can be represented in a form, which is especially convenient for the application in the theory of the optical absorption of polarons.

The projection operator P $(Q=1-P)$ is defined as

$$PA=\frac{\dot{x}\left(\dot{x},A\right)}{\chi} \tag{4.35}$$

with A an operator and

$$\chi=\left(\dot{x},\dot{x}\right). \tag{4.36}$$

The projection operator $Q = 1 - P$ projects an operator onto the space orthogonal to the space containing $\dot{x}$. Here we give some examples of the action of the projection operators:

$$P\dot{x} = \dot{x}, \qquad Q\dot{x} = (1-P)\dot{x} = 0; \tag{4.37}$$

$$Px = \frac{\dot{x}\left(\dot{x}, x\right)}{\chi} = \frac{\dot{x}\left(iLx, x\right)}{\chi} = -\frac{i\dot{x}\left(x, Lx\right)}{\chi} = -\frac{i\dot{x}}{\chi}\left\langle [x,x]\right\rangle = 0, \tag{4.38}$$

$$Qx = (1-P)x = x. \tag{4.39}$$

$$Pa_{\mathbf{k}} = \frac{\dot{x}\left(\dot{x}, a_{\mathbf{k}}\right)}{\chi} = \frac{\dot{x}\left(iLx, a_{\mathbf{k}}\right)}{\chi} = -\frac{i\dot{x}\left(a_{\mathbf{k}}, Lx\right)}{\chi} = -\frac{i\dot{x}}{\chi}\left\langle \left[a_{\mathbf{k}}^{\dagger}, x\right]\right\rangle = 0, \tag{4.40}$$

$$Qa_{\mathbf{k}} = (1-P)a_{\mathbf{k}} = a_{\mathbf{k}} \tag{4.41}$$

The projection operators P and Q are idempotent:

$$P^2 A = \frac{\dot{x}\left(\dot{x}, \frac{\dot{x}(\dot{x},A)}{\chi}\right)}{\chi} = \frac{\dot{x}\left(\dot{x}, A\right)}{\chi} = PA;$$

$$Q^2 = (1-P)^2 = 1 - 2P + P^2 = 1 - P = Q.$$

The Liouville operator can be identiaclly represented as $L = LP + LQ$. Then the operator $\frac{1}{z-L}$in the relaxation function (4.29) can be represented as follows:

$$\frac{1}{z-L} = \frac{1}{z - LQ - LP}.$$

We use the algebraic operator identity:

$$\frac{1}{x+y} = \frac{1}{x} - \frac{1}{x}y\frac{1}{x+y}$$

with $x = z - LQ$ and $y = -LP$:

$$\frac{1}{z-L} = \frac{1}{z-LQ} + \frac{1}{z-LQ}LP\frac{1}{z-L}.$$

Consequently, the relaxation function (4.29) takes the form

$$\Phi\left(z\right) = \left(\dot{x}, \frac{1}{z-LQ}\dot{x}\right) + \left(\dot{x}, \frac{1}{z-LQ}LP\frac{1}{z-L}\dot{x}\right). \tag{4.42}$$

The first term in the r.h.s. of (4.42) simplifies as follows:

$$\left(\dot{x}, \frac{1}{z-LQ}\dot{x}\right) = \left(\dot{x}, \left[\frac{1}{z} + \frac{1}{z^2}LQ + \frac{1}{z^3}LQLQ + ...\right]\dot{x}\right) = \left(\dot{x}, \frac{1}{z}\dot{x}\right)$$

because $Q\dot{x} = 0$. Using the quantity (4.36) we obtain:

$$\left(\dot{x}, \frac{1}{z-LQ}\dot{x}\right) = \frac{\chi}{z}. \tag{4.43}$$

The second term in the r.h.s. of (4.42) contains the operator

$$P\frac{1}{z-L}\dot{x}$$

which according to the definition of the projection operator P (4.35) can be transformed as

$$P\frac{1}{z-L}\dot{x}=\frac{\dot{x}}{\chi}\left(\dot{x},\frac{1}{z-L}\dot{x}\right)=\frac{\dot{x}}{\chi}\Phi\left(z\right). \tag{4.44}$$

It is remarkable that this term is exactly expressed in terms of the sought relaxation function (4.29). Substituting (4.43) and (4.44) in (4.42), we find

$$\begin{aligned}
\Phi\left(z\right)&=\frac{\chi}{z}+\left(\dot{x},\frac{1}{z-LQ}L\frac{\dot{x}}{\chi}\right)\Phi\left(z\right)\Rightarrow \qquad (4.45)\\
z\Phi\left(z\right)&=\chi+\left(\dot{x},\frac{z}{z-LQ}L\frac{\dot{x}}{\chi}\right)\Phi\left(z\right)\\
&=\chi+\left(\dot{x},\frac{z-LQ+LQ}{z-LQ}L\frac{\dot{x}}{\chi}\right)\Phi\left(z\right)\\
&=\chi+\left(\dot{x},\left[1+\frac{LQ}{z-LQ}\right]L\frac{\dot{x}}{\chi}\right)\Phi\left(z\right)\Rightarrow\\
z\Phi\left(z\right)&=\chi+\left[\frac{(\dot{x},L\dot{x})}{\chi}+\frac{1}{\chi}\left(\dot{x},\frac{LQ}{z-LQ}L\dot{x}\right)\right]\Phi\left(z\right).
\end{aligned}$$

Introducring the quantity

$$O=\frac{(\dot{x},L\dot{x})}{\chi} \tag{4.46}$$

and the function called the *memory function*

$$\Sigma(z)=\frac{1}{\chi}\left(\dot{x},LQ\frac{1}{z-LQ}L\dot{x}\right), \tag{4.47}$$

we represent (4.45) in the form of the equation

$$\left[z-O-\Sigma(z)\right]\Phi\left(z\right)=\chi.$$

A solition of this equation gives the relaxation function $\Phi(z)$ represented within the Mori-Zwanzig projection operator technique:

$$\Phi(z)=\frac{\chi}{z-O-\Sigma(z)}. \tag{4.48}$$

The memory function (4.47) can be still transformed to another useful form.First of all, we apply the property of a scalar product (4.14):

$$\Sigma(z) = \frac{1}{\chi}\left(L\dot{x}, Q\frac{1}{z-LQ}L\dot{x}\right) \tag{4.49}$$

$$= \frac{1}{\chi}\left((P+Q)L\dot{x}, Q\frac{1}{z-LQ}L\dot{x}\right) \tag{4.50}$$

$$= \frac{1}{\chi}\left(PL\dot{x}, Q\frac{1}{z-LQ}L\dot{x}\right) + \frac{1}{\chi}\left(QL\dot{x}, Q\frac{1}{z-LQ}L\dot{x}\right). \tag{4.51}$$

For any two operators A and B

$$\begin{aligned}(PA, QB) &= \left(\frac{\dot{x}\,(\dot{x}, A)}{\chi}, \left[B - \frac{\dot{x}\,(\dot{x}, B)}{\chi}\right]\right) \\ &= \frac{(\dot{x}, A)}{\chi}\left[(\dot{x}, B) - \frac{(\dot{x}, \dot{x})\,(\dot{x}, B)}{\chi}\right] \\ &= \frac{(\dot{x}, A)}{\chi}\left[(\dot{x}, B) - (\dot{x}, B)\right] = 0,\end{aligned}$$

therefore the first term on the r.h.s. in (4.49) vanishes, and we obtain

$$\Sigma(z) = \frac{1}{\chi}\left(QL\dot{x}, Q\frac{1}{z-LQ}L\dot{x}\right). \tag{4.52}$$

In this expression, the operator $Q\frac{1}{z-LQ}$ can be represented in the following form, using the fact that Q is the idempotent operator:

$$\begin{aligned}Q\frac{1}{z-LQ} &= Q\left[\frac{1}{z} + \frac{1}{z^2}LQ + \frac{1}{z^3}LQLQ + ...\right] \\ &= \frac{1}{z}Q + \frac{1}{z^2}QLQ + \frac{1}{z^3}QLQLQ + ... \\ &= \frac{1}{z}Q + \frac{1}{z^2}QLQ^2 + \frac{1}{z^3}QLQ^2LQ^2 + ... \\ &= \left[\frac{1}{z} + \frac{1}{z^2}QLQ + \frac{1}{z^3}QLQQLQ + ...\right]Q \Rightarrow \\ Q\frac{1}{z-LQ} &= \frac{1}{z-QLQ}Q. \end{aligned} \tag{4.53}$$

A new Liouville operator can be defined, $\mathcal{L} = QLQ$, which describes the time evolution in the Hilbert space of operators, which is orthogonal complement of $\dot{x}$. Substituting then (4.53) with the operator $\mathcal{L}$ into (4.52), we bring it to the form, which will be used in what follows.

$$\Sigma(z) = \frac{1}{\chi}\left(QL\dot{x}, \frac{1}{z-\mathcal{L}}QL\dot{x}\right). \tag{4.54}$$

For the Hamiltonian (1.16) we obtain the following quantities:

$$\chi = (\dot{x}, \dot{x}) = \left(\frac{p_x}{m_b}, iLx\right) = \frac{i}{m_b\hbar}\,(p_x, Lx)$$

Using (4.15), we find

$$\chi = \frac{i}{m_b\hbar}\left\langle [p_x, x]\right\rangle = \frac{i}{m_b\hbar}\left\langle (-i\hbar)\right\rangle = \frac{1}{m_b} \tag{4.55}$$

and

$$O = \frac{(\dot{x}, L\dot{x})}{\chi} = m_b\frac{1}{\hbar}\left\langle [\dot{x}, \dot{x}]\right\rangle = 0. \tag{4.56}$$

Substituting (4.55) and (4.56) in (4.48), one obtains

$$\Phi(z) = \frac{1}{m_b}\frac{1}{z - \Sigma(z)}. \tag{4.57}$$

The operator

$$\begin{aligned}
L\dot{x} &= L\frac{p_x}{m_b} = \frac{1}{m_b\hbar}\left[H, p_x\right] = \\
&= -\frac{1}{m_b\hbar}\left[p_x, \sum_{\mathbf{k}}(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}})\right] \\
&= \frac{i}{m_b}\sum_{\mathbf{k}} ik_x(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}}) \Rightarrow \\
L\dot{x} &= -\frac{1}{m_b}\sum_{\mathbf{k}} k_x(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}})
\end{aligned} \tag{4.58}$$

does not depend on the velocities. Therefore, multiplying both parts of (4.58) with Q and taking into account (4.39) and (4.41), we obtain

$$QL\dot{x} = -\frac{1}{m_b}\sum_{\mathbf{k}} k_x(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}}),$$

what allows us to represent the memory function in the form

$$\begin{aligned}
\Sigma(z) &= \frac{1}{\chi}\left(QL\dot{x}, \frac{1}{z-\mathcal{L}}QL\dot{x}\right) \\
&= \frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'}\begin{pmatrix} k_x(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}}), \\ \frac{1}{z-\mathcal{L}}k_x'(V_{k'} a_{\mathbf{k}'} e^{i\mathbf{k}'\cdot\mathbf{r}} - V_{k'}^* a_{\mathbf{k}'}^\dagger e^{-i\mathbf{k}'\cdot\mathbf{r}}) \end{pmatrix} \\
&= \frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k_x' V_k V_{k'}^* \begin{pmatrix} (a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}}), \\ \frac{1}{z-\mathcal{L}}(a_{\mathbf{k}'} e^{i\mathbf{k}'\cdot\mathbf{r}} + a_{\mathbf{k}'}^\dagger e^{-i\mathbf{k}'\cdot\mathbf{r}}) \end{pmatrix}.
\end{aligned} \tag{4.59}$$

In transition to (4.59) we have used the property of the amplitude (1.17): $V_k^* = -V_k$ and taken into account that according to the definition (4.8), the first operator enters a scalar product in the hermitian conjugate form. Introducing the operators

$$b_{\mathbf{k}} = a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}};\, b_{\mathbf{k}}^\dagger = a_{\mathbf{k}}^\dagger e^{-i\mathbf{k}\cdot\mathbf{r}},$$

we represent the memory function as

$$\Sigma(z) = \frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k'_x V_k V^*_{k'} \left((b_{\mathbf{k}} + b^\dagger_{\mathbf{k}}), \frac{1}{z-\mathcal{L}}(b_{\mathbf{k}'} + b^\dagger_{\mathbf{k}'}) \right). \tag{4.60}$$

We notice that $Qb_{\mathbf{k}} = Q(a_{\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}}) = a_{\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}} = b_{\mathbf{k}}$. It will be represented through the four relaxation functions:

$$\Sigma(z) = \frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k'_x V_k V^*_{k'} \left[\Phi^{++}_{\mathbf{k}\mathbf{k}'}(z) + \Phi^{--}_{\mathbf{k}\mathbf{k}'}(z) + \Phi^{+-}_{\mathbf{k}\mathbf{k}'}(z) + \Phi^{-+}_{\mathbf{k}\mathbf{k}'}(z) \right], \tag{4.61}$$

$$\Phi^{++}_{\mathbf{k}\mathbf{k}'}(z) = \left(b^\dagger_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b^\dagger_{\mathbf{k}'} \right), \tag{4.62}$$

$$\Phi^{--}_{\mathbf{k}\mathbf{k}'}(z) = \left(b_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b_{\mathbf{k}'} \right), \tag{4.63}$$

$$\Phi^{+-}_{\mathbf{k}\mathbf{k}'}(z) = \left(b^\dagger_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b_{\mathbf{k}'} \right), \tag{4.64}$$

$$\Phi^{-+}_{\mathbf{k}\mathbf{k}'}(z) = \left(b_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b^\dagger_{\mathbf{k}'} \right). \tag{4.65}$$

There exist relations between the above relaxation functions. For example, the relaxation function (4.63), takes the form

$$\begin{aligned} \Phi^{--}_{\mathbf{k}\mathbf{k}'}(z) &= \left(b_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b_{\mathbf{k}'} \right) \\ &= -i\int_0^\infty dte^{izt} \left(e^{i\mathcal{L}t} b_{\mathbf{k}}(0), b_{\mathbf{k}'}(0) \right). \end{aligned}$$

Then the complex conjugate of this relaxation function:

$$\begin{aligned} \left[\Phi^{--}_{\mathbf{k}\mathbf{k}'}(z)\right]^* &= i\int_0^\infty dte^{-iz^*t} \left(e^{i\mathcal{L}t} b_{\mathbf{k}}(0), b_{\mathbf{k}'}(0) \right)^* \\ &= i\int_0^\infty dte^{-iz^*t} \left(b_{\mathbf{k}'}(0), e^{i\mathcal{L}t} b_{\mathbf{k}}(0) \right), \end{aligned}$$

where the property (4.16) has been used. The property (4.13) gives

$$\begin{aligned} \left[\Phi^{--}_{\mathbf{k}\mathbf{k}'}(z)\right]^* &= i\int_0^\infty dte^{-iz^*t} \left(b_{\mathbf{k}'}(0), e^{i\mathcal{H}t/\hbar} b_{\mathbf{k}}(0) e^{-i\mathcal{H}t/\hbar} \right) \\ &= i\int_0^\infty dte^{-iz^*t} \left(e^{i\mathcal{H}t/\hbar} b^\dagger_{\mathbf{k}}(0) e^{i\mathcal{H}t/\hbar}, b^\dagger_{\mathbf{k}'}(0) \right) \\ &= i\int_0^\infty dte^{-iz^*t} \left(e^{i\mathcal{L}t} b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k}'}(0) \right) \\ &= i\int_0^\infty dte^{-iz^*t} \left(b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0) \right) = -\Phi^{++}_{\mathbf{k}\mathbf{k}'}(-z^*), \end{aligned}$$

wherefrom it follows that

$$\Phi^{--}_{\mathbf{kk'}}(z) = -\left[\Phi^{++}_{\mathbf{kk'}}(-z^*)\right]^*. \tag{4.66}$$

Similarly, the relation

$$\Phi^{-+}_{\mathbf{kk'}}(z) = -\left[\Phi^{+-}_{\mathbf{kk'}}(-z^*)\right]^* \tag{4.67}$$

is proven.

e. Memory function In this subsection we indicate which approximations must be made in the calculation of the relaxation functions in order to obtain the FHIP results for the impedance function. Consider the relaxation function (4.62):

$$\begin{aligned}\Phi^{++}_{\mathbf{kk'}}(z) &= \left(b^\dagger_{\mathbf{k}}, \frac{1}{z-\mathcal{L}} b^\dagger_{\mathbf{k'}}\right)\\ &= -i\int_0^\infty dt e^{izt}\left(e^{i\mathcal{L}t} b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k'}}(0)\right) = -i\int_0^\infty dt e^{izt}\left(b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right),\end{aligned}$$

where $b^\dagger_{\mathbf{k}}(t) = e^{i\mathcal{L}t} b^\dagger_{\mathbf{k}}(0)$, and perform a partial integration:

$$\begin{aligned}\Phi^{++}_{\mathbf{kk'}}(z) &= -\frac{1}{z}\int_0^\infty d\left(e^{izt}\right)\left(b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right)\\ &= -\frac{1}{z} e^{izt}\left(b^\dagger_{\mathbf{k}}(t) b^\dagger_{\mathbf{k'}}(0)\right)\Bigg|_0^\infty + \frac{1}{z}\int_0^\infty dt e^{izt}\left(\frac{db^\dagger_{\mathbf{k}}(t)}{dt}, b^\dagger_{\mathbf{k'}}(0)\right)\\ &= \frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k'}}(0)\right) + \frac{1}{z}\int_0^\infty dt e^{izt}\left(i\mathcal{L} b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right)\\ &= \frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k'}}(0)\right) - \frac{i}{z}\int_0^\infty dt e^{izt}\left(\mathcal{L} b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right).\end{aligned} \tag{4.68}$$

Here we supposed that

$$\lim_{t\longrightarrow\infty} e^{izt}\left(b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right) = 0.$$

In the second term in (4.68),

$$\begin{aligned}\left(\mathcal{L} b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k'}}(0)\right) = \left(\mathcal{L} e^{i\mathcal{L}t} b^\dagger_{\mathbf{k}}, b^\dagger_{\mathbf{k'}}\right) &= \left(QLQ e^{iQLQt} b^\dagger_{\mathbf{k}}, b^\dagger_{\mathbf{k'}}(0)\right)\\ &= \left(QLQ e^{iQHQt/\hbar} b^\dagger_{\mathbf{k}} e^{-iQHQt/\hbar}, b^\dagger_{\mathbf{k'}}\right)\\ &= \left(QL e^{iQHQt/\hbar} b^\dagger_{\mathbf{k}} e^{-iQHQt/\hbar}, b^\dagger_{\mathbf{k'}}\right)\end{aligned}$$

because $Qb^\dagger_{\mathbf{k}} = b^\dagger_{\mathbf{k}}$ and $Q^2 = Q$. Further on, we have

$$\begin{aligned}
\left(\mathcal{L}b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right) &= \int_0^\beta d\lambda \left\langle e^{\lambda\hbar L} e^{iQHQt/\hbar} b_{\mathbf{k}} e^{-iQHQt/\hbar} LQ b^\dagger_{\mathbf{k}'} \right\rangle \\
&= \int_0^\beta d\lambda \left\langle e^{\lambda\hbar L} e^{iQHQt/\hbar} b_{\mathbf{k}} e^{-iQHQt/\hbar} L b^\dagger_{\mathbf{k}'} \right\rangle \\
&= \left(L e^{iQHQt/\hbar} b^\dagger_{\mathbf{k}} e^{-iQHQt/\hbar}, b^\dagger_{\mathbf{k}'}\right) \\
&= \left(L b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right).
\end{aligned}$$

So, we find from (4.68)

$$\begin{aligned}
\Phi^{++}_{\mathbf{k}\mathbf{k}'}(z) &= \frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k}'}(0)\right) - \frac{i}{z}\int_0^\infty dt e^{izt}\left(L b^\dagger_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right) \\
&= \frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k}'}(0)\right) - \frac{i}{z}\int_0^\infty dt e^{izt}\left(b^\dagger_{\mathbf{k}}(t), L b^\dagger_{\mathbf{k}'}(0)\right) \\
&= \frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k}'}(0)\right) - \frac{i}{z\hbar}\int_0^\infty dt e^{izt}\left\langle \left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle. \qquad (4.69)
\end{aligned}$$

The first term in the r.h.s. of this expression can be represented as follows:

$$\begin{aligned}
\frac{1}{z}\left(b^\dagger_{\mathbf{k}}(0), b^\dagger_{\mathbf{k}'}(0)\right) &= \frac{1}{z}\int_0^\beta d\lambda \left\langle \left(e^{\lambda\hbar L} b_{\mathbf{k}}\right) b^\dagger_{\mathbf{k}'}\right\rangle = \frac{1}{z}\left\langle \int_0^\beta d\lambda \left(e^{\lambda\hbar L} b_{\mathbf{k}}\right) b^\dagger_{\mathbf{k}'}\right\rangle \\
&= \frac{1}{z}\left\langle \left(\frac{e^{\beta\hbar L}-1}{\hbar L} b_{\mathbf{k}}\right) b^\dagger_{\mathbf{k}'}\right\rangle = \frac{1}{z\hbar}\left\langle e^{\beta H}\frac{1}{L} b_{\mathbf{k}} e^{-\beta H} b^\dagger_{\mathbf{k}'} - \frac{1}{L} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'}\right\rangle \\
&= \frac{1}{z\hbar}\frac{1}{\mathrm{Tr} e^{-\beta H}}\mathrm{Tr}\left\{e^{-\beta H}\left[e^{\beta H}\frac{1}{L} b_{\mathbf{k}} e^{-\beta H} b^\dagger_{\mathbf{k}'} - \frac{1}{L} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'}\right]\right\} \\
&= \frac{1}{z\hbar}\frac{1}{\mathrm{Tr} e^{-\beta H}}\mathrm{Tr}\left\{b^\dagger_{\mathbf{k}'}\frac{1}{L} b_{\mathbf{k}} e^{-\beta H} - e^{-\beta H}\frac{1}{L} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'}\right\} \\
&= \frac{1}{z\hbar}\left\langle b^\dagger_{\mathbf{k}'}\frac{1}{L} b_{\mathbf{k}} - \frac{1}{L} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'}\right\rangle \\
&= \frac{i}{z\hbar}\left\langle b^\dagger_{\mathbf{k}'}\frac{1}{iL} b_{\mathbf{k}} - \frac{1}{iL} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'}\right\rangle \\
&= \frac{i}{z\hbar}\left\langle \int_0^\infty dt e^{iLt} b_{\mathbf{k}} b^\dagger_{\mathbf{k}'} - \int_0^\infty dt b^\dagger_{\mathbf{k}'} e^{iLt} b_{\mathbf{k}}\right\rangle \\
&= \frac{i}{z\hbar}\left\langle \int_0^\infty dt b_{\mathbf{k}}(t) b^\dagger_{\mathbf{k}'} - \int_0^\infty dt b^\dagger_{\mathbf{k}'} b_{\mathbf{k}}(t)\right\rangle \\
&= \frac{i}{z\hbar}\int_0^\infty dt \left\langle \left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle.
\end{aligned}$$

Substituting it in the r.h.s. of (4.69), we find

$$\begin{aligned}
\Phi^{++}_{\mathbf{k}\mathbf{k}'}(z) &= \frac{i}{z\hbar}\int_0^\infty dt \left\langle \left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle - \frac{i}{z\hbar}\int_0^\infty dt e^{izt}\left\langle \left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle \\
&= \frac{i}{z\hbar}\int_0^\infty dt \left(1 - e^{izt}\right)\left\langle \left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle. \qquad (4.70)
\end{aligned}$$

In a similar way one obtains

$$\Phi^{+-}_{\mathbf{k}\mathbf{k}'}(z) = \frac{i}{z\hbar}\int_0^\infty dt\left(1-e^{izt}\right)\left\langle\left[b_{\mathbf{k}}(t), b_{\mathbf{k}'}(0)\right]\right\rangle . \tag{4.71}$$

Inserting the relaxation functions (4.70), (4.71), (4.66) and (4.67), we find the memory function (4.61)

$$\begin{aligned}
\Sigma(z) &= \frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k'_x V_k V^*_{k'} \frac{i}{z\hbar}\int_0^\infty dt\left(1-e^{izt}\right) \\
&\times \left[\begin{array}{c} \left\langle\left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle + \left\langle\left[b_{\mathbf{k}}(t), b_{\mathbf{k}'}(0)\right]\right\rangle \\ -\left\langle\left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle^* - \left\langle\left[b_{\mathbf{k}}(t), b_{\mathbf{k}'}(0)\right]\right\rangle^* \end{array}\right] \\
&= -\frac{1}{m_b}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k'_x V_k V^*_{k'} \frac{2}{z\hbar}\int_0^\infty dt\left(1-e^{izt}\right) \\
&\times \operatorname{Im}\left[\left\langle\left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle + \left\langle\left[b_{\mathbf{k}}(t), b_{\mathbf{k}'}(0)\right]\right\rangle\right],
\end{aligned}$$

wherefrom

$$\Sigma(z) = \frac{1}{z}\int_0^\infty dt\left(1-e^{izt}\right)\operatorname{Im}F(t) \tag{4.72}$$

with

$$F(t) = -\frac{2}{m_b\hbar}\sum_{\mathbf{k}}\sum_{\mathbf{k}'} k_x k'_x V_k V^*_{k'}\left\{\left\langle\left[b_{\mathbf{k}}(t), b^\dagger_{\mathbf{k}'}(0)\right]\right\rangle + \left\langle\left[b_{\mathbf{k}}(t), b_{\mathbf{k}'}(0)\right]\right\rangle\right\}. \tag{4.73}$$

f. Derivation of the memory function To calculate the expectation values in Eq. (4.73), we shall make the following approximations (cf. Ref. [76]). The Liouville operator $\mathcal{L}$, which determines the time evolution of the operator $b^\dagger_{\mathbf{k}}(t) = e^{i\mathcal{L}t}b^\dagger_{\mathbf{k}}(0)$, is replaced by $L_{ph} + L_F$, where L_{ph} is the Liouville operator for free phonons and L_F is the Liouville operator for the Feynman polaron model [41]. The Fröhlich Hamiltonian appearing in the statistical average $\langle \bullet \rangle$ is imilarly replaced by $H_{ph} + H_F$,with H_{ph} the Hamiltonian of free phonons and H_F the Hamiltonian of the Feynman polaron model. With this approximation, e.g., the average

$$\left\langle b_{\mathbf{k}}(t) b^\dagger_{\mathbf{k}'}(0)\right\rangle = \left\langle a_{\mathbf{k}}(t) a^\dagger_{\mathbf{k}'}(0)\right\rangle\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}'\cdot\mathbf{r}}\right\rangle = \delta_{\mathbf{k},\mathbf{k}'}\left\langle a_{\mathbf{k}}(t) a^\dagger_{\mathbf{k}}(0)\right\rangle\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}\cdot\mathbf{r}}\right\rangle . \tag{4.74}$$

The time evolution of the free-phonon annihilation operator (4.10),

$$a_{\mathbf{k}}(t) = e^{iH_{ph}t/\hbar} a_{\mathbf{k}} e^{-iH_{ph}t/\hbar} = \exp\left(i\omega_{\mathbf{k}} a^+_{\mathbf{k}} a_{\mathbf{k}} t\right) a_{\mathbf{k}} \exp\left(-i\omega_{\mathbf{k}} a^+_{\mathbf{k}} a_{\mathbf{k}} t\right)$$

accorting to (4.9) is

$$\begin{aligned}
-i\frac{da_{\mathbf{k}}(t)}{dt} &= \omega_{\mathbf{k}} \exp\left(i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) \left[a_{\mathbf{k}}^{+} a_{\mathbf{k}}, a_{\mathbf{k}}\right] \exp\left(-i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) \\
&= \omega_{\mathbf{k}} \exp\left(i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) \left[a_{\mathbf{k}}^{+}, a_{\mathbf{k}}\right] a_{\mathbf{k}} \exp\left(-i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) \\
&= -\omega_{\mathbf{k}} \exp\left(i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) a_{\mathbf{k}} \exp\left(-i\omega_{\mathbf{k}} a_{\mathbf{k}}^{+} a_{\mathbf{k}} t\right) \Rightarrow \\
a_{\mathbf{k}}(t) &= \exp\left(-i\omega_{\mathbf{k}} t\right) a_{\mathbf{k}}.
\end{aligned}$$

Similarly,

$$a_{\mathbf{k}}^{\dagger}(t) = \exp\left(i\omega_{\mathbf{k}} t\right) a_{\mathbf{k}}^{\dagger}.$$

Hence, we have

$$\left\langle a_{\mathbf{k}}(t) a_{\mathbf{k}}^{\dagger} \right\rangle = \exp\left(-i\omega_{\mathbf{k}} t\right) \left\langle a_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} \right\rangle = \exp\left(-i\omega_{\mathbf{k}} t\right) \left\langle 1 + a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \right\rangle = \exp\left(-i\omega_{\mathbf{k}} t\right) \left[1 + n(\omega_{\mathbf{k}})\right],$$

where $n(\omega_{\mathbf{k}}) = \left[\exp(\beta\hbar\omega_{\mathbf{k}}) - 1\right]^{-1}$ is the average number of phonons with energy $\hbar\omega_{\mathbf{k}}$.

The calculation of the Fourier component of the electron density-density correlation function $\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}\cdot\mathbf{r}} \right\rangle$ in Eq. (4.74) for an electron described by the Feynman polaron model is given below following the approach of Ref. [76].

We calculate the correlation function

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}\cdot\mathbf{r}(\tau)} \right\rangle = \frac{Tr\left(e^{-\beta H_F} e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{i\mathbf{k}\cdot\mathbf{r}(\tau)}\right)}{\mathrm{Tr}\left(e^{-\beta H_F}\right)} \tag{4.75}$$

with the Feynman trial Hamiltonian

$$H_F = \frac{\mathbf{p}^2}{2m} + \frac{\mathbf{p}_f^2}{2m_f} + \frac{1}{2}\chi\left(\mathbf{r} - \mathbf{r}_f\right)^2. \tag{4.76}$$

Here, $\mathbf{r}\,(t)$ denotes the operator in the Heisenberg representation

$$\mathbf{r}\,(t) = e^{\frac{it}{\hbar}H_F} \mathbf{r} e^{-\frac{it}{\hbar}H_F}. \tag{4.77}$$

We show that the correlation function $\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}\cdot\mathbf{r}(\tau)} \right\rangle$ depends on $(\tau - t)$ rather than on t and τ independently:

$$\begin{aligned}
\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)} e^{-i\mathbf{k}\cdot\mathbf{r}(\tau)} \right\rangle &= \frac{\mathrm{Tr}\left(e^{-\beta H_F} e^{\frac{it}{\hbar}H_F} e^{i\mathbf{k}\cdot\mathbf{r}} e^{-\frac{it}{\hbar}H_F} e^{\frac{i\tau}{\hbar}H_F} e^{i\mathbf{k}\cdot\mathbf{r}} e^{-\frac{i\tau}{\hbar}H_F}\right)}{\mathrm{Tr}\left(e^{-\beta H_F}\right)} \\
&= \frac{\mathrm{Tr}\left(e^{-\beta H_F} e^{i\mathbf{k}\cdot\mathbf{r}} e^{\frac{i(\tau-t)}{\hbar}H_F} e^{i\mathbf{k}\cdot\mathbf{r}} e^{-\frac{i(\tau-t)}{\hbar}H_F}\right)}{\mathrm{Tr}\left(e^{-\beta H_F}\right)} \\
&= \left\langle e^{i\mathbf{k}\cdot\mathbf{r}} e^{-i\mathbf{k}\cdot\mathbf{r}(\tau-t)} \right\rangle = \left\langle e^{i\mathbf{k}\cdot\mathbf{r}} e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)} \right\rangle,
\end{aligned} \tag{4.78}$$

where $\sigma = \tau - t$.

The normal coordinates are the center-of-mass vector $\mathbf{R}$ and the vector of the relative motion ρ:

$$\begin{cases} \mathbf{R} = \frac{m\mathbf{r}+m_f\mathbf{r}_f}{m+m_f} \\ \rho = \mathbf{r} - \mathbf{r}_f \end{cases} \tag{4.79}$$

The inverse transformation is:

$$\begin{cases} \mathbf{r} = \mathbf{R} + \frac{m_f}{m+m_f}\rho \\ \mathbf{r}_f = \mathbf{R} - \frac{m}{m+m_f}\rho \end{cases} \tag{4.80}$$

The same transformation as (4.79) takes place for velocities:

$$\begin{cases} \dot{\mathbf{r}} = \dot{\mathbf{R}} + \frac{m_f}{m+m_f}\dot{\rho} \\ \dot{\mathbf{r}}_f = \dot{\mathbf{R}} - \frac{m}{m+m_f}\dot{\rho} \end{cases} \tag{4.81}$$

From (4.79) we derive the transformation for moments

$$\begin{cases} \frac{\mathbf{p}}{m} = \frac{\mathbf{P}}{m+m_f} + \frac{m_f}{m+m_f}\frac{\mathbf{p}_\rho}{\frac{mm_f}{m+m_f}} \\ \frac{\mathbf{p}_f}{m_f} = \frac{\mathbf{P}}{m+m_f} - \frac{m}{m+m_f}\frac{\mathbf{p}_\rho}{\frac{mm_f}{m+m_f}} \end{cases}$$

$$\Downarrow$$

$$\begin{cases} \mathbf{p} = \frac{m}{m+m_f}\mathbf{P} + \mathbf{p}_\rho \\ \mathbf{p}_f = \frac{m_f}{m+m_f}\mathbf{P} - \mathbf{p}_\rho \end{cases} \tag{4.82}$$

The Hamiltonian (4.76) then takes the form

$$H_F = \frac{\mathbf{P}^2}{2M} + \frac{\mathbf{p}_\rho^2}{2\mu} + \frac{1}{2}\mu\bar{\Omega}^2\rho^2 \tag{4.83}$$

with the masses

$$M = m + m_f, \ \mu = \frac{mm_f}{m+m_f} \tag{4.84}$$

and with the frequency

$$\bar{\Omega} = \sqrt{\frac{\chi}{\mu}}. \tag{4.85}$$

The Cartesian coordinates and moments corresponding to the relative motion can be in the standard way expressed in terms of the second quantization operators:

$$\begin{aligned} \rho_j &= \left(\frac{\hbar}{2\mu\bar{\Omega}}\right)^{1/2}\left(C_j + C_j^\dagger\right), \\ p_{\rho,j} &= -i\left(\frac{\mu\hbar\bar{\Omega}}{2}\right)^{1/2}\left(C_j - C_j^\dagger\right). \\ &(j = 1, 2, 3) \end{aligned} \tag{4.86}$$

In these notations, the Hamiltonian (4.83) takes the form

$$H_F = \frac{\mathbf{P}^2}{2M} + \sum_{j=1}^{3} \hbar\bar{\Omega}\left(C_j^\dagger C_j + \frac{1}{2}\right). \tag{4.87}$$

Using (4.87), we find the operators in the Heisenberg representation, (i) for the center-of mass coordinates

$$\begin{aligned} X_j(\sigma) &= e^{\frac{i\sigma}{\hbar}H_F} X_j e^{-\frac{i\sigma}{\hbar}H_F} = e^{i\frac{\sigma}{2M\hbar}P_j^2} X_j e^{-i\frac{\sigma}{2M\hbar}P_j^2} \\ &= X_j + i\frac{\sigma}{2M\hbar}\left[P_j^2, X_j\right] \\ &= X_j + i\frac{\sigma}{2M\hbar}\left(P_j^2 X_j - P_j X_j P_j + P_j X_j P_j - X_j P_j^2\right) \\ &= X_j + i\frac{\sigma}{2M\hbar}\left(P_j\left[P_j, X_j\right] + \left[P_j, X_j\right]P_j\right) \\ &= X_j + i\frac{\sigma}{2M\hbar}P_j\left(-2i\hbar\right) = X_j + \frac{\sigma}{M}P_j, \end{aligned}$$

$$X_j(\sigma) = X_j + \frac{\sigma}{M}P_j, \tag{4.88}$$

(ii) for the operators C_j and $C_j^\dagger$

$$C_j(\sigma) = C_j e^{-i\bar{\Omega}\sigma},\ C_j^\dagger(\sigma) = C_j^\dagger e^{i\bar{\Omega}\sigma}. \tag{4.89}$$

Using the first formula of (4.80) we find

$$\mathbf{r}(\sigma) = \mathbf{R}(\sigma) + \frac{m_f}{m+m_f}\rho(\sigma)$$

$$\Downarrow$$

$$\mathbf{r}(\sigma) = \mathbf{R} + \frac{\sigma}{M}\mathbf{P} + \frac{m_f}{M}\left(\frac{\hbar}{2\mu\bar{\Omega}}\right)^{1/2}\left(\mathbf{C}e^{-i\bar{\Omega}\sigma} + \mathbf{C}^\dagger e^{i\bar{\Omega}\sigma}\right). \tag{4.90}$$

We denote

$$\begin{aligned} a \equiv \frac{m_f}{M}\left(\frac{\hbar}{2\mu\bar{\Omega}}\right)^{1/2} &= \left(\frac{\hbar m_f^2}{2M^2\mu\bar{\Omega}}\right)^{1/2} \\ &= \left(\frac{\hbar m_f^2}{2\left(m+m_f\right)^2 \frac{m m_f}{m+m_f}\bar{\Omega}}\right)^{1/2} = \left(\frac{\hbar m_f}{2mM\bar{\Omega}}\right)^{1/2} \end{aligned}$$

$$\mathbf{r}(\sigma) = \mathbf{R} + \frac{\sigma}{M}\mathbf{P} + a\left(\mathbf{C}e^{-i\bar{\Omega}\sigma} + \mathbf{C}^\dagger e^{i\bar{\Omega}\sigma}\right). \tag{4.91}$$

Therefore, we obtain

$$\begin{aligned} e^{i\mathbf{k}\cdot\mathbf{r}} &= \exp\left(i\mathbf{k}\cdot\mathbf{R}\right)\exp\left(ia\mathbf{k}\cdot\mathbf{C} + ia\mathbf{k}\cdot\mathbf{C}^\dagger\right) \\ &= \prod_{j=1}^{3}\exp\left(ik_j X_j\right)\exp\left(iak_j C_j - iak_j C_j^\dagger\right) \end{aligned}$$

$$\begin{aligned}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)} &= \exp\left[-i\mathbf{k}\cdot\mathbf{R} - i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P} - ia\mathbf{k}\cdot\left(\mathbf{C}e^{-i\bar{\Omega}\sigma} + \mathbf{C}^{\dagger}e^{i\bar{\Omega}\sigma}\right)\right] \\ &= \exp\left(-i\mathbf{k}\cdot\mathbf{R} - i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}\right)\exp\left(-ia\mathbf{k}\cdot\mathbf{C}e^{-i\bar{\Omega}\sigma} - ia\mathbf{k}\cdot\mathbf{C}^{\dagger}e^{i\bar{\Omega}\sigma}\right) \\ &= \prod_{j=1}^{3}\exp\left(-ik_jX_j - i\frac{\sigma}{M}k_jP_j\right)\exp\left(-iak_jC_je^{-i\bar{\Omega}\sigma} - iak_jC_j^{\dagger}e^{i\bar{\Omega}\sigma}\right).\end{aligned} \tag{4.92}$$

The disentangling of the exponents is performed using the formula

$$e^{A+B} = e^{A}T\exp\left(\int_0^1 d\lambda e^{-\lambda A}Be^{\lambda A}\right). \tag{4.93}$$

In the case when $[A,B]$ commutes with both A and B, this formula is reduced to

$$e^{A+B} = e^{A}e^{B}e^{-\frac{1}{2}[A,B]}. \tag{4.94}$$

We perform the necessary commutations:

$$\begin{gathered}-\frac{1}{2}\left[-ik_jX_j, -i\frac{\sigma}{M}k_jP_j\right] = \frac{1}{2}k_j^2\frac{\sigma}{M}\left[X_j, P_j\right] = i\frac{\hbar k_j^2}{2M}\sigma, \\ -\frac{1}{2}\left[iak_jC_j^{\dagger}, iak_jC_j\right] = \frac{1}{2}a^2k_j^2\left[C_j^{\dagger}, C_j\right] = -\frac{1}{2}a^2k_j^2 \\ -\frac{1}{2}\left[-iak_jC_j^{\dagger}e^{i\Omega\sigma}, -iak_jC_je^{-i\Omega\sigma}\right] = \frac{1}{2}a^2k_j^2\left[C_j^{\dagger}, C_j\right] = -\frac{1}{2}a^2k_j^2 \\ \Downarrow\end{gathered}$$

$$\begin{aligned}e^{i\mathbf{k}\cdot\mathbf{r}} &= e^{i\mathbf{k}\cdot\mathbf{R}}e^{ia\mathbf{k}\cdot\mathbf{C}^{\dagger}}e^{ia\mathbf{k}\cdot\mathbf{C}}e^{-\frac{1}{2}a^2k^2}, \\ e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)} &= e^{-i\mathbf{k}\cdot\mathbf{R}}e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}e^{-ia\mathbf{k}\cdot\mathbf{C}^{\dagger}e^{i\bar{\Omega}\sigma}}e^{-ia\mathbf{k}\cdot\mathbf{C}e^{-i\bar{\Omega}\sigma}}e^{i\frac{\hbar k^2}{2M}\sigma - \frac{1}{2}a^2k^2}\end{aligned}$$

$$\Downarrow$$

$$e^{i\mathbf{k}\cdot\mathbf{r}}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)} = e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}e^{ia\mathbf{k}\cdot\mathbf{C}^{\dagger}}e^{ia\mathbf{k}\cdot\mathbf{C}}e^{-ia\mathbf{k}\cdot\mathbf{C}^{\dagger}e^{i\bar{\Omega}\sigma}}e^{-ia\mathbf{k}\cdot\mathbf{C}e^{-i\bar{\Omega}\sigma}}e^{i\frac{\hbar k^2}{2M}\sigma - a^2k^2}.$$

It follows from Eq. (4.94) that when $[A,B]$ commutes with both A and B,

$$e^{A}e^{B} = e^{B}e^{A}e^{[A,B]}. \tag{4.95}$$

Using (4.95), we find

$$\begin{aligned}e^{iak_j\cdot C_j}e^{-iak_j\cdot C_j^{\dagger}e^{i\bar{\Omega}\sigma}} &= e^{-iak_j\cdot C_j^{\dagger}e^{i\bar{\Omega}\sigma}}e^{iak_j\cdot C_j}e^{\left[iak_j\cdot C_j, -iak_j\cdot C_j^{\dagger}e^{i\bar{\Omega}\sigma}\right]} \\ &= e^{-iak_j\cdot C_j^{\dagger}e^{i\bar{\Omega}\sigma}}e^{iak_j\cdot C_j}e^{a^2k_j^2e^{i\bar{\Omega}\sigma}}.\end{aligned}$$

Herefrom, we find

$$e^{i\mathbf{k}\cdot\mathbf{r}}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)} = e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}e^{ia\mathbf{k}\cdot\mathbf{C}^\dagger\left(1-e^{i\bar{\Omega}\sigma}\right)}e^{ia\mathbf{k}\cdot\mathbf{C}\left(1-e^{-i\bar{\Omega}\sigma}\right)}e^{i\frac{\hbar k^2}{2M}\sigma-a^2k^2\left(1-e^{i\bar{\Omega}\sigma}\right)}. \tag{4.96}$$

The correlation function then is

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)}\right\rangle = \left\langle e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}\right\rangle\left\langle e^{ia\mathbf{k}\cdot\mathbf{C}^\dagger\left(1-e^{i\bar{\Omega}\sigma}\right)}e^{ia\mathbf{k}\cdot\mathbf{C}\left(1-e^{-i\bar{\Omega}\sigma}\right)}\right\rangle e^{i\frac{\hbar k^2}{2M}\sigma-a^2k^2\left(1-e^{i\bar{\Omega}\sigma}\right)}, \tag{4.97}$$

since the variables of the center-of mass motion and of the relative motion are averaged independently.

$$\begin{aligned}\left\langle e^{ia\mathbf{k}\cdot\mathbf{C}^\dagger\left(1-e^{i\bar{\Omega}\sigma}\right)}e^{ia\mathbf{k}\cdot\mathbf{C}\left(1-e^{-i\bar{\Omega}\sigma}\right)}\right\rangle &= \left\langle e^{i\mathbf{Q}\cdot\mathbf{C}^\dagger}e^{i\mathbf{Q}^*\cdot\mathbf{C}}\right\rangle \\ &= \frac{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}\mathbf{C}^\dagger\cdot\mathbf{C}}e^{i\mathbf{Q}\cdot\mathbf{C}^\dagger}e^{i\mathbf{Q}^*\cdot\mathbf{C}}\right)}{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}\mathbf{C}^\dagger\cdot\mathbf{C}}\right)}\end{aligned}$$

with

$$\mathbf{Q} = a\left(1-e^{i\bar{\Omega}\sigma}\right)\mathbf{k}.$$

Let us consider the auxiliary expectation value

$$\begin{aligned}\left\langle e^{iQC^\dagger}e^{iQ^*C}\right\rangle &\equiv \frac{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}e^{iQC^\dagger}e^{iQ^*C}\right)}{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\right)} \\ &= \frac{1}{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\right)}\sum_{n=0}^{\infty}\frac{(iQ)^n}{n!}\sum_{m=0}^{\infty}\frac{(iQ^*)^m}{m!}\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\left(C^\dagger\right)^n C^m\right) \\ &= \frac{1}{\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\right)}\sum_{n=0}^{\infty}\frac{(-1)^n|Q|^{2n}}{(n!)^2}\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\left(C^\dagger\right)^n C^n\right).\end{aligned}$$

$$\begin{aligned}\mathrm{Tr}\left(e^{-\beta\hbar\bar{\Omega}C^\dagger C}\left(C^\dagger\right)^n C^n\right) &= \sum_{m=0}^{\infty}\left\langle m\left|e^{-\beta\hbar\bar{\Omega}C^\dagger C}\left(C^\dagger\right)^n C^n\right|m\right\rangle \\ &= \sum_{m=0}^{\infty}e^{-\beta\hbar\bar{\Omega}m}\left\langle m\left|\left(C^\dagger\right)^n C^n\right|m\right\rangle,\end{aligned}$$

where $|m\rangle$ are the eigenstates of $\left(C^\dagger C\right)$. The operators C act on these states as follows:

$$\begin{aligned}C|m\rangle &= \sqrt{m}\,|m-1\rangle, \\ C|0\rangle &= 0.\end{aligned}$$

Therefore, we find

$$\left\langle m\left|\left(C^\dagger\right)^n C^n\right|m\right\rangle = m(m-1)\ldots(m-n+1) = \frac{m!}{(m-n)!}\text{ for } n\le m,$$

$$\left\langle m\left|\left(C^\dagger\right)^n C^n\right|m\right\rangle = 0 \text{ for } n > m.$$

$$\Downarrow$$

$$Tr\left(e^{-\beta\hbar\bar{\Omega}C^{\dagger}C}\left(C^{\dagger}\right)^{n}C^{n}\right)=\sum_{m=n}^{\infty}e^{-\beta\hbar\Omega m}\frac{m!}{(m-n)!},$$

and

$$\begin{aligned}
\mathrm{Tr}\left(e^{-\beta\hbar\Omega C^{\dagger}C}e^{iQC^{\dagger}}e^{iQ^{*}C}\right) &= \sum_{n=0}^{\infty}\frac{(-1)^{n}\left|Q\right|^{2n}}{(n!)^{2}}\sum_{m=n}^{\infty}e^{-\beta\hbar\bar{\Omega}m}\frac{m!}{(m-n)!}\\
&= \sum_{n=0}^{\infty}\frac{(-1)^{n}\left|Q\right|^{2n}}{n!}\sum_{m=n}^{\infty}e^{-\beta\hbar\bar{\Omega}m}\binom{m}{n}\\
&= \sum_{n=0}^{\infty}\frac{(-1)^{n}\left|Q\right|^{2n}}{n!}e^{-\beta\hbar\bar{\Omega}n}\sum_{k=0}^{\infty}e^{-\beta\hbar\bar{\Omega}k}\binom{k+n}{n}\\
&= \sum_{n=0}^{\infty}\frac{(-1)^{n}\left|Q\right|^{2n}}{n!}e^{-\beta\hbar\bar{\Omega}n}\frac{1}{\left(1-e^{-\beta\hbar\bar{\Omega}}\right)^{n+1}}\\
&= \frac{1}{1-e^{-\beta\hbar\bar{\Omega}}}\sum_{n=0}^{\infty}\frac{(-1)^{n}\left|Q\right|^{2n}}{n!}\frac{1}{\left(e^{\beta\hbar\bar{\Omega}}-1\right)^{n}}\\
&= \frac{1}{1-e^{-\beta\hbar\bar{\Omega}}}\exp\left(-\frac{\left|Q\right|^{2}}{e^{\beta\hbar\bar{\Omega}}-1}\right).
\end{aligned}$$

In particular, for $Q=Q^{*}=0$, we have

$$Tr\left(e^{-\beta\hbar\Omega C^{\dagger}C}\right)=\frac{1}{1-e^{-\beta\hbar\Omega}}. \tag{4.98}$$

As a result, the expectation value $\left\langle e^{iQC^{\dagger}}e^{iQ^{*}C}\right\rangle$ is

$$\left\langle e^{iQC^{\dagger}}e^{iQ^{*}C}\right\rangle=\exp\left[-n\left(\Omega\right)\left|Q\right|^{2}\right], \tag{4.99}$$

with

$$n\left(\Omega\right)\equiv\frac{1}{e^{\beta\hbar\Omega}-1}.$$

Using this result, we obtain the expression

$$\begin{aligned}
\left\langle e^{i\mathbf{Q}\cdot\mathbf{C}^{\dagger}}e^{i\mathbf{Q}^{*}\cdot\mathbf{C}}\right\rangle &= \exp\left[-n\left(\Omega\right)\mathbf{Q}\cdot\mathbf{Q}^{*}\right]\\
&= \exp\left[-n\left(\Omega\right)a^{2}k^{2}\left(1-e^{i\bar{\Omega}\sigma}\right)\left(1-e^{-i\bar{\Omega}\sigma}\right)\right]\\
&= \exp\left[-4n\left(\Omega\right)a^{2}k^{2}\sin^{2}\left(\frac{1}{2}\Omega\sigma\right)\right].
\end{aligned}$$

The expectation value $\left\langle e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}\right\rangle$ is

$$\begin{aligned}\left\langle e^{-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}}\right\rangle &= \frac{\int d\mathbf{P}\exp\left(-\beta\frac{P^2}{2M}-i\frac{\sigma}{M}\mathbf{k}\cdot\mathbf{P}\right)}{\int d\mathbf{P}\exp\left(-\beta\frac{P^2}{2M}\right)}\\ &= \frac{\int d\mathbf{P}\exp\left(\frac{(-i\mathbf{P}\beta+\mathbf{k}\sigma)^2}{2M\beta}-\frac{k^2\sigma^2}{2M\beta}\right)}{\int d\mathbf{P}\exp\left(-\beta\frac{P^2}{2M}\right)}\\ &= \exp\left(-\frac{k^2\sigma^2}{2M\beta}\right).\end{aligned}$$

Collecting all factors in Eq. (4.97) together, we find

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)}\right\rangle = \exp\left[i\frac{\hbar k^2}{2M}\sigma-\frac{k^2\sigma^2}{2M\beta}-4n\left(\bar{\Omega}\right)a^2k^2\sin^2\left(\frac{1}{2}\bar{\Omega}\sigma\right)-a^2k^2\left(1-e^{i\bar{\Omega}\sigma}\right)\right]\Rightarrow .$$

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma+t)}\right\rangle = e^{-k^2D(\sigma)} \tag{4.100}$$

with the function

$$D(t)=\frac{\hbar}{2M}\left(-it+\frac{t^2}{\beta\hbar}\right)+a^2\left[1-\exp(i\bar{\Omega}t)+4n\left(\bar{\Omega}\right)\sin^2\left(\frac{\bar{\Omega}t}{2}\right)\right], \tag{4.101}$$

where

$$M=\left(\frac{v}{w}\right)^2 m_b, \bar{\Omega}=v\omega_{\mathrm{LO}}, a^2=\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\frac{v^2-w^2}{v^3}. \tag{4.102}$$

According to (4.78),

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma+t)}\right\rangle = \left\langle e^{i\mathbf{k}\cdot\mathbf{r}}e^{-i\mathbf{k}\cdot\mathbf{r}(\sigma)}\right\rangle.$$

Taking $\sigma=-t$ in (4.100) we finally find the Fourier component of the electron density-density correlation function $\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)}e^{-i\mathbf{k}\cdot\mathbf{r}}\right\rangle$ which enters Eq. (4.74):

$$\left\langle e^{i\mathbf{k}\cdot\mathbf{r}(t)}e^{-i\mathbf{k}\cdot\mathbf{r}}\right\rangle = \exp\left[-k^2D(-t)\right] \tag{4.103}$$

Finally, the correlation functions in (4.73) reduce to

$$\begin{aligned}\left\langle b_{\mathbf{k}}(t)b^{\dagger}_{\mathbf{k}}(0)\right\rangle &= [1+n(\omega_{\mathbf{k}})]\exp\left(-i\omega_{\mathbf{k}}t\right)\exp\left[-k^2D(-t)\right],\\ \left\langle b^{\dagger}_{\mathbf{k}}(0)b_{\mathbf{k}}(t)\right\rangle &= n(\omega_{\mathbf{k}})\exp\left(-i\omega_{\mathbf{k}}t\right)\exp\left[-k^2D(t)\right],\\ \langle b_{\mathbf{k}}(t)b_{\mathbf{k}}(0)\rangle &= 0,\\ \langle b_{\mathbf{k}}(0)b_{\mathbf{k}}(t)\rangle &= 0.\end{aligned}$$

Inserting these equations into Eqs. (4.72) and (4.73), one obtains

$$\Sigma(z)=\frac{1}{z}\int_0^{\infty}dt\left(1-e^{izt}\right)\operatorname{Im}S(t) \tag{4.104}$$

with

$$\operatorname{Im} S(t) = \frac{2}{m_b \hbar} \sum_{\mathbf{k}} k_x^2 \left| V_k \right|^2 \operatorname{Im} \left\{ \begin{array}{c} -\left[1+n(\omega_{\mathbf{k}})\right] \exp\left(-i\omega_{\mathbf{k}} t\right) \exp\left[-k^2 D(-t)\right] \\ +n(\omega_{\mathbf{k}}) \exp\left(-i\omega_{\mathbf{k}} t\right) \exp\left[-k^2 D(t)\right] \end{array} \right\}.$$

Using the property $D(-t)^* =^2 D(t)$ for real vaues of t,one obtains

$$\begin{aligned} -\operatorname{Im} \exp\left(-i\omega_{\mathbf{k}} t\right) \exp\left[-k^2 D(-t)\right]\} &= \operatorname{Im} \exp\left(i\omega_{\mathbf{k}} t\right) \exp\left[-k^2 D(-t)^*\right]\} \\ &= \operatorname{Im} \exp\left(i\omega_{\mathbf{k}} t\right) \exp\left[-k^2 D(t)\right]\} \end{aligned}$$

and consequently

$$S(t) = \frac{2}{m_b \hbar} \sum_{\mathbf{k}} k_x^2 \left| V_k \right|^2 \exp\left[-k^2 D(t)\right] \left\{ \left[1+n(\omega_{\mathbf{k}})\right] \exp\left(i\omega_{\mathbf{k}} t\right) + n(\omega_{\mathbf{k}}) \exp\left(-i\omega_{\mathbf{k}} t\right) \right\}. \tag{4.105}$$

Owing to the rotational invariance of $\left| V_k \right|^2$ and $\omega_{\mathbf{k}}$,we can substitute in (4.105)

$$k_x^2 \to \frac{k^2}{3}.$$

The resulting expression for

$$S(t) = \frac{2}{3m_b \hbar} \sum_{\mathbf{k}} k^2 \left| V_k \right|^2 \exp\left[-k^2 D(t)\right] \left\{ \left[1+n(\omega_{\mathbf{k}})\right] \exp\left(i\omega_{\mathbf{k}} t\right) + n(\omega_{\mathbf{k}}) \exp\left(-i\omega_{\mathbf{k}} t\right) \right\} \tag{4.106}$$

is identical with Eq. (35) of FHIP [45]. In the case of Fröhlich polarons, taking into account (1.17), Eq. (4.106) simplifies to

$$S(t) = \left(\frac{\hbar \omega_{\mathrm{LO}}}{m_b} \right)^{3/2} \frac{\alpha}{3\sqrt{2\pi}} \left\{ \begin{array}{c} \left[1+n(\omega_{\mathrm{LO}})\right] \exp\left(i\omega_{\mathrm{LO}} t\right) \\ +n(\omega_{\mathrm{LO}}) \exp\left(-i\omega_{\mathrm{LO}} t\right) \end{array} \right\} \left[D(t)\right]^{-3/2}. \tag{4.107}$$

B. Calculation of the memory function (Devreese *et. al.* [48])

Upon substituion of (4.30) and (4.57) into Eq. (4.5), we find the absorption coefficient

$$\begin{aligned} \Gamma(\Omega) &= \frac{1}{n\epsilon_0 c} \operatorname{Re} \left[\frac{ie^2}{m_b} \frac{1}{\Omega - \Sigma(\Omega)} \right] = -\frac{1}{n\epsilon_0 c} \frac{e^2}{m_b} \operatorname{Im} \left[\frac{1}{\Omega - \Sigma(\Omega)} \right] \\ &= -\frac{1}{n\epsilon_0 c} \frac{e^2}{m_b} \operatorname{Im} \left\{ \frac{\Omega - \Sigma^*(\Omega)}{\left[\Omega - \operatorname{Re}\Sigma(\Omega)\right]^2 + \left[\operatorname{Im}\Sigma(\Omega)\right]^2} \right\} \\ &= \frac{1}{n\epsilon_0 c} \frac{e^2}{m_b} \frac{\operatorname{Im}\Sigma^*(\Omega)}{\left[\Omega - \operatorname{Re}\Sigma(\Omega)\right]^2 + \left[\operatorname{Im}\Sigma(\Omega)\right]^2} \Rightarrow \end{aligned}$$

$$\Gamma(\Omega) = -\frac{1}{n\epsilon_0 c}\frac{e^2}{m_b}\frac{\operatorname{Im}\Sigma(\Omega)}{\left[\Omega - \operatorname{Re}\Sigma(\Omega)\right]^2 + \left[\operatorname{Im}\Sigma(\Omega)\right]^2} \; . \tag{4.108}$$

This general expression was the strating point for a derivation of the theoretical optical absorption spectrum of a single large polaron, at *all electron-phonon coupling strengths* by Devreese et al. in Ref. [48]. The memory function $\Sigma(\Omega)$ as given by Eq. (4.104) with (4.107) contains the dynamics of the polaron and depends on α, temperature and Ω.Following the notation, introduced in Ref. [45],

$$\Sigma(\Omega) = \frac{\chi^*(\Omega)}{\Omega} \tag{4.109}$$

we reresent Eq. (4.108) in the form used in Ref. [48]:

$$\Gamma(\Omega) = \frac{1}{n\epsilon_0 c}\frac{e^2}{m_b}\frac{\Omega \operatorname{Im}\chi(\Omega)}{\left[\Omega^2 - \operatorname{Re}\chi(\Omega)\right]^2 + \left[\operatorname{Im}\chi(\Omega)\right]^2}. \tag{4.110}$$

According to (4.104) and (4.109),

$$\operatorname{Im}\chi(\Omega) = \operatorname{Im}\int_0^\infty dt \sin(\Omega t) S(t), \quad \operatorname{Re}\chi(\Omega) = \operatorname{Im}\int_0^\infty dt \left[1 - \cos(\Omega t)\right] S(t). \tag{4.111}$$

In the present Notes we limit our attention to the case $T = 0$ ($\beta \to \infty$). It was demonstrated in Ref.[48] that the exact zero-temperature limit arises if the limit $\beta \to \infty$ is taken directly in the expressions (4.111) (see Appendices A and B of Ref.[48]). As follows from (4.107),

$$S(t) = \left(\frac{\hbar\omega_{\mathrm{LO}}}{m_b}\right)^{3/2} \frac{\alpha}{3\sqrt{2}\pi} \exp\left(i\omega_{\mathrm{LO}} t\right) \left[D(t)\right]^{-3/2} \qquad (\beta \to \infty). \tag{4.112}$$

Accorting to (4.101)

$$D(t) = -i\frac{\hbar t}{2M} + a^2\left[1 - \exp(i\bar{\Omega} t)\right] \qquad (\beta \to \infty).$$

Using the Feynman units (where $\hbar = 1, \omega_{\mathrm{LO}} = 1$ and $m_b = 1$), we obtain from (4.102):

$$M = \left(\frac{v}{w}\right)^2, \bar{\Omega} = v, a^2 = \frac{1}{2}\frac{v^2 - w^2}{v^3},$$

and consequently

$$D(t) = \frac{1}{2}\frac{v^2 - w^2}{v^3}(1 - e^{ivt}) - i\frac{1}{2}\left(\frac{w}{v}\right)^2 t = \frac{1}{2}\left(\frac{w}{v}\right)^2\left\{R(1 - e^{ivt}) - it\right\} \qquad (\beta \to \infty) \tag{4.113}$$

with

$$R = \frac{v^2 - w^2}{w^2 v}.$$

and according to (4.112)

$$S(t) = \frac{2\alpha}{3\sqrt{\pi}} \left(\frac{v}{w}\right)^3 e^{it} \left[R(1 - e^{ivt}) - it\right]^{-3/2} \qquad (\beta \to \infty). \tag{4.114}$$

From (4.111) one obtains immediately

$$\operatorname{Im} \chi(\Omega) = \frac{2\alpha}{3\sqrt{\pi}} \left(\frac{v}{w}\right)^3 \operatorname{Im} \int_0^\infty dt \frac{\sin(\Omega t) e^{it}}{\left[R(1 - e^{ivt}) - it\right]^{3/2}}, \tag{4.115}$$

$$\operatorname{Re} \chi(\Omega) = \frac{2\alpha}{3\sqrt{\pi}} \left(\frac{v}{w}\right)^3 \operatorname{Im} \int_0^\infty dt \frac{\left[1 - \cos(\Omega t)\right] e^{it}}{\left[R(1 - e^{ivt}) - it\right]^{3/2}}. \tag{4.116}$$

In the limit $\beta \to \infty$ the function $\operatorname{Im} \chi(\Omega)$ was calculated by FHIP [45]. However, to study the optical absorption to the same approximation as FHIP's treatment of the impedance, we have also to calculate $\operatorname{Re} \chi(\Omega)$ and use this result in (4.110). The calculation of $\operatorname{Re} \chi(\Omega)$, which is a Kramers-Kronig-type transform of $\operatorname{Im} \chi(\Omega)$, is a key ingredient in Ref. [48]. The details of those calculations are presented in the Appendices A, B and C to Ref. [48].

Developing the denominator of both integrals on the right-hand side of (4.115) and (4.116), the calculations are reduced to the evaluation of a sum of integrals of the type

$$\operatorname{Im} \int_0^\infty dt \frac{\sin(\Omega t) e^{i(1+nv)t}}{(R - it)^{3/2+n}}, \qquad \operatorname{Im} \int_0^\infty dt \frac{\cos(\Omega t) e^{i(1+nv)t}}{(R - it)^{3/2+n}}. \tag{4.117}$$

In Appendix B to Ref. [48] it is shown how such integrals are evaluated using a recurrence formula. For $\operatorname{Im} \chi(\Omega)$ a very convenient result was found in [48]:

$$\begin{aligned} \operatorname{Im} \chi(\Omega) = \frac{2\alpha}{3} \left(\frac{v}{w}\right)^3 \sum_{n=0}^{\infty} C^n_{-3/2} (-1)^n \frac{R^n 2^n}{(2n+1)...3 \cdot 1} \\ \times \left|\Omega - 1 - nv\right|^{n+1/2} e^{-|\Omega - 1 - nv| R} \frac{1 + \operatorname{sgn}(\Omega - 1 - nv)}{2}. \end{aligned} \tag{4.118}$$

This expression is a finite sum and not and infinite series. FHIP gave the first two terms of (4.118) explicitly.

Using the same recurrence relation it is seen the analytical expression (see Appendix B to Ref. [48]), which was found for $\operatorname{Re} \chi(\Omega)$ is far more complicated. To circumwent the difficulty with the numerical treatment of $\operatorname{Re} \chi(\Omega)$, the corresponding integrals in (4.117) have been transformed in [48] to integrals with rapildy convergent integrands:

$$\begin{aligned}
&\operatorname{Im}\int_0^\infty dt \frac{[1-\cos(\Omega t)]\, e^{i(1+nv)t}}{(R-it)^{3/2+n}} \\
&= -\frac{1}{\Gamma(n+\frac{3}{2})}\int_0^\infty dx \left[(n+\frac{1}{2})x^{n-1/2}e^{-Rx} - Rx^{n+1/2}e^{-Rx}\right] \\
&\times \ln\left|\left(\frac{(1+nv+x)^2}{\Omega^2-(1+nv+x)^2}\right)\right|^{1/2}. \qquad (4.119)
\end{aligned}$$

The integral on the right-hand side of (4.119) is adequate for computer calculations. In Appendix C to Ref. [48] some supplementary details of the computation of (4.119) are given. Another analytical representation for the memory function (4.104) was derived in Ref. [74].

C. Discussion of optical absorption of polarons at arbitrary coupling

At weak coupling, the optical absorption spectrum (4.108) of the polaron is determined by the absorption of radiation energy, which is reemitted in the form of LO phonons. For $\alpha \gtrsim 5.9$, the polaron can undergo transitions toward a relatively stable RES (see Fig. 7). The RES peak in the optical absorption spectrum also has a phonon sideband-structure, whose average transition frequency can be related to a FC-type transition. Furthermore, at zero temperature, the optical absorption spectrum of one polaron exhibits also a zero-frequency "central peak" [$\sim \delta(\Omega)$]. For non-zero temperature, this "central peak" smears out and gives rise to an "anomalous" Drude-type low-frequency component of the optical absorption spectrum.

For example, in Fig. 7 from Ref. [48], the main peak of the polaron optical absorption for $\alpha = 5$ at $\Omega = 3.51\omega_{\mathrm{LO}}$ is interpreted as due to transitions to a RES. A "shoulder" at the low-frequency side of the main peak is attributed to one-phonon transitions to polaron-"scattering states". The broad structure centered at about $\Omega = 6.3\omega_{\mathrm{LO}}$ is interpreted as a FC band. As seen from Fig. 7, when increasing the electron-phonon coupling constant to α=6, the RES peak at $\Omega = 4.3\omega_{\mathrm{LO}}$ stabilizes. It is in Ref. [48] that the all-coupling optical absorption spectrum of a Fröhlich polaron, together with the role of RES-states, FC-states and scattering states, was first presented.

Recent interesting numerical calculations of the optical conductivity for the Fröhlich polaron performed within the diagrammatic Quantum Monte Carlo method [77], see Fig. 8,



FIG. 7: Optical absorption spectrum of a polaron calculated by Devreese *et al.* [48] $\alpha = 5$ and 6. The RES peak is very intense compared with the FC peak. The frequency $\Omega/\omega_{\mathrm{LO}} = v$ is indicated by the dashed lines.)

fully confirm the essential analytical results derived by Devreese et al. in Ref. [48] for $\alpha \lesssim 3$. In the intermediate coupling regime $3 < \alpha < 6$, the low-energy behavior and the position of the RES-peak in the optical conductivity spectrum of Ref. [77] follow closely the prediction of Ref. [48]. There are some minor qualitative differences between the two approaches in the intermediate coupling regime: in Ref. [77], the dominant ("RES") peak is less intense in the Monte-Carlo numerical simulations and the second ("FC") peak develops less prominently. There are the following qualitative differences between the two approaches in the strong coupling regime: in Ref.[77], the dominant peak broadens and the second peak does not develop, giving instead rise to a flat shoulder in the optical conductivity spectrum at $\alpha = 6$. This behavior has been tentatively attributed to the optical processes with participation of two [73] or more phonons. The above differences can arise also due to the fact that, within the Feynman polaron model, one-phonon processes are assigned more oscillator strength and the RES tends to be more stable as compared to the Monte-Carlo result. The nature of the excited states of a polaron needs further study. An independent numerical simulation might be called for.



FIG. 8: *Left-hand panel*: Monte Carlo optical conductivity spectra of one polaron for the weak-coupling regime (open circles) compared to the second-order perturbation theory (dotted lines) for $\alpha = 0.01$ and $\alpha = 1$ and to the analytical DSG calculations [48] (solid lines). *Right-hand panel*: Monte Carlo optical conductivity spectra for the intermediate coupling regime (open circles) compared to the analytical DSG approach [48] (solid lines). Arrows point to the two- and three-phonon thresholds. (From Ref. [77].)

In Fig. 9, Monte-Carlo optical conductivity spectrum of one polaron for $\alpha = 1$ compares well with that obtained in Ref. [78] within the canonical-transformation formalism taking into account correlation in processes involving two LO phonons. The difference between the results of these two approaches becomes less pronounced when decreasing the value of $\alpha = 1$ and might be indicative of a possible precision loss, which requires an independent check.



FIG. 9: One-polaron optical conductivity $\mathrm{Re}\,\sigma(\omega)$ for $\alpha = 1$ calculated within the Monte Carlo approach [77] (open circles) and derived using the expansion in powers of α up to α^2 [78] (solid curve).

The coupling constant α of the known ionic crystals is too small ($\alpha < 5$) to allow for the experimental detection of sharp RES peaks, and the resonance condition $\Omega = Re\Sigma(\Omega)$ cannot be satisfied for $\alpha \lesssim 5.9$ as shown in Ref, [48]. Nevertheless, for $3 \lesssim \alpha \lesssim 5.9$ the development of RES is already reflected in a broad optical absorption peak. Such a peak, predicted in Ref. [48], was identified, e. g., in the optical absorption of $\mathrm{Pr_2NiO_{4.22}}$ in Ref. [79]. Also, the resonance condition can be fulfilled if an external magnetic field is applied; the magnetic field stabilizes the RES, which then can be detected in a cyclotron resonance peak.

1. *Sum rules for the optical conductivity spectra*

In this section, we analyze the sum rules for the optical conductivity spectra obtained within the DSG approach [48] with those obtained within the diagrammatic Monte Carlo calculation [77]. The values of the polaron effective mass for the Monte Carlo approach are taken from Ref. [28]. In Tables 3 and 4, we represent the polaron ground-state E_0 and the following parameters calculated using the optical conductivity spectra:

$$M_0 \equiv \int_1^{\omega_{\max}} \operatorname{Re}\sigma(\omega)\, d\omega, \tag{4.120}$$

$$M_1 \equiv \int_1^{\omega_{\max}} \omega \operatorname{Re}\sigma(\omega)\, d\omega, \tag{4.121}$$

where $\omega_{\max}$ is the upper value of the frequency available from Ref. [77],

$$\tilde{M}_0 \equiv \frac{\pi}{2m^*} + \int_1^{\omega_{\max}} \operatorname{Re}\sigma(\omega)\, d\omega, \tag{4.122}$$

where m^* is the polaron mass, the optical conductivity is calculated in units $\frac{n_0 e^2}{m_b \omega_{\mathrm{LO}}}$, m^* is measured in units of the band mass m_b, and the frequency is measured in units of ω_{LO}. The values of $\omega_{\max}$ are: $\omega_{\max} = 10$ for $\alpha = 0.01$, 1 and 3, $\omega_{\max} = 12$ for $\alpha = 4.5$, 5.25 and 6, $\omega_{\max} = 18$ for $\alpha = 6.5$, 7 and 8.

Table 3. Polaron parameters obtained within the diagrammatic Monte Carlo approach

α	$M_0^{(\mathrm{MC})}$	$m^{*(\mathrm{MC})}$	$\tilde{M}_0^{(\mathrm{MC})}$	$M_1^{(\mathrm{MC})}/\alpha$	$E_0^{(\mathrm{MC})}$
0.01	0.00249	1.0017	1.5706	0.634	-0.010
1	0.24179	1.1865	1.5657	0.65789	-1.013
3	0.67743	1.8467	1.5280	0.73123	-3.18
4.5	0.97540	2.8742	1.5219	0.862	-4.97
5.25	1.0904	3.8148	1.5022	0.90181	-5.68
6	1.1994	5.3708	1.4919	0.98248	-6.79
6.5	1.30	6.4989	1.5417	1.1356	-7.44
7	1.3558	9.7158	1.5175	1.2163	-8.31
8	1.4195	19.991	1.4981	1.3774	-9.85

Table 4. Polaron parameters obtained within the path-integral approach

α	$M_0^{(\mathrm{DSG})}$	$m^{*(\mathrm{Feynman})}$	$\tilde{M}_0^{(\mathrm{DSG})}$	$M_1^{(\mathrm{DSG})}/\alpha$	$E_0^{(\mathrm{Feynman})}$
0.01	0.00248	1.0017	1.5706	0.633	-0.010
1	0.24318	1.1957	1.5569	0.65468	-1.0130
3	0.69696	1.8912	1.5275	0.71572	-3.1333
4.5	1.0162	3.1202	1.5196	0.83184	-4.8394
5.25	1.1504	4.3969	1.5077	0.88595	-5.7482
6	1.2608	6.8367	1.4906	0.95384	-6.7108
6.5	1.3657	9.7449	1.5269	1.1192	-7.3920
7	1.4278	14.395	1.5369	1.2170	-8.1127
8	1.4741	31.569	1.5239	1.4340	-9.6953

The parameters corresponding to the Monte Carlo calculation are obtained using the numerical data kindly provided by A. Mishchenko. The comparison of the zero frequency moments $\tilde{M}_0^{(\mathrm{MC})}$ and $\tilde{M}_0^{(\mathrm{DSG})}$ with each other and with the value $\pi/2$ corresponding to the sum rule [80]

$$\frac{\pi}{2m^*} + \int_1^\infty \mathrm{Re}\,\sigma(\omega)\,d\omega = \frac{\pi}{2} \tag{4.123}$$

shows that $\left|\tilde{M}_0^{(\mathrm{MC})} - \tilde{M}_0^{(\mathrm{DSG})}\right|$ is smaller than each of the differences $\frac{\pi}{2} - \tilde{M}_0^{(\mathrm{MC})}$, $\frac{\pi}{2} - \tilde{M}_0^{(\mathrm{DSG})}$, which appear due to a finite interval of the integration in (4.120), (4.121).

We analyze also the fulfilment of the ground-state theorem [81]

$$E_0(\alpha) - E_0(0) = -\frac{3}{\pi}\int_0^\alpha \frac{d\alpha'}{\alpha'}\int_0^\infty \omega\,\mathrm{Re}\,\sigma(\omega,\alpha')\,d\omega \tag{4.124}$$

using the first-frequency moments $M_1^{(\mathrm{MC})}$ and $M_1^{(\mathrm{DSG})}$. The results of this comparison are presented in Fig. 10. The dots indicate the polaron ground-state energy calculated using the Feynman variational principle. The solid curve is the value of $E_0(\alpha)$ calculated numerically using the optical conductivity spectra and the ground-state theorem with the DSG optical conductivity [48] for a polaron,

$$E_0^{(\mathrm{DSG})}(\alpha) \equiv -\frac{3}{\pi}\int_0^\alpha \frac{d\alpha'}{\alpha'}\int_0^\infty \omega\,\mathrm{Re}\,\sigma^{(\mathrm{DSG})}(\omega,\alpha')\,d\omega. \tag{4.125}$$

The dashed and the dot-dashed curves are the values obtained using $M_1^{(\mathrm{DSG})}(\alpha)$ and $M_1^{(\mathrm{MC})}(\alpha)$, respectively:

$$\tilde{E}_0^{(\mathrm{DSG})}(\alpha) \equiv -\frac{3}{\pi}\int_0^{\alpha}\frac{d\alpha'}{\alpha'}\int_0^{\omega_{\max}}\omega\,\mathrm{Re}\,\sigma^{(\mathrm{DSG})}(\omega,\alpha')\,d\omega = -\frac{3}{\pi}\int_0^{\alpha}d\alpha'\frac{M_1^{(\mathrm{DSG})}(\alpha')}{\alpha'}, \quad (4.126)$$

$$\tilde{E}_0^{(\mathrm{MC})}(\alpha) \equiv -\frac{3}{\pi}\int_0^{\alpha}\frac{d\alpha'}{\alpha'}\int_0^{\omega_{\max}}\omega\,\mathrm{Re}\,\sigma^{(\mathrm{MC})}(\omega,\alpha')\,d\omega = -\frac{3}{\pi}\int_0^{\alpha}d\alpha'\frac{M_1^{(\mathrm{MC})}(\alpha')}{\alpha'}. \quad (4.127)$$

As seen from the figure, $E_0^{(\mathrm{DSG})}(\alpha)$ to a high degree of accuracy coincides with the variational polaron ground-state energy. Both $\tilde{E}_0^{(\mathrm{DSG})}(\alpha)$ and $\tilde{E}_0^{(\mathrm{MC})}(\alpha)$ differ from $E_0^{(\mathrm{DSG})}(\alpha)$ due to the integration over a finite interval of frequencies. However, $\tilde{E}_0^{(\mathrm{DSG})}(\alpha)$ and $\tilde{E}_0^{(\mathrm{MC})}(\alpha)$ are very close to each other. Herefrom, a conclusion follows that for integrals over the finite frequency region characteristic for the polaron optical absorption (i. e., except the "tails"), the function $\tilde{E}_0^{(\mathrm{MC})}(\alpha)$ (4.127) reproduces very well the function $\tilde{E}_0^{(\mathrm{DSG})}(\alpha)$.



FIG. 10: The ground-state theorem for a polaron using different data for the optical conductivity spectra, DSG from Ref. [48] and MC from Ref. [77]. The notations are explained in the text.

D. Scaling relations

1. *Derivation of the scaling relations*

The form of the Fröhlich Hamiltonian in n dimensions is the same as in 3D,

$$H = \frac{\mathbf{p}^2}{2m_b} + \sum_{\mathbf{k}} \hbar\omega_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \sum_{\mathbf{k}} \left(V_{\mathbf{k}} a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_{\mathbf{k}}^{*} a_{\mathbf{k}}^{\dagger} e^{-i\mathbf{k}\cdot\mathbf{r}} \right), \tag{4.128}$$

except that now all vectors are n-dimensional. In this subsection, dispersionless longitudinal phonons are considered, i.e., $\omega_{\mathbf{k}} = \omega_{\mathrm{LO}}$, and units are chosen such that $\hbar = m_b = \omega_{\mathrm{LO}} = 1$.

The electron-phonon interaction is a representation in second quantization of the electron interaction with the lattice polarization, which in 3D is essentially the Coulomb potential $1/r$. $|V_{\mathbf{k}}|^2$ is proportional to the Fourier transform of this potential, and as a consequence we have in n dimensions

$$|V_{\mathbf{k}}|^2 = \frac{A_n}{L_n k^{n-1}}, \tag{4.129}$$

where L_n is the volume of the n-dimensional crystal. Note that $|V_{\tilde{\mathbf{k}}}|^2$, where $\tilde{\mathbf{k}}$ is an $(n-1)$-dimensional vector, can be obtained from $|V_{\mathbf{k}}|^2$, where $\mathbf{k} = \left(\tilde{\mathbf{k}}, k_n\right)$ is an n-dimensional vector, by summing out one of the dimensions explicitly:

$$|V_{\tilde{\mathbf{k}}}|^2 = \sum_{k_n} |V_{\mathbf{k}}|^2 . \tag{4.130}$$

Inserting Eq. (4.129) into Eq. (4.130), we have

$$\frac{A_{n-1}}{L_{n-1}\tilde{k}^{n-2}} = \sum_{k_n} \frac{A_n}{L_n \left(\tilde{k}^2 + k_n^2\right)^{(n-1)/2}}. \tag{4.131}$$

Replacing the sum in Eq. (4.131) by an integral, i.e.,

$$\frac{L_{n-1}}{L_n} \sum_{k_n} \longrightarrow \frac{1}{2\pi} \int dk_n, \tag{4.132}$$

we obtain

$$\frac{A_{n-1}}{\tilde{k}^{n-2}} = \frac{A_n}{2\pi} \int_{-\infty}^{\infty} \frac{dk_n}{\left(\tilde{k}^2 + k_n^2\right)^{(n-1)/2}}. \tag{4.133}$$

Since

$$\int_{-\infty}^{\infty} \frac{dx}{(z^{\mu} + x^{\mu})^{\rho}} = z^{1-\mu\rho} \frac{\Gamma\left(\frac{1}{\mu}\right) \Gamma\left(\rho - \frac{1}{\mu}\right)}{\Gamma(\rho)}, \tag{4.134}$$

we have

$$\int_{-\infty}^{\infty} \frac{dk_n}{\left(\tilde{k}^2 + k_n^2\right)^{(n-1)/2}} = \frac{1}{\tilde{k}^{n-2}} \frac{\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{n-2}{2}\right)}{\Gamma\left(\frac{n-1}{2}\right)} = \frac{\sqrt{\pi}}{\tilde{k}^{n-2}} \frac{\Gamma\left(\frac{n-2}{2}\right)}{\Gamma\left(\frac{n-1}{2}\right)}, \tag{4.135}$$

where $\Gamma(x)$ is eh Γ function. Inserting Eq. (4.135) into Eq. (4.133), we obtain

$$A_n = \frac{2\sqrt{\pi}\Gamma\left(\frac{n-1}{2}\right)}{\Gamma\left(\frac{n-2}{2}\right)} A_{n-1}. \tag{4.136}$$

In 3D the interaction coefficient is well known, $\left|V_{\mathbf{k}}\right|^2 = 2\sqrt{2}\pi\alpha / L_3 k^2$, so that

$$A_3 = 2\sqrt{2}\pi\alpha. \tag{4.137}$$

Inserting Eq. (4.137) into Eq. (4.136), we immediately obtain

$$A_2 = 2\sqrt{2}\pi\alpha \frac{\Gamma\left(\frac{1}{2}\right)}{2\sqrt{\pi}\Gamma(1)} = \sqrt{2}\pi\alpha. \tag{4.138}$$

Applying Eq. (4.136) $n-2$ times, we further obtain for $n > 3$

$$\begin{aligned} A_n &= \frac{\left(2\sqrt{\pi}\right)^{n-2} \prod\limits_{j=2}^{n-1} \Gamma\left(\frac{j}{2}\right)}{\prod\limits_{j=1}^{n-2} \Gamma\left(\frac{j}{2}\right)} A_2 = \frac{\left(2\sqrt{\pi}\right)^{n-2} \left(\frac{n-1}{2}\right)}{\Gamma\left(\frac{1}{2}\right)} A_2 = 2^{n-2}\pi^{(n-3)/2}\Gamma\left(\frac{n-1}{2}\right) A_2 \\ &= 2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha. \end{aligned} \tag{4.139}$$

So, the interaction coefficient in n dimensions becomes [82]

$$\left|V_{\mathbf{k}}\right|^2 = \frac{2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{L_n k^{n-1}}. \tag{4.140}$$

Following the Feynman approach [41], the upper bound for the polaron ground-state energy can be written down as

$$E = E_0 - \lim_{\beta\to\infty} \frac{1}{\beta} \left\langle S - S_0 \right\rangle_0, \tag{4.141}$$

where S is the exact action functional of the polaron problem, while S_0 is the trial action functional, which corresponds to a model system where an electron is coupled by an elastic force to a fictitious particle (i.e., the model system describes a harmonic oscillator). E_0 is the ground-state energy of the above model system, and

$$\left\langle F \right\rangle_0 \equiv \frac{\int F e^{S_0} \mathcal{D}\mathbf{r}(t)}{\int e^{S_0} \mathcal{D}\mathbf{r}(t)}. \tag{4.142}$$

As indicated above, the Fröhlich Hamiltonian in n dimensions is the same as in 3D, except that now all vectors are n-dimensional [and the coupling coefficient $\left|V_{\mathbf{k}}\right|^2$ is modified in accordance with Eq. (4.140)]. Similarly, the only difference of the model system in n dimensions from the model system in 3D is that now one deals with an n-dimensional harmonic oscillator. So, directly following [41], one can represent $\lim_{\beta\to\infty}\left\langle S-S_0\right\rangle_0/\beta$ as

$$\lim_{\beta\to\infty}\frac{1}{\beta}\left\langle S-S_0\right\rangle_0 = A+B, \tag{4.143}$$

where

$$A=\sum_{\mathbf{k}}\left|V_{\mathbf{k}}\right|^2\int_0^\infty\left\langle e^{i\mathbf{k}\cdot[\mathbf{r}(t)-\mathbf{r}(0)]}\right\rangle_0 e^{-t}dt, \tag{4.144}$$

$$B=\frac{w\left(v^2-w^2\right)}{4}\int_0^\infty\left\langle\left[\mathbf{r}\left(t\right)-\mathbf{r}\left(0\right)\right]^2\right\rangle_0 e^{-wt}dt, \tag{4.145}$$

w and v are variational parameters, which should be determined by minimizing E of Eq. (4.141). Since the averaging $\langle ...\rangle_0$ in Eq. (4.144) is performed with the trial action, which corresponds to a harmonic oscillator, components of the electron coordinates, r_j $(j=1,...,n)$, in $\left\langle e^{i\mathbf{k}\cdot[\mathbf{r}(t)-\mathbf{r}(0)]}\right\rangle_0$ separate [41]:

$$\left\langle e^{i\mathbf{k}\cdot[\mathbf{r}(t)-\mathbf{r}(0)]}\right\rangle_0=\prod_{j=1}^n\left\langle e^{ik_j[r_j(t)-r_j(0)]}\right\rangle_0. \tag{4.146}$$

For the average $\left\langle e^{ik_j[r_j(t)-r_j(0)]}\right\rangle_0$, Feynman obtained [41]

$$\left\langle e^{ik_j[r_j(t)-r_j(0)]}\right\rangle_0=e^{-k_j^2D_0(t)}, \tag{4.147}$$

where

$$D_0\left(t\right)=\frac{w^2}{2v^2}t+\frac{v^2-w^2}{2v^3}\left(1-e^{-vt}\right). \tag{4.148}$$

Inserting Eq. (4.146) with Eq. (4.147) into Eq. (4.144), we obtain

$$A=\int_0^\infty e^{-t}dt\sum_{\mathbf{k}}\left|V_{\mathbf{k}}\right|^2e^{-k^2D_0(t)}. \tag{4.149}$$

Inserting expression (4.140) for $\left|V_{\mathbf{k}}\right|^2$ into Eq. (4.149) and replacing the sum over $\mathbf{k}$ by an integral [see (4.132)], we have

$$\begin{aligned}A&=2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha\int_0^\infty e^{-t}dt\int\frac{e^{-k^2D_0(t)}}{k^{n-1}}\frac{d\mathbf{k}}{\left(2\pi\right)^n}\\&=2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha\int_0^\infty e^{-t}dt\int d\Omega_n\int_0^\infty\frac{e^{-k^2D_0(t)}}{k^{n-1}}\frac{k^{n-1}dk}{\left(2\pi\right)^n},\end{aligned} \tag{4.150}$$

where $d\Omega_n$ is the elemental solid angle in n dimensions. Since the integrand in Eq. (4.150) depends only on the modulus k of $\mathbf{k}$, one have simply $\int d\Omega_n = \Omega_n$ with

$$\Omega_n = \frac{2\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)}. \tag{4.151}$$

So, we obtain for A the result

$$\begin{aligned} A &= \frac{2^{-1/2}\pi^{-1/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{\Gamma\left(\frac{n}{2}\right)} \int_0^\infty e^{-t}dt \int_0^\infty e^{-k^2 D_0(t)}dk = \frac{2^{-1/2}\pi^{-1/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{\Gamma\left(\frac{n}{2}\right)} \int_0^\infty \frac{\sqrt{\pi}e^{-t}}{2\sqrt{D_0(t)}}dt \\ &= \frac{2^{-3/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{\Gamma\left(\frac{n}{2}\right)} \int_0^\infty \frac{e^{-t}}{\sqrt{D_0(t)}}dt. \end{aligned} \tag{4.152}$$

Like in Ref. [41], B can be easily calculated by noticing that

$$\begin{aligned} \left\langle [\mathbf{r}(t) - \mathbf{r}(0)]^2 \right\rangle_0 &= \sum_{j=1}^{n} \left\langle [r_j(t) - r_j(0)]^2 \right\rangle_0 = \sum_{j=1}^{n} \left[-\frac{\partial^2}{\partial k_j^2} \left\langle e^{i\mathbf{k}\cdot[\mathbf{r}(t)-\mathbf{r}(0)]} \right\rangle_0 \right]\Bigg|_{\mathbf{k}=0} \\ &= \sum_{j=1}^{n} 2D_0(t) = 2nD_0(t), \end{aligned} \tag{4.153}$$

so that

$$\begin{aligned} B &= \frac{nw\left(v^2-w^2\right)}{2} \int_0^\infty D_0(t)\, e^{-wt}dt \\ &= \frac{nw\left(v^2-w^2\right)}{2} \int_0^\infty \left[\frac{w^2}{2v^2} t e^{-wt} + \frac{v^2-w^2}{2v^3}\left(e^{-wt} - e^{-(v+w)t}\right) \right] dt \\ &= \frac{nw\left(v^2-w^2\right)}{2} \left[\frac{w^2}{2v^2}\frac{1}{v^2} + \frac{v^2-w^2}{2v^3}\left(\frac{1}{w} - \frac{1}{v+w}\right) \right] \\ &= \frac{n\left(v^2-w^2\right)}{4v}. \end{aligned} \tag{4.154}$$

Inserting Eq. (4.140) with A and B, given by Eqs. (4.152) and (4.154), together with the ground-state energy of the model system [41] (an isotropic n-dimensional harmonic oscillator),

$$E_0 = \frac{n(v-w)}{2}, \tag{4.155}$$

into Eq. (4.141), we obtain

$$\begin{aligned} E &= \frac{n(v-w)}{2} - \frac{n\left(v^2-w^2\right)}{4v} - \frac{2^{-3/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{\Gamma\left(\frac{n}{2}\right)} \int_0^\infty \frac{e^{-t}}{\sqrt{D_0(t)}}dt \\ &= \frac{n(v-w)^2}{4v} - \frac{\Gamma\left(\frac{n-1}{2}\right)\alpha}{2\sqrt{2}\Gamma\left(\frac{n}{2}\right)} \int_0^\infty \frac{e^{-t}}{\sqrt{D_0(t)}}dt. \end{aligned} \tag{4.156}$$

In order to make easier a comparison of E for n dimensions with the Feynman result [41] for 3D,

$$E_{\mathrm{3D}}(\alpha)=\frac{3(v-w)^2}{4v}-\frac{1}{\sqrt{2\pi}}\alpha\int_0^\infty\frac{e^{-t}}{\sqrt{D_0(t)}}dt, \tag{4.157}$$

it is convenient to rewrite Eq. (4.156) in the form

$$E_{n\mathrm{D}}(\alpha)=\frac{n}{3}\left[\frac{3(v-w)^2}{4v}-\frac{1}{\sqrt{2\pi}}\frac{3\sqrt{\pi}\Gamma\left(\frac{n-1}{2}\right)}{2n\Gamma\left(\frac{n}{2}\right)}\alpha\int_0^\infty\frac{e^{-t}}{\sqrt{D_0(t)}}dt\right]. \tag{4.158}$$

It is worth recalling that the parameters w and v must be determined by minimizing E. Thus, in the case of Eq. (4.158) one has to minimize the expression in the square brackets. The only difference of this expression from the r.h.s. of Eq. (4.157) is that α is multiplied by the factor

$$a_n=\frac{3\sqrt{\pi}\Gamma\left(\frac{n-1}{2}\right)}{2n\Gamma\left(\frac{n}{2}\right)}. \tag{4.159}$$

This means that the minimizing parameters w and v in nD at a given α will be exactly the same as those calculated in 3D for the Fröhlich constant as large as $a_n\alpha$:

$$v_{n\mathrm{D}}(\alpha)=v_{\mathrm{3D}}(a_n\alpha),\ w_{n\mathrm{D}}(\alpha)=w_{\mathrm{3D}}(a_n\alpha). \tag{4.160}$$

Therefore, comparing Eq. (4.158) to Eq. (4.157), we obtain the scaling relation [82–84]

$$E_{n\mathrm{D}}(\alpha)=\frac{n}{3}E_{\mathrm{3D}}(a_n\alpha), \tag{4.161}$$

where a_n is given by Eq. (4.159). As discussed in Ref. [82], the above scaling relation is not an exact relation. It is valid for the Feynman polaron energy and also for the ground-state energy to order α. The next-order term (i.e., α^2) no longer satisfies Eq. (4.161). The reason is that in the exact calculation (to order α^2) the electron motion in the different space directions is coupled by the electron-phonon interaction. No such a coupling appears in the Feynman polaron model [see, e.g., Eq. (4.146)]; and this is the underlying reason for the validity of the scaling relation for the Feynman approximation.

In Refs. [82, 84], scaling relations are obtained also for the impedance function, the effective mass and the mobility of a polaron. The inverse of the impedance function $Z(\omega)$ is given by

$$\frac{1}{Z(\omega)}=\frac{i}{\omega-\Sigma(\omega)}, \tag{4.162}$$

where the memory function $\Sigma(\omega)$ can be expressed as [85]

$$\Sigma(z)=\frac{1}{z}\int_0^\infty dt\left(1-e^{izt}\right)\mathrm{Im}S(t), \tag{4.163}$$

with $z = \omega + i0^{+}$ and

$$S(t) = \sum_{\mathbf{k}} 2k_1^2 \left|V_{\mathbf{k}}\right|^2 e^{-k^2 D(t)} T(t), \tag{4.164}$$

$$T(t) = [1 + n(1)] e^{t} + n(1) e^{-it}, \tag{4.165}$$

$$D(t) = \frac{w^2}{2v^2}\left(-it + \frac{t^2}{\beta}\right) + \frac{v^2 - w^2}{2v^3}\left[1 - e^{-ivt} + 4n(v)\sin^2\left(\frac{vt}{2}\right)\right]. \tag{4.166}$$

Here, β is the inverse temperature and $n(\omega)$ is the occupation number of phonons with frequency ω (recall that in our units $\omega_{\mathrm{LO}} = 1$).

As implied from Eqs. (4.162) and (4.163), scaling of $\Sigma(\omega)$ and $Z(\omega)$ is determined by scaling of $S(t)$. For an isotropic crystal, since $\left|V_{\mathbf{k}}\right|^2$, $D(t)$ and $T(t)$ do not depend on the direction of $\mathbf{k}$, one can write $\sum_{\mathbf{k}} k_1^2 \left|V_{\mathbf{k}}\right|^2 e^{-k^2 D(t)} T(t) = \sum_{\mathbf{k}} k_2^2 \left|V_{\mathbf{k}}\right|^2 e^{-k^2 D(t)} T(t) = ... = \sum_{\mathbf{k}} k_n^2 \left|V_{\mathbf{k}}\right|^2 e^{-k^2 D(t)} T(t)$, so that

$$S(t) = \frac{2}{n}\sum_{\mathbf{k}} k^2 \left|V_{\mathbf{k}}\right|^2 e^{-k^2 D(t)} T(t). \tag{4.167}$$

Inserting expression (4.140) for $\left|V_{\mathbf{k}}\right|^2$ and replacing the sum over $\mathbf{k}$ by an integral, we have

$$\begin{aligned} S(t) &= \frac{2}{n}\frac{2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{(2\pi)^n}\int d\Omega_n \int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk \\ &= \frac{2}{n}\frac{2^{n-3/2}\pi^{(n-1)/2}\Gamma\left(\frac{n-1}{2}\right)\alpha}{(2\pi)^n}\frac{2\pi^{n/2}}{\Gamma\left(\frac{n}{2}\right)}\int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk \\ &= \sqrt{\frac{2}{\pi}}\frac{\Gamma\left(\frac{n-1}{2}\right)\alpha}{n\Gamma\left(\frac{n}{2}\right)}\int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk. \end{aligned} \tag{4.168}$$

In particular, for 3D one has from Eq. (4.168)

$$S_{\mathrm{3D}}(\alpha; t) = \frac{2\sqrt{2}}{3\pi}\alpha\int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk. \tag{4.169}$$

For nD, Eq. (4.168) can be rewritten is the form

$$\begin{aligned} S_{n\mathrm{D}}(\alpha; t) &= \frac{2\sqrt{2}}{3\pi}\frac{3\sqrt{\pi}\Gamma\left(\frac{n-1}{2}\right)}{2n\Gamma\left(\frac{n}{2}\right)}\alpha\int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk \\ &= \frac{2\sqrt{2}}{3\pi}a_n\alpha\int_0^{\infty} k^2 e^{-k^2 D(t)} T(t)\, dk. \end{aligned} \tag{4.170}$$

So, the only difference of the expression for $S_{\mathrm{3D}}(t)$ from $S_{\mathrm{3D}}(t)$ is that α is multiplied by a_n. Since for the minimizing parameters w and v, which enter $D(t)$, scaling is determined by the same product α with a_n [see Eq. (4.160)], we can write

$$S_{n\mathrm{D}}(\alpha; t) = S_{\mathrm{3D}}(a_n\alpha; t), \tag{4.171}$$

so that [84]

$$\Sigma_{n\mathrm{D}}(\alpha;\omega) = \Sigma_{3\mathrm{D}}(a_n\alpha;\omega), \tag{4.172}$$

and

$$Z_{n\mathrm{D}}(\alpha;\omega) = Z_{3\mathrm{D}}(a_n\alpha;\omega). \tag{4.173}$$

The polaron mass at zero temperature can be obtained from the impedance function as [85, 86]

$$\frac{m^*}{m_b} = 1 - \lim_{\omega\to 0}\frac{\mathrm{Re}\Sigma(\omega)}{\omega}, \tag{4.174}$$

so that from the scaling relation (4.172) for the memory function we also have a scaling relation for the polaron mass [84]:

$$\frac{m^*_{n\mathrm{D}}(\alpha)}{(m_b)_{n\mathrm{D}}} = \frac{m^*_{3\mathrm{D}}(a_n\alpha)}{(m_b)_{3\mathrm{D}}}. \tag{4.175}$$

Since the mobility can be obtained from the memory function as [87]

$$\frac{1}{\mu} = -\frac{m_b}{e}\lim_{\omega\to 0}\frac{\mathrm{Im}\Sigma(\omega)}{\omega}, \tag{4.176}$$

fulfilment of the scaling relation (4.172) implies also a scaling relation for the mobility [84]:

$$\mu_{n\mathrm{D}}(\alpha) = \mu_{3\mathrm{D}}(a_n\alpha). \tag{4.177}$$

In the important particular case of 2D, the above scaling relations take the form [82–84]:

$$E_{2\mathrm{D}}(\alpha) = \frac{2}{3}E_{3\mathrm{D}}\left(\frac{3\pi}{4}\alpha\right), \tag{4.178}$$

$$Z_{2\mathrm{D}}(\alpha;\omega) = Z_{3\mathrm{D}}\left(\frac{3\pi}{4}\alpha;\omega\right), \tag{4.179}$$

$$\frac{m^*_{2\mathrm{D}}(\alpha)}{(m_b)_{n\mathrm{D}}} = \frac{m^*_{3\mathrm{D}}\left(\frac{3\pi}{4}\alpha\right)}{(m_b)_{3\mathrm{D}}}, \tag{4.180}$$

$$\mu_{2\mathrm{D}}(\alpha) = \mu_{3\mathrm{D}}\left(\frac{3\pi}{4}\alpha\right). \tag{4.181}$$

2. Check of the scaling relation for the path integral Monte Carlo result for the polaron free energy

The fulfilment of the PD-scaling relation [84] is checked for the path integral Monte Carlo results [43] for the polaron free energy.

The path integral Monte Carlo results of Ref.[43] for the polaron free energy in 3D and in 2D are given for a few values of temperature and for some selected values of α. For a check of the scaling relation, the values of the polaron free energy at $\beta = 10$ are taken from Ref. [43] in 3D (Table I, for 4 values of α) and in 2D (Table II, for 2 values of α) and plotted in Fig. 11, upper panel, with squares and open circles, correspondingly.

The PD-scaling relation for the polaron ground-state energy as derived in Ref. [84] reads:

$$E_{2D}\left(\alpha\right) \equiv \frac{2}{3} E_{3D}\left(\frac{3\pi\alpha}{4}\right). \tag{4.182}$$

In Fig. 11, lower panel, the available data for the free energy from Ref [43] are plotted in the following form *inspired by the l.h.s. and the r.h.s parts of Eq. (1)*: $F_{2D}\left(\alpha\right)$ (squares) and $\frac{2}{3}F_{3D}\left(\frac{3\pi\alpha}{4}\right)$(open triangles). As follows from the figure, t*he path integral Monte Carlo results for the polaron free energy in 2D and 3D very closely follow the PD-scaling relation of the form given by Eq. (4.182):*

$$F_{2D}\left(\alpha\right) \equiv \frac{2}{3} F_{3D}\left(\frac{3\pi\alpha}{4}\right). \tag{4.183}$$



FIG. 11: *Upper panel:* The values of the polaron free energy in 3D (squares) and 2D (open circles) obtained by Ciuchi'2001 [43] for $\beta = 10$. The data for $F_{3D}(\alpha)$ are interpolated using a polynomial fit to the available four points (dotted line). *Lower panel:* Demonstration of the PD-scaling cf. PD'1987: the values of the polaron free energy in 2D obtained by Ciuchi'2001 [43] for $\beta = 10$ (squares) are very close to the **PD-scaled** according to PD'1987 [84] values of the polaron free energy in 3D from Ciuchi'2001 for $\beta = 10$ (open triangles). The data for $\frac{2}{3}F_{3D}\left(\frac{3\pi\alpha}{4}\right)$ are interpolated using a polynomial fit to the available four points (solid line).

Appendix 1. Weak coupling: LLP approach

Inspired by the work of Tomonaga on quantum electrodynamics (Q. E. D.), Lee, Low and Pines (L.L.P.) [29] derived (1.22) and $m^* = m_b(1 + \alpha/6)$ from a canonical transformation formulation, which establishes (1.22) as a variational upper bound for the ground-state energy.

The wave equation corresponding to the Fröhlich Hamiltonian (1.16) is

$$H\Phi = E\Phi. \tag{4.184}$$

We shall take advantage of the fact that the total momentum of the system

$$\mathbf{P}_{op} = \sum_{\mathbf{k}} \hbar \mathbf{k} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}} + \mathbf{p} \tag{4.185}$$

(where $\mathbf{p} = -i\hbar\nabla$ is the momentum of the electron) is a constant of motion because it commutes with the Hamiltonian (1.16)

Indeed,

$$\begin{aligned}
[\mathbf{p}, H] &= [\mathbf{p}, \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}})] = \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} \left[\mathbf{p}, e^{i\mathbf{k}\cdot\mathbf{r}}\right] + V_k^* a^{\dagger}_{\mathbf{k}} \left[\mathbf{p}, e^{-i\mathbf{k}\cdot\mathbf{r}}\right]) \\
&= \sum_{\mathbf{k}} \hbar \mathbf{k} (V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a^{\dagger}_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}});
\end{aligned}$$

$$\begin{aligned}
\left[\sum_{\mathbf{k}} \hbar \mathbf{k} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}}, H\right] &= \left[\sum_{\mathbf{k}} \hbar \mathbf{k} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}}, \sum_{\mathbf{k}'} (V_{k'} a_{\mathbf{k}'} e^{i\mathbf{k}'\cdot\mathbf{r}} + V_{k'}^* a^{\dagger}_{\mathbf{k}'} e^{-i\mathbf{k}'\cdot\mathbf{r}})\right] = \\
&= \sum_{\mathbf{k}} \hbar \mathbf{k} \left[a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}}, (V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* a^{\dagger}_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}})\right] = \\
&= \sum_{\mathbf{k}} \hbar \mathbf{k} \left\{V_k \left[a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}}, a_{\mathbf{k}}\right] e^{i\mathbf{k}\cdot\mathbf{r}} + V_k^* \left[a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}}, a^{\dagger}_{\mathbf{k}}\right] e^{-i\mathbf{k}\cdot\mathbf{r}}\right\} = \\
&= -\sum_{\mathbf{k}} \hbar \mathbf{k} \left(V_k a_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}} - V_k^* a^{\dagger}_{\mathbf{k}} e^{-i\mathbf{k}\cdot\mathbf{r}}\right);
\end{aligned}$$

$$[\mathbf{P}_{op}, H] = \left[\sum_{\mathbf{k}} \hbar \mathbf{k} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}} + \mathbf{p}, H\right] = 0. \tag{4.186}$$

Because of the commutation (4.186), the operators H and $\mathbf{P}_{op}$ have a common set of basis functions: $H\Phi = E\Phi$ and $\mathbf{P}_{op}\Phi = \mathbf{P}\Phi$.

It is possible to transform to a representation in which $\mathbf{P}_{op}$ becomes a "c" number $\mathbf{P}$, and in which the Hamiltonian no longer contains the electron coordinates. The unitary (canonical) transformation required is $\Phi = S_1\psi$, where

$$S_1 = \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a^{\dagger}_{\mathbf{k}} a_{\mathbf{k}})\cdot\mathbf{r}\right]. \tag{4.187}$$

Derivation of the transformations of the operators.

$$\begin{aligned}
\mathbf{p} \longrightarrow & S_1^{-1}\mathbf{p}S_1 = \\
&= \exp\left[-\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \mathbf{p} \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \\
&= \exp\left[-\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] (-i\hbar\nabla) \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \\
&= \exp\left[-\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \left\{ \begin{array}{c} (\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}) \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \\ + \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] (-i\hbar\nabla) \end{array} \right\} \\
&= \mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \mathbf{p}, \qquad (4.188)
\end{aligned}$$

$$\begin{aligned}
\mathbf{P}_{op} \longrightarrow & S_1^{-1}\mathbf{P}_{op}S_1 = \\
&= \exp\left[-\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \left(\sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \mathbf{p}\right) \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}})\cdot\mathbf{r}\right] \\
&= \exp\left[\frac{i}{\hbar} \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}\cdot\mathbf{r}\right] \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} \exp\left[-\frac{i}{\hbar} \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}\cdot\mathbf{r}\right] + S_1^{-1}\mathbf{p}S_1 \\
&= \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \mathbf{P} - \sum_{\mathbf{k}} \hbar\mathbf{k}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}} + \mathbf{p} = \mathbf{P} + \mathbf{p}, \qquad (4.189)
\end{aligned}$$

$$
\begin{aligned}
a_{\mathbf{k}} &\longrightarrow S_1^{-1} a_{\mathbf{k}} S_1 = \\
&= \exp\left[-\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}) \cdot \mathbf{r}\right] a_{\mathbf{k}} \exp\left[\frac{i}{\hbar}(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}) \cdot \mathbf{r}\right] \\
&= \exp\left[\frac{i}{\hbar} \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] a_{\mathbf{k}} \exp\left[-\frac{i}{\hbar} \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] \\
&= \exp\left[i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] a_{\mathbf{k}} \exp\left[-i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] \\
&= \exp\left[i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] a_{\mathbf{k}} \sum_{n=0}^{\infty} \frac{1}{n!} \left(-i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right)^n \\
&= \exp\left[i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right] \sum_{n=0}^{\infty} \frac{1}{n!} a_{\mathbf{k}} \left(-i \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} \cdot \mathbf{r}\right)^n \\
&= \exp\left[i \mathbf{k} \cdot \mathbf{r} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right] \sum_{n=0}^{\infty} \frac{1}{n!} \left(-i \mathbf{k} \cdot \mathbf{r}\right)^n a_{\mathbf{k}} \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right)^n \overset{\text{see}(*)}{=} \\
&= \exp\left[i \mathbf{k} \cdot \mathbf{r} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right] \sum_{n=0}^{\infty} \frac{1}{n!} [-i \mathbf{k} \cdot \mathbf{r} \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + 1\right)]^n a_{\mathbf{k}} \\
&= \exp\left[i \mathbf{k} \cdot \mathbf{r} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right] \exp\left[-i \mathbf{k} \cdot \mathbf{r} (a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + 1)\right] a_{\mathbf{k}} \\
&= a_{\mathbf{k}} \exp\left(-i \mathbf{k} \cdot \mathbf{r}\right). \qquad (4.190)
\end{aligned}
$$

Here the property was used:

$$
a_{\mathbf{k}} \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right)^n = \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + 1\right)^n a_{\mathbf{k}}. \qquad (*)
$$

It is evident for $n = 0$.For $n = 1$ it is demonstrated as follows:

$$
a_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} = a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + 1 \Longrightarrow a_{\mathbf{k}} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} = (a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + 1) a_{\mathbf{k}};
$$

then for $n \geq 2$ the validity of (*) is straightforwardly demonstrated by induction.

Finally,

$$
a_{\mathbf{k}}^{\dagger} \longrightarrow S_1^{-1} a_{\mathbf{k}}^{\dagger} S_1 = [S_1^{-1} a_{\mathbf{k}} S_1]^{\dagger} = a_{\mathbf{k}}^{\dagger} \exp\left(i \mathbf{k} \cdot \mathbf{r}\right). \qquad (4.191)
$$

Using (4.188), (4.189), (4.190) and (4.191), one arrives at

$$
H \longrightarrow \mathcal{H} = S_1^{-1} H S_1 = \frac{\left(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}\right)^2}{2 m_b} + \sum_{\mathbf{k}} \hbar \omega_{\mathrm{LO}} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \sum_{\mathbf{k}} (V_k a_{\mathbf{k}} + V_k^* a_{\mathbf{k}}^{\dagger}), \qquad (4.192)
$$

where $\mathbf{p}$ is set 0. [2] The wave equation (4.184) takes the form

$$HS_1\psi = ES_1\psi \Longrightarrow \mathcal{H}\psi = E\psi. \tag{4.193}$$

Our aim is to calculate for a given momentum $\mathbf{P}$ the lowest eigenvalue $E(P)$ of the Hamiltonian (4.192). For the low-lying energy levels of the electron $E(P)$ may be well represented by the first two terms of a power series expansion in P^2 : $E(P) = E_0 + P^2/2m_p + O(P^4)$,where m_p is the effective mass of the polaron.

The canonical transformation (4.187) formally eliminates the electron operators from the Hamiltonian. LLP use further a variational method of calculation. The trial wave function is chosen as

$$\psi = S_2\psi_0 \tag{4.194}$$

where ψ_0 is the eigenstate of the unperturbed Hamiltonian with no phonons present (vacuum state). Specifically, ψ_0 is defined by

$$a_{\mathbf{k}}\psi_0 = 0, \qquad (\psi_0, \psi_0) = 1 \tag{4.195}$$

and the second canonical transformation:

$$S_2 = \exp\left[\sum_{\mathbf{k}}(a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right], \tag{4.196}$$

where $f_{\mathbf{k}}$ are treated as variational functions and will be chosen to minimize the energy. The physical significance of Eq. (4.196) is that it "dresses" the electron with the virtual phonon field, which describes the polarization. Viewed as a unitary transformation, S_2 is a *displacement* operator on $a_{\mathbf{k}}$ and $a_{\mathbf{k}}^{\dagger}$:

---

[2] Transformation of the equation $\mathbf{P}_{op}\Phi = \mathbf{P}\Phi$ leads to $S_1^{-1}\mathbf{P}_{op}S_1\psi = \mathbf{P}\psi$.At the same time, applying Eq. (4.189), we obtain $S_1^{-1}\mathbf{P}_{op}S_1 = \mathbf{P} + \mathbf{p}$.Setting the gauge $\mathbf{p}\psi = 0$ eliminates the operator $\mathbf{p}$ from the problem.

$$\begin{aligned}
a_{\mathbf{k}} &\longrightarrow S_2^{-1} a_{\mathbf{k}} S_2 = \\
&= \exp\left[-\sum_{\mathbf{k}} (a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] a_{\mathbf{k}} \exp\left[\sum_{\mathbf{k}} (a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] \\
&= \exp\left[-(a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] a_{\mathbf{k}} \exp\left[(a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] \overset{\text{see}(**)}{=} \\
&= a_{\mathbf{k}} + \left[a_{\mathbf{k}}, (a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] + \frac{1}{2}\left[\left[a_{\mathbf{k}}, (a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right], (a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} - a_{\mathbf{k}} f_{\mathbf{k}}^{*})\right] + ... \\
&= a_{\mathbf{k}} + f_{\mathbf{k}}, \qquad (4.197)
\end{aligned}$$

$$a_{\mathbf{k}}^{\dagger} \longrightarrow S_2^{-1} a\dagger_{\mathbf{k}} S_2 = a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^{*}. \qquad (4.198)$$

Here the relation was used

$$\exp[-V]\, a \exp[V] = a + [a, V] + \frac{1}{2}\left[[a, V], V\right] + \frac{1}{3!}\left[\left[[a, V], V\right], V\right] + ... \qquad (**)$$

Further we seek to minimize the expression for the energy,

$$E = (\psi, H\psi) = \left(\psi_0, S_2^{-1} \mathcal{H} S_2 \psi_0\right). \qquad (4.199)$$

In virtue of (4.197) and (4.198), we obtain:

$$\begin{aligned}
S_2^{-1} \mathcal{H} S_2 &= \frac{\left[\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} \left(a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^{*}\right)(a_{\mathbf{k}} + f_{\mathbf{k}})\right]^2}{2m_b} \\
&\quad + \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} \left(a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^{*}\right)(a_{\mathbf{k}} + f_{\mathbf{k}}) + \sum_{\mathbf{k}} \left[V_k (a_{\mathbf{k}} + f_{\mathbf{k}}) + V_k^{*} \left(a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^{*}\right)\right] \\
&= \frac{\left[(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}}) - \sum_{\mathbf{k}} \hbar \mathbf{k} \left|f_{\mathbf{k}}\right|^2 - \sum_{\mathbf{k}} \hbar \mathbf{k} \left(a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} + a_{\mathbf{k}} f_{\mathbf{k}}^{*}\right)\right]^2}{2m_b} \\
&\quad + \sum_{\mathbf{k}} \hbar\omega_{\mathrm{LO}} \left(a_{\mathbf{k}}^{\dagger} a_{\mathbf{k}} + \left|f_{\mathbf{k}}\right|^2 + a_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} + a_{\mathbf{k}} f_{\mathbf{k}}^{*}\right) \\
&\quad + \sum_{\mathbf{k}} \left[V_k (a_{\mathbf{k}} + f_{\mathbf{k}}) + V_k^{*} \left(a_{\mathbf{k}}^{\dagger} + f_{\mathbf{k}}^{*}\right)\right] \\
&= H_0 + H_1,
\end{aligned}$$

where

$$
\begin{aligned}
H_0 = {} & \frac{\left[\left(\mathbf{P} - \sum_{\mathbf{k}} \hbar \mathbf{k} a^\dagger_{\mathbf{k}} a_{\mathbf{k}} \right) \right]^2 + \left[\sum_{\mathbf{k}} \hbar \mathbf{k} \left| f_{\mathbf{k}} \right|^2 \right]^2}{2 m_b} + \sum_{\mathbf{k}} \left[V_k f_{\mathbf{k}} + V_k^* f^*_{\mathbf{k}} \right] \\
& + \sum_{\mathbf{k}} \left| f_{\mathbf{k}} \right|^2 \left\{ \hbar \omega_{\mathrm{LO}} - \frac{\hbar \mathbf{k} \cdot \mathbf{P}}{m_b} + \frac{\hbar^2 k^2}{2 m_b} \right\} + \frac{\hbar^2}{m_b} \sum_{\mathbf{k}} \mathbf{k} a^\dagger_{\mathbf{k}} a_{\mathbf{k}} \cdot \sum_{\mathbf{k}'} \mathbf{k}' \left| f_{\mathbf{k}'} \right|^2 \\
& + \sum_{\mathbf{k}} a_{\mathbf{k}} \left\{ V_k + f^*_{\mathbf{k}} \left[\hbar \omega_{\mathrm{LO}} - \frac{\hbar \mathbf{k} \cdot \mathbf{P}}{m_b} + \frac{\hbar^2 k^2}{2 m_b} + \frac{\hbar^2 \mathbf{k}}{m_b} \cdot \sum_{\mathbf{k}'} \mathbf{k}' \left| f_{\mathbf{k}'} \right|^2 \right] \right\} \\
& + \sum_{\mathbf{k}} a^\dagger_{\mathbf{k}} \left\{ V_k^* + f_{\mathbf{k}} \left[\hbar \omega_{\mathrm{LO}} - \frac{\hbar \mathbf{k} \cdot \mathbf{P}}{m_b} + \frac{\hbar^2 k^2}{2 m_b} + \frac{\hbar^2 \mathbf{k}}{m_b} \cdot \sum_{\mathbf{k}'} \mathbf{k}' \left| f_{\mathbf{k}'} \right|^2 \right] \right\} \\
& + \sum_{\mathbf{k}} \hbar \omega_{\mathrm{LO}} a^\dagger_{\mathbf{k}} a_{\mathbf{k}}; \qquad (4.200)
\end{aligned}
$$

$$
\begin{aligned}
H_1 = {} & \sum_{\mathbf{k},\mathbf{k}'} \frac{\hbar^2 \mathbf{k} \cdot \mathbf{k}'}{2 m_b} \left\{ a_{\mathbf{k}} a_{\mathbf{k}'} f^*_{\mathbf{k}} f^*_{\mathbf{k}'} + 2 a^\dagger_{\mathbf{k}} a_{\mathbf{k}'} f_{\mathbf{k}} f^*_{\mathbf{k}'} + a^\dagger_{\mathbf{k}} a^\dagger_{\mathbf{k}'} f_{\mathbf{k}} f_{\mathbf{k}'} \right\} + \\
& + \sum_{\mathbf{k},\mathbf{k}'} \frac{\hbar^2 \mathbf{k} \cdot \mathbf{k}'}{2 m_b} \left\{ a^\dagger_{\mathbf{k}} a_{\mathbf{k}} a_{\mathbf{k}'} f^*_{\mathbf{k}'} + a^\dagger_{\mathbf{k}'} a^\dagger_{\mathbf{k}} a_{\mathbf{k}} f_{\mathbf{k}'} \right\}.
\end{aligned}
$$

Using (4.195), we obtain from (4.199) that

$$
\begin{aligned}
E = H_0 = {} & \frac{P^2 + \left[\sum_{\mathbf{k}} \hbar \mathbf{k} \left| f_{\mathbf{k}} \right|^2 \right]^2}{2 m_b} + \sum_{\mathbf{k}} \left[V_k f_{\mathbf{k}} + V_k^* f^*_{\mathbf{k}} \right] \\
& + \sum_{\mathbf{k}} \left| f_{\mathbf{k}} \right|^2 \left\{ \hbar \omega_{\mathrm{LO}} - \frac{\hbar \mathbf{k} \cdot \mathbf{P}}{m_b} + \frac{\hbar^2 k^2}{2 m_b} \right\}. \qquad (4.201)
\end{aligned}
$$

We minimize (4.201) by imposing

$$
\frac{\delta E}{\delta f_{\mathbf{k}}} = 0, \qquad \frac{\delta E}{\delta f^\dagger_{\mathbf{k}}} = 0.
$$

This results in

$$
V_k + f^*_{\mathbf{k}} \left\{ \hbar \omega_{\mathrm{LO}} - \frac{\hbar \mathbf{k} \cdot \mathbf{P}}{m_b} + \frac{\hbar^2 k^2}{2 m_b} + \frac{\hbar^2}{m_b} \left[\sum_{\mathbf{k}'} \hbar \mathbf{k}' \left| f_{\mathbf{k}'} \right|^2 \right] \cdot \mathbf{k} \right\} = 0 \qquad (4.202)
$$

and the appropriate complex conjugate equation for $f_{\mathbf{k}}$. Upon comparing (4.202) and (4.200), we see that the linear terms in $a^\dagger_{\mathbf{k}}$ and $a_{\mathbf{k}}$ are identically zero if (4.202) is satisfied, and hence that H_0 is diagonal in a representation in which $a^\dagger_{\mathbf{k}} a_{\mathbf{k}}$ is diagonal. So, the variational calculation by LLP is equivalent to the use of (4.200) as the total Hamiltonian provided $f^*_{\mathbf{k}}$

satisfies (4.202). An estimate of the accuracy of the LLP variational procedure was obtained by an estimate of the effect of H_1using a perturbation theory.

Now we evaluate the energy of the ground state of the system, which is given by Eq. (4.201) with $f^*_{\mathbf{k}}$ satisfying Eq. (4.202). The only preferred direction in this problem is $\mathbf{P}$. Therefore one may introduce the parameter η defined as

$$\eta\mathbf{P} = \sum_{\mathbf{k}} \hbar\mathbf{k} \left| f_{\mathbf{k}} \right|^2 . \tag{4.203}$$

Then Eq. (4.202) leads to

$$f^*_{\mathbf{k}} = -V_k \Bigg/ \left[\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b} \right] , \tag{4.204}$$

and we obtain the following implicit equation for η:

$$\begin{aligned} \eta\mathbf{P} &= \sum_{\mathbf{k}} \hbar\mathbf{k} \left| V_k \right|^2 \Bigg/ \left[\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b} \right]^2 \\ &= \frac{V}{(2\pi)^3} \int d^3k \hbar\mathbf{k} \left(\frac{\hbar\omega_{\mathrm{LO}}}{k} \right)^2 \frac{4\pi\alpha}{V} \left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}} \right)^{\frac{1}{2}} \Bigg/ \left[\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b} \right]^2 . \end{aligned}$$

Let us introduce spherical coordinates with a polar axis along $\mathbf{P}$ and denote $x = \cos(\mathbf{k}\hat{\ }\mathbf{P})$:

$$\begin{aligned} \eta P &= \frac{\alpha\hbar^3\omega_{\mathrm{LO}}^2}{2\pi^2} \left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}} \right)^{\frac{1}{2}} 2\pi \int_{-1}^{1} dx x \int_0^\infty dk k \Bigg/ \left[\hbar\omega_{\mathrm{LO}} - \frac{\hbar k P x}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b} \right]^2 \\ &= \frac{\alpha\hbar}{2\pi^2} \left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}} \right)^{\frac{1}{2}} 2\pi \int_{-1}^{1} dx x \int_0^\infty dk k \Bigg/ \left[1 - 2\frac{\hbar k P x}{2m_b\hbar\omega_{\mathrm{LO}}}(1-\eta) + \frac{\hbar^2 k^2}{2m_b\hbar\omega_{\mathrm{LO}}} \right]^2 . \end{aligned}$$

Further, we introduce the parameter

$$q = \frac{P}{(2m_b\hbar\omega_{\mathrm{LO}})^{1/2}}(1-\eta) \tag{4.205}$$

and a new variable

$$\kappa = \frac{\hbar k}{(2m_b\hbar\omega_{\mathrm{LO}})^{1/2}}.$$

This gives

$$
\begin{aligned}
\eta &= \frac{\alpha\hbar}{\pi}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\frac{2m_b\hbar\omega_{\mathrm{LO}}}{\hbar^2P}\int_{-1}^{1}dxx\int_0^{\infty}d\kappa\kappa\Big/\left[1-2q\kappa x+\kappa^2\right]^2\\
&= \frac{\alpha}{\pi}\frac{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}{P}\int_{-1}^{1}dxx\int_0^{\infty}d\kappa\kappa\Big/\left[(\kappa-qx)^2+(1-q^2x^2)\right]^2\\
&= \frac{\alpha}{\pi}\frac{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}{P}\int_{-1}^{1}dxx\int_{-qx}^{\infty}d\kappa\left(\kappa+qx\right)\Big/\left[\kappa^2+(1-q^2x^2)\right]^2\\
&= \frac{\alpha}{\pi}\frac{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}{P}\int_{-1}^{1}dxx\left\{\begin{array}{r} -\frac{1}{2[\kappa^2+(1-q^2x^2)]} \\ +qx\left[\frac{\kappa}{2(1-q^2x^2)[\kappa^2+(1-q^2x^2)]}+\frac{1}{2(1-q^2x^2)^{3/2}}\arctan\left(\frac{\kappa}{[1-q^2x^2]^{1/2}}\right)\right]\end{array}\right\}_{-qx}^{\infty}\\
&= \frac{\alpha}{\pi}\frac{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}{P}\int_{-1}^{1}dxx\left\{\frac{1}{2}+\frac{qx\pi}{4(1-q^2x^2)^{3/2}}+\frac{q^2x^2}{2(1-q^2x^2)}+\frac{qx}{2(1-q^2x^2)^{3/2}}\arcsin\left(qx\right)\right\}.
\end{aligned}
$$

$$\Downarrow$$

$$
\begin{aligned}
\eta &= \frac{\alpha}{\pi}\frac{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}{P}\frac{q\pi}{4}\int_{-1}^{1}\frac{x^2}{(1-q^2x^2)^{3/2}}dx\\
&= \frac{\alpha}{4}\left(1-\eta\right)\int_{-1}^{1}\frac{x^2}{(1-q^2x^2)^{3/2}}dx\\
&= \frac{\alpha}{2}\left(1-\eta\right)\frac{q-\sqrt{1-q^2}\arcsin\left(q\right)}{q^3\sqrt{1-q^2}}\\
&= \frac{\alpha\left(1-\eta\right)}{2\left(\frac{P}{\left(2m_b\hbar\omega_{\mathrm{LO}}\right)^{1/2}}(1-\eta)\right)^3}\left(\frac{q}{\sqrt{1-q^2}}-\arcsin\left(q\right)\right)\\
&= \frac{\alpha}{2\left(1-\eta\right)^2}\left(\frac{2m_b\hbar\omega_{\mathrm{LO}}}{P^2}\right)^{3/2}\left(\frac{q}{\sqrt{1-q^2}}-\arcsin\left(q\right)\right)
\end{aligned}
$$

So, we have arrived at the equation

$$\eta\left(1-\eta\right)^2=\frac{\alpha}{2}\left(\frac{2m_b\hbar\omega_{\mathrm{LO}}}{P^2}\right)^{3/2}\left(\frac{q}{\sqrt{1-q^2}}-\arcsin\left(q\right)\right), \tag{4.206}$$

or equivalently, using the definition (4.205),

$$\frac{\eta}{1-\eta}=\frac{\alpha}{2q^3}\left(\frac{q}{\sqrt{1-q^2}}-\arcsin\left(q\right)\right). \tag{4.207}$$

Using Eqs. (4.203) and (4.204), we can simplify the energy (4.201) as follows:

$$
\begin{aligned}
E = \frac{P^2+\left(\eta\mathbf{P}\right)^2}{2m_b} &- 2\sum_{\mathbf{k}}\frac{\left|V_k\right|^2}{\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}}\\
&+\sum_{\mathbf{k}}\frac{\left|V_k\right|^2}{\left(\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}\right)^2}\left(\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}+\frac{\hbar^2k^2}{2m_b}\right)
\end{aligned}
$$

$$= \frac{(1+\eta^2)\, P^2}{2m_b} - 2\sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}$$

$$+ \sum_{\mathbf{k}} \frac{|V_k|^2}{\left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}\right)^2} \left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta+\eta) + \frac{\hbar^2k^2}{2m_b}\right)$$

$$= \frac{(1+\eta^2)\, P^2}{2m_b} - 2\sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}$$

$$+ \sum_{\mathbf{k}} \frac{|V_k|^2}{\left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}\right)^2} \left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}\right)$$

$$- \sum_{\mathbf{k}} \frac{|V_k|^2}{\left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}\right)^2} \left(\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}\eta\right)$$

$$= \frac{(1+\eta^2)\, P^2}{2m_b} - 2\sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}$$

$$+ \sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}$$

$$- \left(\frac{\mathbf{P}}{m_b}\eta\right) \cdot \sum_{\mathbf{k}} \frac{\hbar\mathbf{k}\,|V_k|^2}{\left(\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}\right)^2}$$

$$= \frac{(1+\eta^2)\, P^2}{2m_b} - \sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}} - \left(\frac{\mathbf{P}}{m_b}\eta\right) \cdot \sum_{\mathbf{k}} \hbar\mathbf{k}\,|f_{\mathbf{k}}|^2$$

$$= \frac{(1+\eta^2)\, P^2}{2m_b} - \left(\frac{\mathbf{P}}{m_b}\eta\right) \cdot \eta\mathbf{P} - \sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}$$

$$\Downarrow$$

$$E = \frac{P^2}{2m_b}\left(1-\eta^2\right) - \sum_{\mathbf{k}} \frac{|V_k|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2k^2}{2m_b}}. \tag{4.208}$$

The sum over $\mathbf{k}$ in Eq. (4.208) is calculated as follows:

$$\begin{aligned}
&\sum_{\mathbf{k}}\frac{\left|V_k\right|^2}{\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}}\\
&=\frac{V}{(2\pi)^3}\int d\mathbf{k}\frac{\left(\frac{\hbar\omega_{\mathrm{LO}}}{k}\left(\frac{4\pi\alpha}{V}\right)^{\frac{1}{2}}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{4}}\right)^2}{\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}}\\
&=\frac{V}{(2\pi)^3}\hbar^2\omega_{\mathrm{LO}}^2\left(\frac{4\pi\alpha}{V}\right)\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int d\mathbf{k}\frac{1}{k^2\left[\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}\right]}\\
&=\frac{m_b\omega_{\mathrm{LO}}^2\alpha}{\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int d\mathbf{k}\frac{1}{k^2\left(k^2-2\frac{\mathbf{k}\cdot\mathbf{P}}{\hbar}(1-\eta)+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}\right)}.
\end{aligned}$$

For the calculation of this integral, we can use the auxiliary identity

$$\frac{1}{ab}=\int_0^1\frac{1}{\left[ax+b\left(1-x\right)\right]^2}. \tag{4.209}$$

Setting

$$\begin{aligned}
a&=k^2-2\frac{\mathbf{k}\cdot\mathbf{P}}{\hbar}(1-\eta)+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar},\\
b&=k^2,
\end{aligned}$$

we find

$$\begin{aligned}
&\sum_{\mathbf{k}}\frac{\left|V_k\right|^2}{\hbar\omega_{\mathrm{LO}}-\frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta)+\frac{\hbar^2k^2}{2m_b}}\\
&=\frac{m_b\omega_{\mathrm{LO}}^2\alpha}{\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int_0^1dx\int d\mathbf{k}\frac{1}{\left[x\left(k^2-2\frac{\mathbf{k}\cdot\mathbf{P}}{\hbar}(1-\eta)+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}\right)+\left(1-x\right)k^2\right]^2}\\
&=\frac{m_b\omega_{\mathrm{LO}}^2\alpha}{\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int_0^1dx\int d\mathbf{k}\frac{1}{\left(k^2-2\frac{\mathbf{k}\cdot\mathbf{P}}{\hbar}(1-\eta)x+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}x\right)^2}\\
&=\frac{m_b\omega_{\mathrm{LO}}^2\alpha}{\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int_0^1dx\int d\mathbf{k}\frac{1}{\left(\left(\mathbf{k}-\frac{\mathbf{P}}{\hbar}(1-\eta)x\right)^2+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}x-\frac{P^2}{\hbar^2}(1-\eta)^2x^2\right)^2}\\
&=\frac{m_b\omega_{\mathrm{LO}}^2\alpha}{\pi^2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}}\int_0^1dx\int d\mathbf{k}\frac{1}{\left(k^2+\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}x-\frac{P^2}{\hbar^2}(1-\eta)^2x^2\right)^2}
\end{aligned}$$

As long as $P^2/\left(2m_b\right)$ is sufficiently small so that no spontaneous emission can occur (roughly $P^2/\left(2m_b\right)\lesssim\hbar\omega_{\mathrm{LO}}$), the quantity

$$A\equiv\frac{2m_b\omega_{\mathrm{LO}}}{\hbar}x-\frac{P^2}{\hbar^2}(1-\eta)^2x^2$$

is supposed to be positive for $0 < x < 1$. Therefore, we can use the integral

$$\int \frac{1}{\left(k^2+A\right)^2} d\mathbf{k} = \frac{\pi^2}{\sqrt{A}},$$

what gives

$$\begin{aligned}
&\sum_{\mathbf{k}} \frac{\left|V_k\right|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b}} \\
&= \frac{m_b \omega_{\mathrm{LO}}^2 \alpha}{\pi^2} \left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{\frac{1}{2}} \int_0^1 dx \frac{\pi^2}{\sqrt{\frac{2m_b\omega_{\mathrm{LO}}}{\hbar} x - \frac{P^2}{\hbar^2}(1-\eta)^2 x^2}} \\
&= \frac{1}{2}\alpha\hbar\omega_{\mathrm{LO}} \int_0^1 dx \frac{1}{\sqrt{x - \frac{(1-\eta)^2 P^2}{2m_b\hbar\omega_{\mathrm{LO}}} x^2}} \\
&= \frac{1}{2}\alpha\hbar\omega_{\mathrm{LO}} \int_0^1 dx \frac{1}{\sqrt{x - q^2 x^2}}.
\end{aligned}$$

We change the variable $x = t^2$, what gives

$$\int_0^1 \frac{1}{\sqrt{x - q^2 x^2}} dx = 2\int_0^1 \frac{1}{\sqrt{1 - q^2 t^2}} dt = \frac{2}{q} \arcsin q,$$

and hence

$$\sum_{\mathbf{k}} \frac{\left|V_k\right|^2}{\hbar\omega_{\mathrm{LO}} - \frac{\hbar\mathbf{k}\cdot\mathbf{P}}{m_b}(1-\eta) + \frac{\hbar^2 k^2}{2m_b}} = \frac{\alpha\hbar\omega_{\mathrm{LO}}}{q} \arcsin q. \tag{4.210}$$

As a result, the energy (4.208) is expressed in a closed form

$$E = \frac{P^2}{2m_b}\left(1-\eta^2\right) - \frac{\alpha\hbar\omega_{\mathrm{LO}}}{q} \arcsin q. \tag{4.211}$$

Further, we expand the r.h.s. of Eq. (4.207) to the second order in powers of P (or, what is the same, in powers of q) using the relation

$$\frac{q}{\sqrt{1-q^2}} - \arcsin(q) = \frac{1}{3}q^3 + O\left(q^5\right) \tag{4.212}$$

what results in

$$\frac{\eta}{1-\eta} = \frac{\alpha}{2q^3}\left[\frac{1}{3}q^3 + O\left(q^5\right)\right] = \frac{\alpha}{6} + O\left(q^2\right).$$

Solving this equation for η, we find

$$\eta = \frac{\alpha/6}{1+\alpha/6} + O\left(\frac{P^2}{2m_b\hbar\omega_{\mathrm{LO}}}\right). \tag{4.213}$$

We also apply the expansion in powers of q up to $\sim q^2$ to the energy (4.211):

$$\begin{aligned}
E &= \frac{P^2}{2m_b}\left(1-\eta^2\right) - \frac{\alpha\hbar\omega_{\mathrm{LO}}}{q}\left(q + \frac{1}{6}q^3 + O\left(q^5\right)\right) \\
&= \frac{P^2}{2m_b}\left(1-\eta^2\right) - \alpha\hbar\omega_{\mathrm{LO}} - \frac{1}{6}\alpha\hbar\omega_{\mathrm{LO}}q^2 + \hbar\omega_{\mathrm{LO}}O\left(q^4\right) \\
&= -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{2m_b}\left(1-\eta^2\right) - \frac{\alpha P^2\left(1-\eta\right)^2}{12m_b} + \hbar\omega_{\mathrm{LO}}O\left(q^4\right) \\
&= -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{2m_b}\left(1-\eta^2\right) - \frac{\alpha P^2\left(1-\eta\right)^2}{12m_b} + \hbar\omega_{\mathrm{LO}}O\left(q^4\right) \\
&= -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{12m_b}\left(1-\eta\right)\left(\left(6+\alpha\right)\eta - \alpha + 6\right) + \hbar\omega_{\mathrm{LO}}O\left(q^4\right) \\
&= -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{12m_b}\left(1 - \frac{\alpha/6}{1+\alpha/6}\right)\left(\left(6+\alpha\right)\frac{\alpha/6}{1+\alpha/6} - \alpha + 6\right) + \hbar\omega_{\mathrm{LO}}O\left(q^4\right) \\
&= -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{2m_b\left(1+\alpha/6\right)} + \hbar\omega_{\mathrm{LO}}O\left(q^4\right).
\end{aligned}$$

Finally, we have arrived at the LLP polaron energy

$$E = -\alpha\hbar\omega_{\mathrm{LO}} + \frac{P^2}{2m_b\left(1+\alpha/6\right)} + \hbar\omega_{\mathrm{LO}}O\left(\left(\frac{P^2}{2m_b\hbar\omega_{\mathrm{LO}}}\right)^2\right). \tag{4.214}$$

Appendix 2. Expansion in Stieltjes continuous fractions [52]

In this derivation it is shown that the approximation used in the evaluation of the function, which determines the polaron mass [see Eqs. (40) and (B1) from Ref. [52]]

$$g(k,z)=\int_{-\infty}^{0} d\tau e^{iz\tau}\exp\left[-k^2C(0)\right]\exp\left[k^2C(\tau)\right] \tag{4.215}$$

with

$$C(\tau)=\frac{1}{3}\sum_{k'}\frac{k'^2}{m^2}\left|f_{k'}\right|^2\frac{e^{i\gamma_{k'}\tau}}{\gamma_{k'}^2} \tag{4.216}$$

is equivalent to an expansion in a continued fraction limited to the first step. Moreover, it is proved that the choice of the coefficients of the continued fraction can be justified by a variational principle, at least when z is real and positive.

Expanding the last exponential of Eq. (4.215) in a power series leads to

$$\begin{aligned} g(k,z)=\exp\left[-k^2C(0)\right]&\int_{-\infty}^{0} d\tau e^{iz\tau}\sum_{n=0}^{\infty}\frac{1}{n!}\left(\frac{1}{3m^2}\right)^n \\ &\times\sum_{\vec{k}_1,\ldots,\vec{k}_n}\frac{k_1^2k_2^2\ldots k_n^2\left|f_{k_1}\right|^2\left|f_{k_2}\right|^2\ldots\left|f_{k_n}\right|^2}{\gamma_{k_1}^2\gamma_{k_2}^2\ldots\gamma_{k_n}^2} \\ &\times\exp\left[i(\gamma_{k_1}+\gamma_{k_2}+\ldots\gamma_{k_n}^2)\tau\right] \\ &=-i\exp\left[-k^2C(0)\right]\sum_{n=0}^{\infty}\frac{(3m^2)^{-n}}{n!} \\ &\times\sum_{\vec{k}_1,\ldots,\vec{k}_n}\frac{k_1^2k_2^2\ldots k_n^2\left|f_{k_1}\right|^2\left|f_{k_2}\right|^2\ldots\left|f_{k_n}\right|^2}{\gamma_{k_1}^2\gamma_{k_2}^2\ldots\gamma_{k_n}^2}\frac{1}{\gamma_{k_1}+\gamma_{k_2}+\ldots+\gamma_{k_n}+z}. \end{aligned} \tag{4.217}$$

The multiple sum over the k's is in fact an integral with 3n variables. It is possible to change the variables in that one of the new variables is

$$x_n=\gamma_{k_1}+\gamma_{k_2}+\ldots+\gamma_{k_n}. \tag{4.218}$$

Then the multiple sum which appears in the last term of Eq. (4.217) is of the following type:

$$J(z)=\int_{n\omega}^{\infty}\frac{L(x_n)}{x_n+z}dx_n, \tag{4.219}$$

where

$$L(x_n) = 0$$

is the result of the integration over the $n-1$ other variables. An expansion of integrals of the type (4.219) into Stieltjes continued fractions is known to give good results when z is real and not located on the cut of $J(z)$, i.e., when

$$z > -n\omega. \tag{4.220}$$

The first nontrivial step in the continued fraction expansion is

$$J(z) = \frac{a_0}{a_1 + z} \tag{4.221}$$

with

$$a_0 = \int_{n\omega}^{\infty} L(x_n)dx_n, \tag{4.222}$$

$$a_1 = \frac{\int_{n\omega}^{\infty} x_n L(x_n)dx_n}{\int_{n\omega}^{\infty} L(x_n)dx_n}. \tag{4.223}$$

A *variational principle* can be established, which gives a rather strong argument in favour of the approximation (4.221). Let us introduce a variational parameter $\bar{x}$ writing

$$J(z) = \int_{n\omega}^{\infty} \frac{L(x_n)}{(x_n - \bar{x}) + (z + \bar{x})} dx_n. \tag{4.224}$$

Performing two steps of the division, this relation becomes

$$\begin{aligned} J(z) = {} & \frac{1}{z+\bar{x}} \int_{n\omega}^{\infty} L(x_n)dx_n \\ & - \frac{1}{(z+\bar{x})^2} \int_{n\omega}^{\infty} (x_n - \bar{x})L(x_n)dx_n + K(z,\bar{x}) \end{aligned} \tag{4.225}$$

with

$$K(z,\bar{x}) = \frac{1}{(z+\bar{x})^2}\int_{n\omega}^{\infty}\frac{(x_n-\bar{x})^2 L(x_n)}{x_n+z}dx_n. \tag{4.226}$$

The approximation consists of neglecting the term $K(z,\bar{x})$ in Eq. (4.225). As this term is positive [cf. (4.220)], the best approximation is obtained when it is minimum. Therefore let us use the freedom in the choice of $\bar{x}$ to minimize the expression (4.226),

$$\begin{aligned} \frac{\partial K(z,\bar{x})}{\partial \bar{x}} &= -2\frac{K(z,\bar{x})}{z+\bar{x}} \\ -2\frac{1}{(z+\bar{x})^2}\int_{n\omega}^{\infty}\frac{(x_n-\bar{x})^2 L(x_n)}{x_n+z}dx_n &= 0, \end{aligned} \tag{4.227}$$

which gives

$$-2\frac{1}{(z+\bar{x})^2}\int_{n\omega}^{\infty}\left(\frac{x_n-\bar{x}}{z+\bar{x}}+1\right)\frac{x_n-\bar{x}}{x_n+z}L(x_n)dx_n = 0 \tag{4.228}$$

or

$$-2\frac{1}{(z+\bar{x})^3}\int_{n\omega}^{\infty}(x_n-\bar{x})L(x_n)dx_n = 0. \tag{4.229}$$

This provides us with the best value of the variational parameter

$$\bar{x} = \frac{\int_{n\omega}^{\infty}x_n L(x_n)dx_n}{\int_{n\omega}^{\infty}L(x_n)dx_n}, \tag{4.230}$$

which is a_1 [cf. Eq. (4.223)].

With this value of $\bar{x}$ and neglecting $K(z,\bar{x})$, the expression (4.225) of the calculated quantity $J(z)$ becomes

$$J(z) = \frac{1}{z+\bar{x}}\int_{n\omega}^{\infty}L(x_n)dx_n = J(z) = \frac{a_0}{a_1+z}, \tag{4.231}$$

which is the first step (4.221) of a Stieltjes continued fraction.

To prove that this value of $\bar{x}$ gives a minimum of $K(z,\bar{x})$, let us calculate the second derivative

$$\begin{aligned}\frac{\partial^2 K(z,\bar{x})}{\partial \bar{x}^2} &= \frac{6}{(z+\bar{x})^3}\int\limits_{n\omega}^{\infty}(x_n-\bar{x})L(x_n)dx_n \\ &\quad + \frac{2}{(z+\bar{x})^3}\int\limits_{n\omega}^{\infty}L(x_n)dx_n. \end{aligned} \tag{4.232}$$

Now the parameter $\bar{x}$ is replaced by its expression (4.230). The relation (4.232) becomes

$$\frac{\partial^2 K(z,\bar{x})}{\partial \bar{x}^2} = \frac{2}{(z+a_1)^3}\int\limits_{n\omega}^{\infty}L(x_n)dx_n, \tag{4.233}$$

which is positive of $z \geqslant -n\omega$, since it follows from relation (4.230) that $a_1 > n\omega$.

Our approximation is related to that used by Feynman which is based on the following inequality:

$$\left\langle e^{-sx}\right\rangle \geqslant e^{-s\langle x\rangle}, \tag{4.234}$$

where the brackets denote the expectation value of the random variable x. For instance,

$$\left\langle e^{-sx}\right\rangle = \frac{\displaystyle\int_a^{\infty} L(x)e^{-sx}dx}{\displaystyle\int_a^{\infty} L(x)ds}, \tag{4.235}$$

where L(x) is the non-normalized probability density of x. The Laplace transform of Eq. (4.234) gives

$$\int_0^{\infty} e^{-sz}\left\langle e^{-sx}\right\rangle ds \geqslant \int_0^{\infty} e^{-sz}e^{-s\langle x\rangle}ds, \tag{4.236}$$

which after integration becomes

$$\int_a^{\infty}\frac{L(x)}{x+z}dx \geqslant \frac{\displaystyle\int_a^{\infty} L(x)ds}{\langle x\rangle + z} = \frac{a_0}{a_1+z}. \tag{4.237}$$

The last inequality shows the relation with our procedure.

Part II

Many polarons

V. OPTICAL CONDUCTIVITY OF AN INTERACTING MANY-POLARON GAS

A. Kubo formula for the optical conductivity of the many-polaron gas

The derivations in the present section are based on Ref. [51]. The Hamiltonian of a system of N interacting continuum polarons is given by:

$$
\begin{aligned}
H_0 = & \sum_{j=1}^{N} \frac{p_j^2}{2m_b} + \sum_{\mathbf{q}} \hbar\omega_{\mathrm{LO}} b_{\mathbf{q}}^{+} b_{\mathbf{q}} \\
& + \sum_{\mathbf{q}} \sum_{j=1}^{N} \left(e^{i\mathbf{q}\cdot\mathbf{r}_j} b_{\mathbf{q}} V_{\mathbf{q}} + e^{-i\mathbf{q}\cdot\mathbf{r}_j} b_{\mathbf{q}}^{+} V_{\mathbf{q}}^{*} \right) + \frac{e^2}{2\varepsilon_\infty} \sum_{j=1}^{N} \sum_{\ell(\neq j)=1}^{N} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|},
\end{aligned} \tag{5.1}
$$

where $\mathbf{r}_j, \mathbf{p}_j$ represent the position and momentum of the N constituent electrons (or holes) with band mass m_b; $b_{\mathbf{q}}^{+}, b_{\mathbf{q}}$ denote the creation and annihilation operators for longitudinal optical (LO) phonons with wave vector $\mathbf{q}$ and frequency ω_{LO}; $V_{\mathbf{q}}$ describes the amplitude of the interaction between the electrons and the phonons; and e is the elementary electron charge. This Hamiltonian can be subdivided into the following parts:

$$
H = H_e + H_{e-e} + H_{ph} + H_{e-ph} \tag{5.2}
$$

where

$$
H_e = \sum_{j=1}^{N} \frac{p_j^2}{2m_b} \tag{5.3}
$$

is the kinetic energy of electrons,

$$
H_{e-e} = \frac{e^2}{2\varepsilon_\infty} \sum_{j=1}^{N} \sum_{\ell(\neq j)=1}^{N} \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} \tag{5.4}
$$

is the potential energy of the Coulomb electron-electron interaction,

$$
H_{ph} = \sum_{\mathbf{q}} \hbar\omega_{\mathrm{LO}} b_{\mathbf{q}}^{+} b_{\mathbf{q}} \tag{5.5}
$$

is the Hamlitonian of phonons, and

$$H_{e-ph} = \sum_{\mathbf{q}} \sum_{j=1}^{N} \left(e^{i\mathbf{q}\cdot\mathbf{r}_j} b_{\mathbf{q}} V_{\mathbf{q}} + e^{-i\mathbf{q}\cdot\mathbf{r}_j} b_{\mathbf{q}}^{+} V_{\mathbf{q}}^{*} \right) \tag{5.6}$$

is the Hamiltonian of the electron-phonon interaction. Further on, we use the second quantization formalism for electrons, in which the terms H_e, H_{e-e} and H_{e-ph} are

$$H_e = \sum_{\mathbf{k},\sigma} \frac{\hbar^2 k^2}{2m_b} a_{\mathbf{k},\sigma}^{+} a_{\mathbf{k},\sigma}, \tag{5.7}$$

$$H_{e-e} = \frac{1}{2} \sum_{\mathbf{q}\neq 0} v_{\mathbf{q}} \sum_{\substack{\mathbf{k},\sigma \\ \mathbf{k}',\sigma'}} a_{\mathbf{k}+\mathbf{q},\sigma}^{+} a_{\mathbf{k}'-\mathbf{q},\sigma'}^{+} a_{\mathbf{k}',\sigma'} a_{\mathbf{k},\sigma} = \frac{1}{2} \sum_{\mathbf{q}\neq 0} v_{\mathbf{q}} : \rho_{\mathbf{q}} \rho_{-\mathbf{q}} :, \tag{5.8}$$

$$H_{e-ph} = \sum_{\mathbf{q}} \left(V_{\mathbf{q}} b_{\mathbf{q}} \rho_{\mathbf{q}} + V_{\mathbf{q}}^{*} b_{\mathbf{q}}^{+} \rho_{-\mathbf{q}} \right), \tag{5.9}$$

where : ... : is the symbol of the normal product of operators,

$$v_{\mathbf{q}} = \frac{4\pi e^2}{\varepsilon_{\infty} q^2 V} \tag{5.10}$$

is the Fourier component of the Coulomb potential, and

$$\rho_{\mathbf{q}} = \sum_{j=1}^{N} e^{i\mathbf{q}\cdot\mathbf{r}_j} = \sum_{\mathbf{k},\sigma} a_{\mathbf{k}+\mathbf{q},\sigma}^{+} a_{\mathbf{k},\sigma} \tag{5.11}$$

is the Fourier component of the electron density.

The ground state energy of the many-polaron Hamiltonian (5.1) has been studied by L. Lemmens, J. T. Devreese and F. Brosens (LDB) [89], for weak and intermediate strength of the electron-phonon coupling. They introduce a variational wave function:

$$|\psi_{\mathrm{LDB}}\rangle = U\,|\phi\rangle \left|\psi_{el}^{(0)}\right\rangle, \tag{5.12}$$

where $\left|\psi_{el}^{(0)}\right\rangle$ represents the ground-state many-body wave function for the electron (or hole) system and $|\phi\rangle$ is the phonon vacuum, U is a many-body unitary operator which determines a canonical transformation for a fermion gas interacting with a boson field:

$$U = \exp\left\{ \sum_{j=1}^{N} \sum_{\mathbf{q}} \left(f_{\mathbf{q}} a_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{r}_j} - f_{\mathbf{q}}^{*} a_{\mathbf{q}}^{+} e^{-i\mathbf{q}\cdot\mathbf{r}_j} \right) \right\}. \tag{5.13}$$

In Ref. [89], this canonical transformation was used to establish a many-fermion theory. The $f_{\mathbf{q}}$ were determined variationally [89] resulting in

$$f_{\mathbf{q}} = \frac{V_{\mathbf{q}}}{\hbar\omega_{\mathrm{LO}} + \dfrac{\hbar^2 q^2}{2m_b S(\mathbf{q})}}, \tag{5.14}$$

for a system with total momentum $\mathbf{P} = \sum_j \mathbf{p}_j = 0$. In this expression, $S(\mathbf{q})$ represents the static structure factor of the constituent interacting many electron or hole system :

$$NS(\mathbf{q}) = \left\langle \sum_{j=1}^{N} \sum_{j'=1}^{N} e^{i\mathbf{q}\cdot(\mathbf{r}_j - \mathbf{r}_{j'})} \right\rangle . \tag{5.15}$$

The angular brackets $\langle ... \rangle$ represent the expectation value with respect to the ground state.

The many-polaron optical conductivity is the response of the current-density, in the system described by the Hamiltonian (5.1), to an applied electric field (along the x-axis) with frequency ω. This applied electric field introduces a perturbation term in the Hamiltonian (5.1), which couples the vector potential of the incident electromagnetic field to the current-density. Within linear response theory, the optical conductivity can be expressed through the Kubo formula as a current-current correlation function:

$$\sigma(\omega) = i\frac{Ne^2}{Vm_b\omega} + \frac{1}{V\hbar\omega} \int_0^\infty e^{i\omega t} \left\langle [J_x(t), J_x(0)] \right\rangle dt. \tag{5.16}$$

In this expression, V is the volume of the system, and J_x is the x-component of the current operator $\mathbf{J}$, which is related to the momentum operators of the charge carriers:

$$\mathbf{J} = \frac{q}{m_b} \sum_{j=1}^{N} \mathbf{p}_j = \frac{q}{m_b}\mathbf{P}, \tag{5.17}$$

with q the charge of the charge carriers ($+e$ for holes, $-e$ for electrons) and $\mathbf{P}$ the total momentum operator of the charge carriers. The real part of the optical conductivity at temperature zero, which is proportional to the optical absorption coefficient, can be written as a function of the total momentum operator of the charge carriers as follows :

$$\operatorname{Re}\sigma(\omega) = \frac{1}{V\hbar\omega} \frac{e^2}{m_b^2} \operatorname{Re}\left\{ \int_0^\infty e^{i\omega t} \left\langle [P_x(t), P_x(0)] \right\rangle dt \right\} . \tag{5.18}$$

B. Force-force correlation function

Let us integrate over time in (5.18) twice by parts as follows:

$$\begin{aligned}
&\int_0^\infty dt \left\langle \left[P_x(t), P_x\right]\right\rangle e^{i\omega t-\delta t} \\
&= \frac{1}{i\omega-\delta}\left\{ \left\langle \left[P_x(t), P_x\right]\right\rangle e^{i\omega t-\delta t}\Big|_{t=0}^{\infty} - \int_0^\infty dt \left\langle \left[\frac{d}{dt}P_x(t), P_x\right]\right\rangle e^{i\omega t-\delta t}\right\} \\
&= -\frac{1}{i\omega-\delta}\int_0^\infty dt \left\langle \left[\frac{d}{dt}\left(e^{\frac{it}{\hbar}H}P_x e^{-\frac{it}{\hbar}H}\right), P_x\right]\right\rangle e^{i\omega t-\delta t} \\
&= -\frac{1}{i\omega-\delta}\int_0^\infty dt \left\langle \left[\left(e^{\frac{it}{\hbar}H}\frac{i}{\hbar}\left[H,P_x\right]e^{-\frac{it}{\hbar}H}\right), P_x\right]\right\rangle e^{i\omega t-\delta t} \\
&= -\frac{1}{i\omega-\delta}\int_0^\infty dt \left\langle \left[\frac{i}{\hbar}\left[H,P_x\right], e^{-\frac{it}{\hbar}H}P_x e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t} \\
&= -\frac{1}{i\omega-\delta}\int_0^\infty dt \left\langle \left[F_x(0), e^{-\frac{it}{\hbar}H}P_x e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t} \\
&= -\left(\frac{1}{i\omega-\delta}\right)^2 \left\{ \left\langle \left[F_x(0), e^{-\frac{it}{\hbar}H}P_x e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t}\Big|_{t=0}^{\infty} \right. \\
&\left. - \int_0^\infty dt \left\langle \left[F_x(0), \frac{d}{dt}e^{-\frac{it}{\hbar}H}P_x e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t}\right\}
\end{aligned}$$

$$\begin{aligned}
&= -\left(\frac{1}{i\omega-\delta}\right)^2 \left\{ -\left\langle \left[F_x, P_x\right]\right\rangle + \int_0^\infty dt \left\langle \left[F_x(0), e^{-\frac{it}{\hbar}H}\left(\frac{i}{\hbar}\left[H,P_x\right]\right)e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t}\right\} \\
&= -\left(\frac{1}{i\omega-\delta}\right)^2 \left\{ -\left\langle \left[F_x, P_x\right]\right\rangle + \int_0^\infty dt \left\langle \left[F_x(0), e^{-\frac{it}{\hbar}H}F_x(0)\, e^{\frac{it}{\hbar}H}\right]\right\rangle e^{i\omega t-\delta t}\right\} \\
&= -\left(\frac{1}{i\omega-\delta}\right)^2 \left\{ -\left\langle \left[F_x, P_x\right]\right\rangle + \int_0^\infty dt \left\langle \left[e^{\frac{it}{\hbar}H}F_x(0)\, e^{-\frac{it}{\hbar}H}, F_x(0)\right]\right\rangle e^{i\omega t-\delta t}\right\} \\
&= \frac{1}{(\omega+i\delta)^2}\left\{ -\left\langle \left[F_x, P_x\right]\right\rangle + \int_0^\infty dt \left\langle \left[F_x(t), F_x(0)\right]\right\rangle e^{i\omega t-\delta t}\right\},
\end{aligned}$$

where $\mathbf{F} \equiv \frac{i}{\hbar}\left[H, \mathbf{P}\right]$ is the operator of the force applied to the center of mass of the electrons.

Since both F_x and P_x are hermitian operators, the average $\left\langle \left[F_x, P_x\right]\right\rangle$ is imaginary. Hence, for $\omega \neq 0$, this term does not give a contribution into $\operatorname{Re}\sigma(\omega)$. As a result, integrating by parts twice, the real part of the optical conductivity of the many-polaron system is written with a force-force correlation function:

$$\operatorname{Re}\sigma(\omega) = \frac{1}{V\hbar\omega^3}\frac{e^2}{m_b^2}\operatorname{Re}\left\{\int_0^\infty e^{i\omega t}\left\langle \left[F_x(t), F_x(0)\right]\right\rangle dt\right\}. \tag{5.19}$$

The force operator is determined as

$$F_x = \frac{i}{\hbar}\left[H, P_x\right] = \frac{i}{\hbar}\left[H_e + H_{e-e} + H_{ph} + H_{e-ph}, P_x\right].$$

Further, we use the commutators:

$$\left[a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma}, P_x\right] = \sum_{\mathbf{k}'} \hbar k'_x \left[a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma}, a^+_{\mathbf{k}',\sigma}a_{\mathbf{k}',\sigma}\right]$$

$$= \sum_{\mathbf{k}'} \hbar k'_x \begin{pmatrix} a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma}a^+_{\mathbf{k}',\sigma}a_{\mathbf{k}',\sigma} + a^+_{\mathbf{k}+\mathbf{q},\sigma}a^+_{\mathbf{k}',\sigma}a_{\mathbf{k},\sigma}a_{\mathbf{k}',\sigma} \\ -a^+_{\mathbf{k}',\sigma}a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k}',\sigma}a_{\mathbf{k},\sigma} - a^+_{\mathbf{k}',\sigma}a_{\mathbf{k}',\sigma}a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma} \end{pmatrix}$$

$$= \sum_{\mathbf{k}'} \hbar k'_x \left(\delta_{\mathbf{k}\mathbf{k}'}a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k}',\sigma} - \delta_{\mathbf{k}',\mathbf{k}+\mathbf{q}}a^+_{\mathbf{k}',\sigma}a_{\mathbf{k},\sigma}\right)$$

$$= \sum_{\mathbf{k}'} \hbar k'_x \left(\delta_{\mathbf{k}\mathbf{k}'}a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma} - \delta_{\mathbf{k}',\mathbf{k}+\mathbf{q}}a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma}\right)$$

$$= a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma} \sum_{\mathbf{k}'} \hbar k'_x \left(\delta_{\mathbf{k}\mathbf{k}'} - \delta_{\mathbf{k}',\mathbf{k}+\mathbf{q}}\right) = -\hbar q_x a^+_{\mathbf{k}+\mathbf{q},\sigma}a_{\mathbf{k},\sigma},$$

$$[\rho_{\mathbf{q}}, P_x] = -\hbar q_x \rho_{\mathbf{q}}.$$

Hence, $[H_e, P_x] = 0$, $[H_{e-e}, P_x] = 0$,

$$[H_{e-ph}, P_x] = \sum_{\mathbf{q}} \left(V_{\mathbf{q}} b_{\mathbf{q}} [\rho_{\mathbf{q}}, P_x] + V^*_{\mathbf{q}} b^+_{\mathbf{q}} [\rho_{-\mathbf{q}}, P_x]\right)$$

$$= -\hbar \sum_{\mathbf{q}} q_x \left(V_{\mathbf{q}} b_{\mathbf{q}} \rho_{\mathbf{q}} - V^*_{\mathbf{q}} b^+_{\mathbf{q}} \rho_{-\mathbf{q}}\right),$$

So, the commutator of the Hamiltonian (5.1) with the total momentum operator of the charge carriers leads to the expression for the force

$$\mathbf{F} = -i \sum_{\mathbf{q}} \mathbf{q} \left(V_{\mathbf{q}} b_{\mathbf{q}} \rho_{\mathbf{q}} - V^*_{\mathbf{q}} b^+_{\mathbf{q}} \rho_{-\mathbf{q}}\right). \tag{5.20}$$

This result for the force operator clarifies the significance of using the force-force correlation function rather than the momentum-momentum correlation function. The operator product $F_x(t)F_x(0)$ is proportional to $|V_{\mathbf{k}}|^2$, the charge carrier - phonon interaction strength. This will be a distinct advantage for any expansion of the final result in the charge carrier - phonon interaction strength, since one power of $|V_{\mathbf{k}}|^2$ is factored out beforehand. Substituting (5.20) into (5.19), the real part of the optical conductivity then takes the form:

$$\operatorname{Re}\sigma(\omega) = \frac{1}{V\hbar\omega^3}\frac{e^2}{m_b^2} \operatorname{Re} \int_0^\infty dt e^{i\omega t - \delta t} \sum_{\mathbf{q},\mathbf{q}'} q_x q'_x$$

$$\times \left\langle \left[\left[V_{\mathbf{q}} b_{\mathbf{q}}(t) + V^*_{-\mathbf{q}} b^+_{-\mathbf{q}}(t)\right] \rho_{\mathbf{q}}(t), \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V^*_{\mathbf{q}'} b^+_{\mathbf{q}'}\right) \rho_{-\mathbf{q}'}\right]\right\rangle. \tag{5.21}$$

Up to this point, no approximations other than *linear response theory* have been made.

C. Canonical transformation

The expectation value appearing in the right hand side of expression (5.21) for the real part of the optical conductivity is calculated now with respect to the LDB many-polaron wave function (5.12). The unitary operator (5.13) can be written as

$$U = \exp \sum_{\mathbf{q}} A_{\mathbf{q}} \rho_{\mathbf{q}}, \quad A_{\mathbf{q}} = f_{\mathbf{q}} b_{\mathbf{q}} - f_{-\mathbf{q}}^{*} b_{-\mathbf{q}}^{+}, \tag{5.22}$$

The transformed Hamiltonian (5.2) is denoted as

$$\tilde{H} = U^{-1} H U. \tag{5.23}$$

The momentum operator of an electron $\mathbf{p}_j$, the operator of the total momentum of electrons $\mathbf{P}$ and the phonon creation and annihilation operators are transformed by the unitary transformation (5.22) as follows:

$$U^{-1} \mathbf{p}_j U = \mathbf{p}_j + \sum_{\mathbf{q}} \hbar \mathbf{q} A_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{r}_j}, \tag{5.24}$$

$$U^{-1} \mathbf{P} U = \mathbf{P} + \sum_{\mathbf{q}} \hbar \mathbf{q} A_{\mathbf{q}} \rho_{\mathbf{q}}, \tag{5.25}$$

$$U^{-1} b_{\mathbf{q}} U = b_{\mathbf{q}} - f_{\mathbf{q}}^{*} \rho_{-\mathbf{q}}, \quad U^{-1} b_{\mathbf{q}}^{+} U = b_{\mathbf{q}}^{+} - f_{\mathbf{q}} \rho_{\mathbf{q}}. \tag{5.26}$$

As a result, after the transformation (5.22), the Hamiltonian takes the form (see Ref. [89]):

$$\tilde{H} = H_e + \tilde{H}_{e-e} + H_{ph} + \tilde{H}_{e-ph} + H_N + H_{ppe}, \tag{5.27}$$

where the terms are

$$\tilde{H}_{e-e} = \frac{1}{2} \sum_{\mathbf{q}\neq 0} \tilde{v}_{\mathbf{q}} : \rho_{\mathbf{q}} \rho_{-\mathbf{q}} :, \quad \tilde{v}_{\mathbf{q}} = v_{\mathbf{q}} + 2\left(\hbar\omega_{\mathrm{LO}} \left|f_{\mathbf{q}}\right|^2 - V_{\mathbf{q}} f_{\mathbf{q}}^{*} - V_{\mathbf{q}}^{*} f_{\mathbf{q}}\right), \tag{5.28}$$

$$\begin{aligned} \tilde{H}_{e-ph} = & \sum_{\mathbf{q}} \left[\left(V_{\mathbf{q}} - \hbar\omega_{\mathrm{LO}} f_{\mathbf{q}}\right) b_{\mathbf{q}} \rho_{\mathbf{q}} + \left(V_{\mathbf{q}}^{*} - \hbar\omega_{\mathrm{LO}} f_{\mathbf{q}}^{*}\right) b_{\mathbf{q}}^{+} \rho_{-\mathbf{q}} \right] \\ & + \frac{\hbar^2}{2m_b} \sum_{\mathbf{q}} A_{\mathbf{q}} \sum_{\mathbf{k},\sigma} \left(\mathbf{q}^2 + 2\mathbf{k}\cdot\mathbf{q}\right) a_{\mathbf{k}+\mathbf{q},\sigma}^{+} a_{\mathbf{k},\sigma}, \end{aligned} \tag{5.29}$$

$$H_N = \hat{N} \sum_{\mathbf{q}} \left(\hbar\omega_{\mathrm{LO}} \left|f_{\mathbf{q}}\right|^2 - V_{\mathbf{q}} f_{\mathbf{q}}^{*} - V_{\mathbf{q}}^{*} f_{\mathbf{q}}\right), \quad \left(\hat{N} \equiv \sum_{\mathbf{k},\sigma} a_{\mathbf{k},\sigma}^{+} a_{\mathbf{k},\sigma}\right), \tag{5.30}$$

$$H_{ppe} = \frac{\hbar^2}{2m_b} \sum_{\mathbf{q}\mathbf{q}'} \mathbf{q} \cdot \mathbf{q}' A_{\mathbf{q}} A_{\mathbf{q}'} \rho_{\mathbf{q}+\mathbf{q}'}. \tag{5.31}$$

The exact expression for the real part of the conductivity (5.21) after the replacement of $|\Psi_0\rangle$ by $|\Psi_{LDB}\rangle = U|\phi\rangle \left|\psi_{el}^{(0)}\right\rangle$ is transformed to the approximate one

$$\begin{aligned}
&\operatorname{Re}\sigma(\omega) \\
&= \frac{1}{V\hbar\omega^3} \frac{e^2}{m_b^2} \operatorname{Re} \int_0^\infty dt e^{i\omega t - \delta t} \sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \\
&\times \left\langle \psi_{el}^{(0)} \middle| \left\langle \phi \middle| U^{-1} \left[\begin{array}{c} e^{\frac{it}{\hbar}H} \left[V_{\mathbf{q}} b_{\mathbf{q}} + V_{-\mathbf{q}}^* b_{-\mathbf{q}}^+ \right] \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}H}, \\ \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V_{\mathbf{q}'}^* b_{\mathbf{q}'}^+ \right) \rho_{-\mathbf{q}'} \end{array} \right] U \middle| \phi \right\rangle \middle| \psi_{el}^{(0)} \right\rangle
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{V\hbar\omega^3} \frac{e^2}{m_b^2} \operatorname{Re} \int_0^\infty dt e^{i\omega t - \delta t} \sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \\
&\times \left\langle \psi_{el}^{(0)} \middle| \left\langle \phi \middle| \left[\begin{array}{c} e^{\frac{it}{\hbar}\tilde{H}} U^{-1} \left[V_{\mathbf{q}} b_{\mathbf{q}} + V_{-\mathbf{q}}^* b_{-\mathbf{q}}^+ \right] U \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}}, \\ U^{-1} \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V_{\mathbf{q}'}^* b_{\mathbf{q}'}^+ \right) U \rho_{-\mathbf{q}'} \end{array} \right] \middle| \phi \right\rangle \middle| \psi_{el}^{(0)} \right\rangle
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{V\hbar\omega^3} \frac{e^2}{m_b^2} \operatorname{Re} \int_0^\infty dt e^{i\omega t - \delta t} \sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \\
&\times \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| \left[e^{\frac{it}{\hbar}\tilde{H}} \left(V_{\mathbf{q}} \left(b_{\mathbf{q}} - f_{\mathbf{q}}^* \rho_{-\mathbf{q}} \right) + V_{-\mathbf{q}}^* \left(b_{-\mathbf{q}}^+ - f_{-\mathbf{q}} \rho_{-\mathbf{q}} \right) \right) \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}}, \right. \\
&\left. \left(V_{-\mathbf{q}'} \left(b_{-\mathbf{q}'} - f_{-\mathbf{q}'}^* \rho_{\mathbf{q}'} \right) + V_{\mathbf{q}'}^* \left(b_{\mathbf{q}'}^+ - f_{\mathbf{q}'} \rho_{\mathbf{q}'} \right) \right) \rho_{-\mathbf{q}'} \right] \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle.
\end{aligned}$$

So, we have arrived at the expression

$$\begin{aligned}
\operatorname{Re}\sigma(\omega) = \frac{1}{V\hbar\omega^3} \frac{e^2}{m_b^2} \operatorname{Re} \int_0^\infty dt e^{i\omega t - \delta t} \sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \\
\times \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| \left[e^{\frac{it}{\hbar}\tilde{H}} \left(V_{\mathbf{q}} \left(b_{\mathbf{q}} - f_{\mathbf{q}}^* \rho_{-\mathbf{q}} \right) + V_{-\mathbf{q}}^* \left(b_{-\mathbf{q}}^+ - f_{-\mathbf{q}} \rho_{-\mathbf{q}} \right) \right) \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}}, \right. \\
\left. \left(V_{-\mathbf{q}'} \left(b_{-\mathbf{q}'} - f_{-\mathbf{q}'}^* \rho_{\mathbf{q}'} \right) + V_{\mathbf{q}'}^* \left(b_{\mathbf{q}'}^+ - f_{\mathbf{q}'} \rho_{\mathbf{q}'} \right) \right) \rho_{-\mathbf{q}'} \right] \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle.
\end{aligned}$$

Since $\rho_{\mathbf{q}}\rho_{-\mathbf{q}} = \rho_{-\mathbf{q}}\rho_{\mathbf{q}}$, and $V_{\mathbf{q}} f_{\mathbf{q}}^* = V_{-\mathbf{q}} f_{-\mathbf{q}}^*$, the terms proportional to $\rho_{-\mathbf{q}}\rho_{\mathbf{q}}$ vanish after the summation over $\mathbf{q}$:

$$\sum_{\mathbf{q}} q_x V_{\mathbf{q}} f_{\mathbf{q}}^* \rho_{-\mathbf{q}} \rho_{\mathbf{q}} \overset{\mathbf{q} \to -\mathbf{q}}{=} -\sum_{\mathbf{q}} q_x V_{\mathbf{q}} f_{\mathbf{q}}^* \rho_{-\mathbf{q}} \rho_{\mathbf{q}} = 0. \tag{5.32}$$

Hence we obtain the real part of the optical conductivity in the form

$$\begin{aligned}\operatorname{Re}\sigma(\omega) = {} & \frac{1}{V\hbar\omega^3}\frac{e^2}{m_b^2}\operatorname{Re}\int_0^\infty dte^{i\omega t-\delta t}\sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \\ & \times \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| \left[e^{\frac{it}{\hbar}\tilde{H}} \left(V_{\mathbf{q}} b_{\mathbf{q}} + V_{-\mathbf{q}}^* b_{-\mathbf{q}}^+ \right) \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}}, \right. \\ & \left. \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V_{\mathbf{q}'}^* b_{\mathbf{q}'}^+ \right) \rho_{-\mathbf{q}'} \right] \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle. \end{aligned} \tag{5.33}$$

Introducing the factor

$$\begin{aligned}\mathcal{J}(\mathbf{q},\mathbf{q}') = {} & \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| \left[e^{\frac{it}{\hbar}\tilde{H}} \left(V_{\mathbf{q}} b_{\mathbf{q}} + V_{-\mathbf{q}}^* b_{-\mathbf{q}}^+ \right) \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}}, \right. \\ & \left. \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V_{\mathbf{q}'}^* b_{\mathbf{q}'}^+ \right) \rho_{-\mathbf{q}'} \right] \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle, \end{aligned} \tag{5.34}$$

the optical conductivity can be written as

$$\operatorname{Re}\sigma(\omega) = \frac{1}{V\hbar\omega^3}\frac{e^2}{m_b^2}\operatorname{Re}\int_0^\infty dte^{i\omega t-\delta t}\sum_{\mathbf{q},\mathbf{q}'} q_x q_x' \mathcal{J}(\mathbf{q},\mathbf{q}'). \tag{5.35}$$

In the case of a weak electron-phonon coupling, we can neglect in the exponent $e^{-\frac{it}{\hbar}\tilde{H}}$ of (5.33) the terms $\tilde{H}_{e-ph}$ and H_{ppe} [i. e., the renormalized Hamiltonian of the electron-phonon interaction (5.28) and (5.31)]. Namely, we replace $\tilde{H}$ in Eq. (5.33) by the Hamiltonian

$$\tilde{H}_0 = H_e + \tilde{H}_{e-e} + H_{ph} + H_N. \tag{5.36}$$

In this case, we find

$$\begin{aligned}\mathcal{J}(\mathbf{q},\mathbf{q}') = {} & \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| \left[e^{\frac{it}{\hbar}\tilde{H}_0} \left(V_{\mathbf{q}} b_{\mathbf{q}} + V_{-\mathbf{q}}^* b_{-\mathbf{q}}^+ \right) \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0}, \right. \\ & \left. \left(V_{-\mathbf{q}'} b_{-\mathbf{q}'} + V_{\mathbf{q}'}^* b_{\mathbf{q}'}^+ \right) \rho_{-\mathbf{q}'} \right] \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle \end{aligned}$$

$$\begin{aligned} & = |V_{\mathbf{q}}|^2 \delta_{\mathbf{q}\mathbf{q}'} \left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| e^{i\tilde{H}_0 t/\hbar} \rho_{\mathbf{q}} b_{\mathbf{q}} e^{-i\tilde{H}_0 t/\hbar} \rho_{-\mathbf{q}} b_{\mathbf{q}}^+ \\ & -\rho_{\mathbf{q}} b_{\mathbf{q}} e^{i\tilde{H}_0 t/\hbar} \rho_{-\mathbf{q}} b_{\mathbf{q}}^+ e^{-i\tilde{H}_0 t/\hbar} \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle \\ & = 2i |V_{\mathbf{q}}|^2 \delta_{\mathbf{q}\mathbf{q}'} \operatorname{Im} \left[\left\langle \psi_{el}^{(0)} \right| \left\langle \phi \right| e^{i\tilde{H}_0 t/\hbar} \rho_{\mathbf{q}} b_{\mathbf{q}} e^{-i\tilde{H}_0 t/\hbar} \rho_{-\mathbf{q}} b_{\mathbf{q}}^+ \left| \phi \right\rangle \left| \psi_{el}^{(0)} \right\rangle \right]. \end{aligned}$$

The time-dependent phonon operators are

$$e^{i\tilde{H}_0 t/\hbar} b_{\mathbf{q}} e^{-i\tilde{H}_0 t/\hbar} = b_{\mathbf{q}} e^{-i\omega_{\mathrm{LO}} t},$$

so that we have

$$\begin{aligned}\mathcal{J}(\mathbf{q},\mathbf{q}') &= 2i|V_{\mathbf{q}}|^2\delta_{\mathbf{q}\mathbf{q}'}\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}\left\langle\psi_{el}^{(0)}\left|\left\langle\phi\left|e^{i\tilde{H}_0t/\hbar}\rho_{\mathbf{q}}e^{-i\tilde{H}_0t/\hbar}\rho_{-\mathbf{q}}b_{\mathbf{q}}b_{\mathbf{q}}^{+}\right|\phi\right\rangle\right|\psi_{el}^{(0)}\right\rangle\right]\\ &= 2i|V_{\mathbf{q}}|^2\delta_{\mathbf{q}\mathbf{q}'}\operatorname{Im}\left[\left\langle\psi_{el}^{(0)}\left|e^{i\tilde{H}_et/\hbar}\rho_{\mathbf{q}}e^{-i\tilde{H}_et/\hbar}\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\left\langle\phi\left|b_{\mathbf{q}}b_{\mathbf{q}}^{+}\right|\phi\right\rangle\right],\end{aligned}$$

where $\tilde{H}_e = H_e + \tilde{H}_{e-e} + H_N$.

Taking the expectation value with respect to the phonon vacuum, we find

$$\mathcal{J}(\mathbf{q},\mathbf{q}') = 2i|V_{\mathbf{q}}|^2\delta_{\mathbf{q}\mathbf{q}'}\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}\left\langle\psi_{el}^{(0)}\left|e^{i\tilde{H}_et/\hbar}\rho_{\mathbf{q}}e^{-i\tilde{H}_et/\hbar}\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\right]. \tag{5.37}$$

The optical conductivity (5.33) then takes the form:

$$\begin{aligned}\operatorname{Re}\sigma(\omega) &= -\frac{2e^2}{V\hbar m_b^2\omega^3}\operatorname{Im}\int_0^\infty dte^{i\omega t-\delta t}\sum_{\mathbf{q}}q_x^2|V_{\mathbf{q}}|^2\\ &\times\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}\left\langle\psi_{el}^{(0)}\left|e^{i\tilde{H}_et/\hbar}\rho_{\mathbf{q}}e^{-i\tilde{H}_et/\hbar}\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\right]\end{aligned} \tag{5.38}$$

For an isotropic electron-phonon system, q_x^2 in 3D can be replaced by $\frac{1}{3}\left(q_x^2+q_y^2+q_z^2\right)=\frac{1}{3}q^2$, what gives us the result

$$\operatorname{Re}\sigma_{3\mathrm{D}}(\omega) = -\frac{2}{3V\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}}q^2|V_{\mathbf{q}}|^2\operatorname{Im}\int_0^\infty dte^{i\omega t-\delta t}\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}F(\mathbf{q},t)\right], \tag{5.39}$$

where the two-point correlation function is

$$F(\mathbf{q},t) = \left\langle\psi_{el}^{(0)}\left|e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle. \tag{5.40}$$

The same derivation for the 2D case, provides the expression

$$\operatorname{Re}\sigma_{2\mathrm{D}}(\omega) = -\frac{1}{A\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}}q^2|V_{\mathbf{q}}|^2\operatorname{Im}\int_0^\infty dte^{i\omega t-\delta t}\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}F(\mathbf{q},t)\right], \tag{5.41}$$

where A is the surface of the 2D system.

D. Dynamic structure factor

To find the formula for the real part of the optical conductivity in its final form, we introduce the standard expression for the dynamic structure factor of the system of charge carriers interacting through a Coulomb potential,

$$S(\mathbf{q},\omega) = \frac{1}{2N}\int_{-\infty}^{\infty}\left\langle\psi_{el}^{(0)}\left|\sum_{j,\ell}e^{i\mathbf{q}\cdot(\mathbf{r}_j(t)-\mathbf{r}_\ell(0))}\right|\psi_{el}^{(0)}\right\rangle e^{i\omega t}dt. \tag{5.42}$$

The dynamic structure factor is expressed in terms of the two-point correlation function as follows:

$$S(\mathbf{q},\omega) = \frac{1}{2N}\int\limits_{-\infty}^{\infty} \left\langle \psi_{el}^{(0)} \left| e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}} \right| \psi_{el}^{(0)} \right\rangle e^{i\omega t} dt$$
$$= \frac{1}{2N}\int\limits_{-\infty}^{\infty} F\left(\mathbf{q},t\right) e^{i\omega t} dt = \frac{1}{2N} F\left(\mathbf{q},\omega\right)$$
$$\Downarrow$$
$$S(\mathbf{q},\omega) = \frac{1}{2N} F\left(\mathbf{q},\omega\right), \tag{5.43}$$

where $F\left(\mathbf{q},\omega\right)$ is the Fourier image of $F\left(\mathbf{q},t\right)$:

$$F\left(\mathbf{q},\omega\right) = \int\limits_{-\infty}^{\infty} F\left(\mathbf{q},t\right) e^{i\omega t} dt. \tag{5.44}$$

The function $F\left(\mathbf{q},t\right)$ obeys the following property:

$$\begin{aligned} F^{*}\left(\mathbf{q},t\right) &= \left\langle \psi_{el}^{(0)} \left| \rho_{-\mathbf{q}}^{+} e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{\mathbf{q}}^{+} e^{-\frac{it}{\hbar}\tilde{H}_e} \right| \psi_{el}^{(0)} \right\rangle \\ &= \left\langle \psi_{el}^{(0)} \left| \rho_{\mathbf{q}} e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_e} \right| \psi_{el}^{(0)} \right\rangle \\ &= \left\langle \psi_{el}^{(0)} \left| \rho_{\mathbf{q}} e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}} \right| \psi_{el}^{(0)} \right\rangle e^{-\frac{it}{\hbar}\tilde{E}_0}, \end{aligned}$$

where $\tilde{E}_0$ is the eigenvalue of the Hamiltonian $\tilde{H}_e$:

$$\tilde{H}_e \left| \psi_{el}^{(0)} \right\rangle = \tilde{E}_0 \left| \psi_{el}^{(0)} \right\rangle.$$

Herefrom, we find that

$$\begin{aligned} F^{*}\left(\mathbf{q},t\right) &= e^{-\frac{it}{\hbar}\tilde{E}_0} \left\langle \psi_{el}^{(0)} \left| \rho_{\mathbf{q}} e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}} \right| \psi_{el}^{(0)} \right\rangle \\ &= \left\langle \psi_{el}^{(0)} \left| e^{-\frac{it}{\hbar}\tilde{H}_e}\rho_{\mathbf{q}} e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{-\mathbf{q}} \right| \psi_{el}^{(0)} \right\rangle = F\left(\mathbf{q},-t\right). \end{aligned} \tag{5.45}$$

From (5.45), for the function

$$B\left(\mathbf{q},t\right) \equiv \operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t} F\left(\mathbf{q},t\right)\right] \tag{5.46}$$

the following equality is derived:

$$\begin{aligned} B\left(\mathbf{q},-t\right) &= \operatorname{Im}\left[e^{i\omega_{\mathrm{LO}}t} F\left(\mathbf{q},-t\right)\right] \\ &= \operatorname{Im}\left[e^{i\omega_{\mathrm{LO}}t} F^{*}\left(\mathbf{q},t\right)\right] \\ &= -\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t} F\left(\mathbf{q},t\right)\right] = -B\left(\mathbf{q},t\right), \end{aligned}$$

$$B(\mathbf{q},-t) = -B(\mathbf{q},t). \tag{5.47}$$

The integral in Eq. (5.39)

$$\operatorname{Im}\int_0^\infty dt e^{i\omega t-\delta t}\operatorname{Im}\left[e^{-i\omega_{\mathrm{LO}}t}F(\mathbf{q},t)\right] = \operatorname{Im}\int_0^\infty dt e^{i\omega t-\delta t}B(\mathbf{q},t)$$

is then transformed as follows:

$$\begin{aligned}
\operatorname{Im}\int_0^\infty dt e^{i\omega t-\delta t}B(\mathbf{q},t) &= \frac{1}{2i}\left[\int_0^\infty dt e^{i\omega t-\delta t}B(\mathbf{q},t) - \int_0^\infty dt e^{-i\omega t-\delta t}B(\mathbf{q},t)\right]\\
&= \frac{1}{2i}\left[\int_0^\infty dt e^{i\omega t-\delta t}B(\mathbf{q},t) - \int_{-\infty}^0 dt e^{i\omega t+\delta t}B(\mathbf{q},-t)\right]\\
&= \frac{1}{2i}\left[\int_0^\infty dt e^{i\omega t-\delta t}B(\mathbf{q},t) + \int_{-\infty}^0 dt e^{i\omega t+\delta t}B(\mathbf{q},t)\right]\\
&= \frac{1}{2i}\int_{-\infty}^\infty dt e^{i\omega t-\delta|t|}B(\mathbf{q},t)\\
&= \frac{1}{2i}\int_{-\infty}^\infty dt e^{i\omega t-\delta|t|}\frac{1}{2i}\left[e^{-i\omega_{\mathrm{LO}}t}F(\mathbf{q},t) - e^{i\omega_{\mathrm{LO}}t}F^*(\mathbf{q},t)\right]\\
&= -\frac{1}{4}\int_{-\infty}^\infty dt e^{i\omega t-\delta|t|}\left[e^{-i\omega_{\mathrm{LO}}t}F(\mathbf{q},t) - e^{i\omega_{\mathrm{LO}}t}F(\mathbf{q},-t)\right].
\end{aligned}$$

We can show that, as far as $\left|\psi_{el}^{(0)}\right\rangle$ is the *ground* state, the integral $\int_{-\infty}^\infty dt e^{i\omega t-\delta|t|}F(\mathbf{q},-t)$ for positive ω is equal to zero. Let $\left\{\left|\psi_{el}^{(n)}\right\rangle\right\}$ is the total basis set of the eigenfunctions of the Hamiltonian $\tilde{H}_e$. Using these functions we expand $F(\mathbf{q},t)$:

$$\begin{aligned}
F(\mathbf{q},t) &= \sum_n \left\langle\psi_{el}^{(0)}\left|e^{\frac{it}{\hbar}\tilde{H}_e}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_e}\right|\psi_{el}^{(n)}\right\rangle\left\langle\psi_{el}^{(n)}\left|\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\\
&= \sum_n \left|\left\langle\psi_{el}^{(n)}\left|\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\right|^2 e^{\frac{it}{\hbar}\left(\tilde{E}_0-\tilde{E}_n\right)},
\end{aligned}$$

$$\begin{aligned}
\int_{-\infty}^\infty dt e^{i\omega t-\delta|t|}F(\mathbf{q},-t) &= \sum_n \left|\left\langle\psi_{el}^{(n)}\left|\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\right|^2 \int_{-\infty}^\infty dt e^{i\omega t+\frac{i}{\hbar}\left(\tilde{E}_n-\tilde{E}_0\right)t-\delta|t|}\\
&= \sum_n \left|\left\langle\psi_{el}^{(n)}\left|\rho_{-\mathbf{q}}\right|\psi_{el}^{(0)}\right\rangle\right|^2 2\pi\delta\left(\omega+\frac{\tilde{E}_n-\tilde{E}_0}{\hbar}\right) = 0,
\end{aligned}$$

because for $\omega > 0$, $\omega + \frac{\tilde{E}_n-\tilde{E}_0}{\hbar}$ is never equal to zero.

So, rewriting expression (5.39) with the dynamic structure factor of the electron (or hole) gas results in:

$$\begin{aligned}
\operatorname{Re}\sigma_{3\mathrm{D}}(\omega) &= \frac{1}{6V\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}} q^2\left|V_{\mathbf{q}}\right|^2\int_{-\infty}^\infty dt e^{i(\omega-\omega_{\mathrm{LO}})t-\delta|t|}F(\mathbf{q},t),\\
\operatorname{Re}\sigma_{2\mathrm{D}}(\omega) &= \frac{1}{4A\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}} q^2\left|V_{\mathbf{q}}\right|^2\int_{-\infty}^\infty dt e^{i(\omega-\omega_{\mathrm{LO}})t-\delta|t|}F(\mathbf{q},t),
\end{aligned}$$

and we obtain

$$\operatorname{Re}\sigma_{\mathrm{3D}}(\omega) = \frac{n_0}{3\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}} q^2|V_{\mathbf{q}}|^2 S(\mathbf{q},\omega-\omega_{\mathrm{LO}}), \tag{5.48a}$$

$$\operatorname{Re}\sigma_{\mathrm{2D}}(\omega) = \frac{n_0}{2\hbar\omega^3}\frac{e^2}{m_b^2}\sum_{\mathbf{q}} q^2|V_{\mathbf{q}}|^2 S(\mathbf{q},\omega-\omega_{\mathrm{LO}}), \tag{5.48b}$$

where

$$n_0 = \begin{cases} N/V \text{ in 3D,} \\ N/A \text{ in 2D} \end{cases}$$

is the density of charge carriers.

For an isotropic medium, the dynamic structure factor does not depend on the direction of $\mathbf{q}$, so that $S(\mathbf{q},\omega) = S(q,\omega)$, where $q = |\mathbf{q}|$. Let us simplify the expression (5.48a) using explicitly the amplitudes of the Fröhlich electron-phonon interaction. The modulus squared of the Fröhlich electron-phonon interaction amplitude is given by

$$|V_{\mathbf{q}}|^2 = \begin{cases} \dfrac{(\hbar\omega_{\mathrm{LO}})^2}{q^2}\dfrac{4\pi\alpha}{V}\left(\dfrac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{1/2} & \text{in 3D} \\ \dfrac{(\hbar\omega_{\mathrm{LO}})^2}{q}\dfrac{2\pi\alpha}{A}\left(\dfrac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{1/2} & \text{in 2D,} \end{cases} \tag{5.49}$$

where α is the (dimensionless) Fröhlich coupling constant determining the coupling strength between the charge carriers and the longitudinal optical phonons [83, 84]. In 3D and 2D, respectively, the sums over $\mathbf{q}$ is transformed to the integrals as follows:

$$\text{3D: } \sum_{\mathbf{q}} \ldots = \frac{V}{(2\pi)^3}\int d\mathbf{q}\ldots$$

$$\text{2D: } \sum_{\mathbf{q}} \ldots = \frac{A}{(2\pi)^2}\int d\mathbf{q}\ldots$$

$$\Downarrow$$

$$\operatorname{Re}\sigma_{\mathrm{3D}}(\omega) = \frac{n_0}{3\hbar\omega^3}\frac{e^2}{m_b^2}\frac{V}{(2\pi)^3}\int d\mathbf{q}q^2\left|\frac{\hbar\omega_{\mathrm{LO}}}{iq}\left(\frac{4\pi\alpha}{V}\right)^{1/2}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{1/4}\right|^2 S(q,\omega-\omega_{\mathrm{LO}})$$

$$\Downarrow$$

$$\operatorname{Re}\sigma_{\mathrm{3D}}(\omega) = \frac{n_0e^2}{m_b^2}\frac{2\alpha}{3\pi}\frac{\hbar\omega_{\mathrm{LO}}^2}{\omega^3}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{1/2}\int_0^\infty q^2 S(q,\omega-\omega_{\mathrm{LO}})dq. \tag{5.51}$$

In the same way, we transform $\operatorname{Re}\sigma_{\mathrm{2D}}(\omega)$:

$$\operatorname{Re}\sigma_{\mathrm{2D}}(\omega) = \frac{n_0e^2}{m_b^2}\frac{\alpha}{2}\frac{\hbar\omega_{\mathrm{LO}}^2}{\omega^3}\left(\frac{\hbar}{2m_b\omega_{\mathrm{LO}}}\right)^{1/2}\int_0^\infty q^2 S(q,\omega-\omega_{\mathrm{LO}})dq.$$

Using the Feynman units ($\hbar = 1$, $m_b = 1$, $\omega_{\mathrm{LO}} = 1$), $\mathrm{Re}\,\sigma(\omega)$ is

$$\mathrm{Re}\,\sigma_{3\mathrm{D}}(\omega) = n_0 e^2 \frac{\sqrt{2}\alpha}{3\pi} \frac{1}{\omega^3} \int_0^\infty q^2 S(q, \omega - 1) dq, \tag{5.52}$$

$$\mathrm{Re}\,\sigma_{2\mathrm{D}}(\omega) = n_0 e^2 \frac{\alpha}{2\sqrt{2}} \frac{1}{\omega^3} \int_0^\infty q^2 S(q, \omega - 1) dq. \tag{5.53}$$

From these expressions, it is clear that the scaling relation

$$\mathrm{Re}\,\sigma_{2\mathrm{D}}(\omega, \alpha) = \mathrm{Re}\,\sigma_{3\mathrm{D}}(\omega, \frac{3\pi}{4}\alpha) \tag{5.54}$$

which holds for the one-polaron case introduced in ref. [83, 84], is also valid for the many-polaron case if the corresponding 2D or 3D dynamic structure factor is used.

1. *Calculation of the dynamic structure factor using the retarded Green's functions*

The dynamic structure factor $S(\mathbf{q}, \omega)$ is expressed through the two-point correlation function by Eq. (5.43). The correlation function $F(\mathbf{q}, \omega)$ can be found using the retarded Green's function of the density operators

$$G^R(\mathbf{q}, t) = -i\Theta(t) \left\langle \psi_{el}^{(0)} \middle| \left[e^{\frac{it}{\hbar}\tilde{H}_0} \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0}, \rho_{-\mathbf{q}} \right] \middle| \psi_{el}^{(0)} \right\rangle, \tag{5.55}$$

where $\Theta(t)$ is the step function. Let us consider the more general case of a finite temperature,

$$G^R(\mathbf{q}, t) = -i\Theta(t) \left\langle \left[e^{\frac{it}{\hbar}\tilde{H}_0} \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0}, \rho_{-\mathbf{q}} \right] \right\rangle, \tag{5.56}$$

where the average is

$$\langle \ldots \rangle \equiv \frac{Tr\left(e^{-\beta\tilde{H}_0} \ldots\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}, \ \beta = \frac{1}{k_B T}. \tag{5.57}$$

The Fourier image $G^R(\mathbf{q}, \omega)$ of the retarded Green's function $G^R(\mathbf{q}, t)$ is

$$\begin{aligned} G^R(\mathbf{q}, \omega) &= \int_{-\infty}^\infty G^R(\mathbf{q}, t) e^{i\omega t} dt \\ &= -i \int_0^\infty \left\langle \left[e^{\frac{it}{\hbar}\tilde{H}_0} \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0}, \rho_{-\mathbf{q}} \right] \right\rangle e^{i\omega t} dt \\ &= -i \int_0^\infty \left(\left\langle e^{\frac{it}{\hbar}\tilde{H}_0} \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0} \rho_{-\mathbf{q}} \right\rangle - \left\langle \rho_{-\mathbf{q}} e^{\frac{it}{\hbar}\tilde{H}_0} \rho_{\mathbf{q}} e^{-\frac{it}{\hbar}\tilde{H}_0} \right\rangle \right) e^{i\omega t} dt \end{aligned}$$

The imaginary part of $G^R(\mathbf{q},\omega)$ then is

$$\begin{aligned}
\operatorname{Im} G^R(\mathbf{q},\omega) &= -\operatorname{Re}\int_0^\infty \left(\left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle - \left\langle \rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right\rangle\right) e^{i\omega t}dt \\
&= -\frac{1}{2}\int_0^\infty \left(\left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle - \left\langle \rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right\rangle\right) e^{i\omega t}dt \\
&- \frac{1}{2}\int_0^\infty \left(\left\langle \rho_{\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right\rangle - \left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}\right\rangle\right) e^{-i\omega t}dt \\
&= -\frac{1}{2}\int_{-\infty}^\infty \left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle e^{i\omega t}dt + \frac{1}{2}\int_{-\infty}^\infty \left\langle \rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right\rangle e^{i\omega t}dt.
\end{aligned}$$

In the second integral here, we replace t by $(t' + i\hbar\beta)$:

$$\begin{aligned}
&\int_{-\infty}^\infty \frac{Tr\left(e^{-\beta\tilde{H}_0}\rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}e^{i\omega t}dt \\
&= \int_{-\infty-i\hbar\beta}^{\infty-i\hbar\beta} \frac{Tr\left(e^{-\beta\tilde{H}_0}\rho_{-\mathbf{q}}e^{\frac{it'}{\hbar}\tilde{H}_0-\beta\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it'}{\hbar}\tilde{H}_0+\beta\tilde{H}_0}\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}e^{i\omega(t'+i\hbar\beta)}dt' \\
&= e^{-\beta\hbar\omega}\int_{-\infty-i\hbar\beta}^{\infty-i\hbar\beta} \frac{Tr\left(e^{\frac{it'}{\hbar}\tilde{H}_0-\beta\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it'}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}e^{i\omega t'}dt'.
\end{aligned}$$

As far as the integral over t converges (i. e., $\left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle$ tends to zero at $|t| \to \infty$), we can shift the integration contour to the real axis, what gives us the result

$$\begin{aligned}
&\int_{-\infty}^\infty \frac{Tr\left(e^{-\beta\tilde{H}_0}\rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}e^{i\omega t}dt \\
&= e^{-\beta\hbar\omega}\int_{-\infty}^\infty \frac{Tr\left(e^{\frac{it}{\hbar}\tilde{H}_0-\beta\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right)}{Tr\left(e^{-\beta\tilde{H}_0}\right)}e^{i\omega t}dt, \\
&\Updownarrow \\
&\int_{-\infty}^\infty \left\langle \rho_{-\mathbf{q}}e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\right\rangle e^{i\omega t}dt \\
&= e^{-\beta\hbar\omega}\int_{-\infty}^\infty \left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle e^{i\omega t}dt.
\end{aligned}$$

Herefrom, we find that

$$\operatorname{Im} G^R(\mathbf{q},\omega) = -\frac{1}{2}\left(1 - e^{-\beta\hbar\omega}\right)\int_{-\infty}^\infty \left\langle e^{\frac{it}{\hbar}\tilde{H}_0}\rho_{\mathbf{q}}e^{-\frac{it}{\hbar}\tilde{H}_0}\rho_{-\mathbf{q}}\right\rangle e^{i\omega t}dt$$

$$\Downarrow$$

$$\operatorname{Im} G^R(\mathbf{q},\omega) = -\frac{1}{2}\left(1 - e^{-\beta\hbar\omega}\right)F(\mathbf{q},\omega).$$

So, the equation follows from the analytical properties of the Green's functions:

$$F(\mathbf{q},\omega) = -\frac{2\,\mathrm{Im}\,G^R(\mathbf{q},\omega)}{1-e^{-\beta\hbar\omega}}. \tag{5.58}$$

The formula (5.58) is related to arbitrary temperatures. In the zero-temperature limit ($\beta \to \infty$), the factor $\left(1-e^{-\beta\hbar\omega}\right)^{-1}$ (5.58) turns into the Heavicide step function $\Theta(\omega)$, what leads to the formula

$$F(\mathbf{q},\omega)|_{T=0} = -2\Theta(\omega)\,\mathrm{Im}\,G^R(\mathbf{q},\omega)\big|_{T=0} \tag{5.59}$$

$$\Downarrow$$

$$S(\mathbf{q},\omega)|_{T=0} = -\frac{1}{N}\Theta(\omega)\,\mathrm{Im}\,G^R(\mathbf{q},\omega)\big|_{T=0} \tag{5.60}$$

The retarded Green function is related to the dielectric function of the electron gas by the following equation:

$$G^R(\mathbf{q},\omega) = \frac{1}{v_{\mathbf{q}}}\left[\frac{1}{\varepsilon(\mathbf{q},\omega)}-1\right]. \tag{5.61}$$

Within the random phase approximation (RPA), following [70], the expression for $G^R(\mathbf{q},\omega)$ is

$$G^R(\mathbf{q},\omega) = \left[1-v_{\mathbf{q}}P(\mathbf{q},\omega)\right]^{-1}\hbar P(\mathbf{q},\omega), \tag{5.62}$$

where the polarization function $P(\mathbf{q},\omega)$ is (see, e. g., p. 434 of [70])

$$P(\mathbf{q},\omega) = \frac{1}{\hbar}\sum_{\mathbf{k},\sigma}\frac{f_{\mathbf{k}+\mathbf{q},\sigma}-f_{\mathbf{k},\sigma}}{\omega-\frac{\hbar\mathbf{k}^2}{2m_b}+\frac{\hbar(\mathbf{k}+\mathbf{q})^2}{2m_b}+i\delta},\ \ \delta\to+0 \tag{5.63}$$

with the Fermi distribution function of electrons $f_{\mathbf{k},\sigma}$.

For a finite temperature, the explicit analytic expression for the imaginary part of the structure factor $P_{\mathrm{3D}}(\mathbf{q},\omega)$ is obtained (see [70]),

$$\begin{aligned}\mathrm{Im}\,P_{\mathrm{3D}}(\mathbf{q},\omega) &= \frac{Vm_b^2}{2\pi\hbar^4\beta q}\ln\frac{1+\exp\left\{\beta\left[\mu-E^{(+)}(q,\omega)\right]\right\}}{1+\exp\left\{\beta\left[\mu-E^{(-)}(q,\omega)\right]\right\}},\\ E^{(\pm)}(q,\omega) &\equiv \frac{\left(\hbar\omega\pm\frac{\hbar^2q^2}{2m_b}\right)^2}{4\frac{\hbar^2q^2}{2m_b}},\end{aligned} \tag{5.64}$$

with the chemical potential μ. The real part of the structure factor is obtained using the Kramers-Kronig dispersion relation:

$$\mathrm{Re}\,P(\mathbf{q},\omega) = \frac{1}{\pi}\int_{-\infty}^{\infty}\mathcal{P}\left(\frac{1}{\omega'-\omega}\right)\mathrm{Im}\,P(\mathbf{q},\omega')\,d\omega'. \tag{5.65}$$

Analytical expressions for both real and imaginary parts of $P\left(\mathbf{q},\omega\right)$ can be written down for the zero temperature (see [70]),

$$\operatorname{Re}P_{\mathrm{3D}}\left(q,\omega\right)=-\frac{Vm_b}{4\pi^2\hbar^2q}\left\{\begin{array}{c}\left[k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega-\frac{\hbar q^2}{2m_b}\right)^2\right]\ln\left|\frac{\omega-\frac{\hbar q^2}{2m_b}-\frac{\hbar k_F q}{m_b}}{\omega-\frac{\hbar q^2}{2m_b}+\frac{\hbar k_F q}{m_b}}\right|\\+\left[k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega+\frac{\hbar q^2}{2m_b}\right)^2\right]\ln\left|\frac{\omega+\frac{\hbar q^2}{2m_b}+\frac{\hbar k_F q}{m_b}}{\omega+\frac{\hbar q^2}{2m_b}-\frac{\hbar k_F q}{m_b}}\right|\\+2k_Fq\end{array}\right\},$$

$$\operatorname{Im}P_{\mathrm{3D}}\left(q,\omega\right)=-\frac{Vm_b}{4\pi\hbar^2q}\left\{\begin{array}{c}\left[k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega-\frac{\hbar q^2}{2m_b}\right)^2\right]\Theta\left(k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega-\frac{\hbar q^2}{2m_b}\right)^2\right)\\-\left[k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega+\frac{\hbar q^2}{2m_b}\right)^2\right]\Theta\left(k_F^2-\frac{m_b^2}{\hbar^2q^2}\left(\omega+\frac{\hbar q^2}{2m_b}\right)^2\right)\end{array}\right\},\tag{5.66}$$

where $k_F=\left(3\pi^2N/V\right)^{1/3}$ is the Fermi wave number.

After substituting into Eq. (5.58) the retarded Green's function (5.62) in terms of the polarization function we arrive at the formula

$$F\left(\mathbf{q},\omega\right)=-\frac{2\hbar}{1-e^{-\beta\hbar\omega}}\operatorname{Im}\frac{P\left(\mathbf{q},\omega\right)}{1-v_{\mathbf{q}}P\left(\mathbf{q},\omega\right)}$$

$$\Downarrow$$

$$S\left(\mathbf{q},\omega\right)=-\frac{\hbar}{N\left(1-e^{-\beta\hbar\omega}\right)}\frac{\operatorname{Im}P\left(\mathbf{q},\omega\right)}{\left[1-v_{\mathbf{q}}\operatorname{Re}P\left(\mathbf{q},\omega\right)\right]^2+\left[v_{\mathbf{q}}\operatorname{Im}P\left(\mathbf{q},\omega\right)\right]^2},\tag{5.67}$$

$$\left.S\left(\mathbf{q},\omega\right)\right|_{T=0}=-\frac{\hbar}{N}\Theta\left(\omega\right)\frac{\operatorname{Im}P\left(\mathbf{q},\omega\right)}{\left[1-v_{\mathbf{q}}\operatorname{Re}P\left(\mathbf{q},\omega\right)\right]^2+\left[v_{\mathbf{q}}\operatorname{Im}P\left(\mathbf{q},\omega\right)\right]^2}.\tag{5.68}$$

With this dynamic structure factor, the optical conductivity (5.52) (in the Feynman units) takes the form

$$\begin{aligned}\operatorname{Re}\sigma_{\mathrm{3D}}(\omega)=&-e^2\frac{\sqrt{2}\alpha}{3\pi V}\frac{1}{\omega^3}\Theta\left(\omega-1\right)\\&\times\int_0^\infty\frac{\operatorname{Im}P_{\mathrm{3D}}\left(q,\omega-1\right)}{\left[1-v_q\operatorname{Re}P_{\mathrm{3D}}\left(q,\omega-1\right)\right]^2+\left[v_q\operatorname{Im}P_{\mathrm{3D}}\left(q,\omega-1\right)\right]^2}q^2dq.\end{aligned}\tag{5.69}$$

Correspondingly, in the 2D case we obtain the expression

$$\begin{aligned}\operatorname{Re}\sigma_{\mathrm{2D}}(\omega)=&-e^2\frac{\alpha}{2\sqrt{2}V}\frac{1}{\omega^3}\Theta\left(\omega-1\right)\\&\times\int_0^\infty\frac{\operatorname{Im}P_{\mathrm{2D}}\left(q,\omega-1\right)}{\left[1-v_q\operatorname{Re}P_{\mathrm{2D}}\left(q,\omega-1\right)\right]^2+\left[v_q\operatorname{Im}P_{\mathrm{2D}}\left(q,\omega-1\right)\right]^2}q^2dq.\end{aligned}\tag{5.70}$$

2. Plasmon-phonon contribution

The RPA dynamic structure factor for the electron (or hole) system can be separated in two parts, one related to continuum excitations of the electrons (or holes) S_{cont}, and one related to the undamped plasmon branch:

$$S_{\text{RPA}}(q,\omega) = A_{\text{pl}}(q)\delta\left(\omega - \omega_{\text{pl}}\left(q\right)\right) + S_{\text{cont}}(q,\omega), \tag{5.71}$$

where $\omega_{\text{pl}}\left(q\right)$ is the wave number dependent plasmon frequency and A_{pl} is the strength of the undamped plasmon branch.

In Eqs. (5.69), (5.70), the contribution of the continuum excitations corresponds to the region (q,ω) where $\operatorname{Im} P\left(q,\omega\right) \neq 0$. The contribution related to the undamped plasmons is provided by a region of $\left(q,\omega\right)$, where the equations

$$\left\{\begin{array}{c} \operatorname{Im} P\left(q,\omega\right) = 0 \\ 1 - v_q \operatorname{Re} P\left(q,\omega\right) = 0 \end{array}\right. . \tag{5.72}$$

are fulfilled simultaneously. Using (5.61), we find that Eqs. (5.73) are equivalent to

$$\operatorname{Im} \frac{1}{\varepsilon\left(q,\omega\right)} = 0, \quad \operatorname{Re} \frac{1}{\varepsilon\left(q,\omega\right)} = 0. \tag{5.73}$$

In the region where $\operatorname{Im} P\left(\mathbf{q},\omega\right) = 0$, the expression $\frac{\operatorname{Im} P(q,\omega)}{\left[1 - v_q \operatorname{Re} P(q,\omega)\right]^2 + \left[v_q \operatorname{Im} P(q,\omega)\right]^2}$ is proportional to the delta function, which gives a finite contribution to the memory function after the integration over q:

$$\left.\frac{\operatorname{Im} P\left(q,\omega\right)}{\left[1 - v_q \operatorname{Re} P\left(q,\omega\right)\right]^2 + \left[v_q \operatorname{Im} P\left(q,\omega\right)\right]^2}\right|_{\operatorname{Im} P(q,\omega)=0}$$
$$= \frac{1}{\pi v_q}\delta\left(1 - v_q \operatorname{Re} P\left(q,\omega\right)\right). \tag{5.74}$$

Using Eq. (5.74), the coefficients $A_{\text{pl}}(q)$ in Eq. (5.71) can be expressed in terms of the polarization function $P\left(q,\omega\right)$ as follows:

$$\left.\frac{\operatorname{Im} P\left(q,\omega\right)}{\left[1 - v_q \operatorname{Re} P\left(q,\omega\right)\right]^2 + \left[v_q \operatorname{Im} P\left(q,\omega\right)\right]^2}\right|_{\operatorname{Im} P(q,\omega)=0}$$
$$= \left.\frac{1}{\pi v_q^2 \left|\frac{\partial}{\partial\omega} \operatorname{Re} P\left(q,\omega\right)\right|}\right|_{\omega=\omega_{\text{pl}}(q)} \delta\left(\omega - \omega_{\text{pl}}\left(q\right)\right)$$

$$\Downarrow$$

$$A_{\mathrm{pl}}(q) = \left.\frac{1}{\pi v_q^2 \left|\frac{\partial}{\partial\omega}\operatorname{Re}P(q,\omega)\right|}\right|_{\omega=\omega_{\mathrm{pl}}(q)}. \tag{5.75}$$

The contribution derived from the undamped plasmon branch $A_{\mathrm{pl}}(q)\delta(\omega-\omega_{\mathrm{pl}}(q))$ is denoted in Ref. [51] as the *'plasmon-phonon' contribution*. The physical process related to this contribution is the emission of both a phonon and a plasmon in the scattering process.

E. Comparison to the infrared spectrum of $\mathbf{Nd}_{2-x}\mathbf{Ce}_x\mathbf{CuO}_{2-y}$

Calvani and collaborators have performed doping-dependent measurements of the infrared absorption spectra of the high-T$_c$ material $\mathrm{Nd}_{2-x}\mathrm{Ce}_x\mathrm{CuO}_{2-y}$ (NCCO). The region of the spectrum examined by these authors (50-10000 cm^{-1}) is very rich in absorption features: they observe is a "Drude-like" component at the lowest frequencies, and a set of sharp absorption peaks related to phonons and infrared active modes (up to about 1000 cm^{-1}) possibly associated to small (Holstein) polarons. Three distinct absorption bands can be distinguished: the 'd-band' (around 1000 cm^{-1}), the Mid-Infrared band (MIR, around 5000 cm^{-1}) and the Charge-Transfer band (around 10^4 cm^{-1}). Of all these features, the d-band and, at a higher temperatures, the Drude-like component have (hypothetically) been associated with large polaron optical absorption [90].

For the lowest levels of Ce doping, the d-band can be most clearly distinguished from the other features. The experimental optical absorption spectrum (up to 3000 cm^{-1}) of $\mathrm{Nd}_2\mathrm{CuO}_{2-\delta}$ ($\delta < 0.004$), obtained by Calvani and co-workers [90], is shown in Fig. 12 (shaded area) together with the theoretical curve obtained by the present method (full, bold curve) and, for reference, the one-polaron optical absorption result (dotted curve). At lower frequencies (600-1000 cm^{-1}) a marked difference between the single polaron optical absorption and the many-polaron result is manifest. The experimental d-band can be clearly identified, rising in intensity at about 600 cm^{-1}, peaking around 1000 cm^{-1}, and then decreasing in intensity above that frequency. At a density of $n_0 = 1.5\times10^{17}$ cm^{-3}, we found a remarkable agreement between our theoretical predictions and the experimental curve.



FIG. 12: The infrared absorption of $Nd_2CuO_{2-\delta}$ ($\delta < 0.004$) is shown as a function of frequency, up to 3000 cm^{-1}. The experimental results of Calvani and co-workers [90] is represented by the thin black curve and by the shaded area. The so-called 'd-band' rises in intensity around 600 cm^{-1} and increases in intensity up to a maximum around 1000 cm^{-1}. The dotted curve shows the single polaron result. The full black curve represents the theoretical results obtained in the present work for the interacting many-polaron gas with $n_0 = 1.5 \times 10^{17}$ cm^{-3}, $\alpha = 2.1$ and $m_b = 0.5\ m_e$. (From Ref. [51].)

F. Experimental data on the optical absorption in manganites: interpretation in terms of a many-polaron response

In Refs. [91, 92], the experimental results on the optical spectroscopy of $La_{2/3}Sr_{1/3}MnO_3$ (LSMO) and $La_{2/3}Ca_{1/3}MnO_3$ (LCMO) thin films in the mid-infrared frequency region are presented. The optical conductivity spectra of LCMO films are interpreted in [91, 92] in terms of the optical response of small polarons, while the optical conductivity spectra of LSMO films are explained using the large-polaron picture (see Fig. 13).

The real part of the optical conductivity $\operatorname{Re}\sigma(\omega)$ is expressed in Ref. [92] by the formula

$$\operatorname{Re}\sigma(\omega) = \alpha n_p \frac{2}{3}\frac{e^2}{m^2}\frac{(\hbar\omega_0)^2}{\pi\hbar\omega^3}\sqrt{\frac{\hbar}{2m\omega_0}}\int_0^\infty q^2 S^{(\mathrm{H})}(q,\omega-\omega_0)dq, \tag{5.76a}$$

where α is the electron-phonon coupling constant, n_p is the polaron density, m is the electron band mass, ω_0 is the LO-phonon frequency, and $S^{(\mathrm{H})}(q,\omega)$ is the dynamic structure factor,



FIG. 13: Comparison of the low-temperature MIR optical conductivity to $\sigma(\omega)$ from various model calculations: the solid line refers to the weak-coupling approach of Tempere and Devreese [51] modified for an on-site Hubbard interaction, the dashed line is the result of the phenomenological approach for self-trapped large polarons by Emin [93], the dotted curve is the weak-coupling single-polaron result [69]. (From Ref. [91].)

determined in Ref. [92] through the dielectric function of an electron gas $\varepsilon(\mathbf{q},\omega)$:

$$S^{(\mathrm{H})}(\mathbf{q},\omega)=\frac{\hbar}{n_p}\frac{q^2}{4\pi e^2}\,\mathrm{Im}\left[-\frac{1}{\varepsilon(\mathbf{q},\omega)}\right]. \tag{5.77}$$

In Ref. [51], the other definition for the dynamic structure factor is used, which is equivalent to that given by (5.77) (see Ref. [70]) with the factor N (the number of electrons)

$$S^{(\mathrm{TD})}(\mathbf{q},\omega)=\frac{1}{2}\int_{-\infty}^{+\infty}\left\langle\varphi_{\mathrm{el}}\left|\rho_{\mathbf{q}}(t)\,\rho_{-\mathbf{q}}(0)\right|\varphi_{\mathrm{el}}\right\rangle e^{i\omega t}dt=NS^{(\mathrm{H})}(\mathbf{q},\omega). \tag{5.78}$$

Here, φ_{el} denotes the ground state of the electron subsystem (without the electron-phonon interaction), $\rho_{\mathbf{q}}=\sum_{j=1}^{N}e^{i\mathbf{q}\cdot\mathbf{r}_j}$ is the Fourier component of the electron density.

In Ref. [51], the dynamic structure factor is calculated within the random-phase approximation (RPA) taking into account the Coulomb interaction between electrons with the Fourier component of the Coulomb potential

$$v_{\mathbf{q}}=\frac{4\pi e^2}{q^2\varepsilon_\infty}, \tag{5.79}$$

where ε_∞ is the high-frequency dielectric constant of the crystal. In Refs. [91, 92], the dynamic structure factor is calculated taking into account the local Hubbard electron-electron interaction instead of the Coulomb interaction. The local Hubbard interaction is used in the small-polaron formalism and describes the potential energy of two electrons on one and the same site (see, e. g., Refs. [94, 95]). In its simplest form the Hubbard interaction is (see Eq. (1) of Ref. [94])

$$V_H = U \sum_i n_{i\uparrow} n_{i\downarrow}, \tag{5.80}$$

where U is the coupling constant of the Hubbard interaction, $n_{i\uparrow}$ is the electron occupation number for the i-th site.

In Refs. [91, 92], there are no details of the calculation using the interaction term (5.80). The following procedure can be supposed. The transition from the summation over the lattice sites to the integral over the crystal volume V is performed taking into account the normalization condition

$$N_{\uparrow(\downarrow)} = \sum_i n_{i\uparrow(\downarrow)} = \int_V \tilde{n}_{\uparrow(\downarrow)}(\mathbf{r})\, d\mathbf{r},$$

where the density $\tilde{n}_{\uparrow(\downarrow)}(\mathbf{r})$ is to be determined through $n_{i\uparrow(\downarrow)}$. As far as the lattice cell volume $\Omega_0 \ll V$, the integral $\int_V n_{\uparrow(\downarrow)}(\mathbf{r})\, d\mathbf{r}$ can be written as the sum over the lattice sites:

$$\int_V n_{\uparrow(\downarrow)}(\mathbf{r})\, d\mathbf{r} = \Omega_0 \sum_i \tilde{n}_{\uparrow(\downarrow)}(\mathbf{r}_i),$$

where $\{\mathbf{r}_i\}$ are the vectors of the lattice. Therefore, from the equality

$$\sum_i n_{i\uparrow(\downarrow)} = \Omega_0 \sum_i \tilde{n}_{\uparrow(\downarrow)}(\mathbf{r}_i)$$

we find that

$$n_{i\uparrow(\downarrow)} = \tilde{n}_{\uparrow(\downarrow)}(\mathbf{r}_i).$$

The potential (5.80) is then transformed from the sum over sites to the integral:

$$V_H = \frac{U}{\Omega_0} \sum_i n_{i\uparrow} n_{i\downarrow} \Omega_0 = \int_V U\Omega_0 \delta(\mathbf{r} - \mathbf{r}')\, \tilde{n}_\uparrow(\mathbf{r})\, \tilde{n}_\downarrow(\mathbf{r}')\, d\mathbf{r} d\mathbf{r}'. \tag{5.81}$$

Consequently, in the continuum approach the Hubbard model is described by the δ-like interparticle potential $U\Omega_0\delta(\mathbf{r} - \mathbf{r}')$.

This development of the approach [51] performed in Refs. [91, 92] seems to be contradictory by the following reason. For a many-polaron system, both the electron-phonon and electron-electron interactions are provided by the electrostatic potentials. Therefore, it would be consistent to consider them both within one and the same approach. Namely, the Coulomb electron-electron interaction with the potential (5.79) is relevant for large and small polarons with the Fröhlich electron-phonon interaction, while the Hubbard electron-electron interaction is relevant for small Holstein polarons. Nevertheless, as recognized in Ref. [91], this model "reproduces the observed shape of the polaron peak quite convincingly" and provides a better agreement with the experiment [91, 92] than the phenomenological approach [93] and the one-polaron theory [69].

VI. INTERACTING POLARONS IN A QUANTUM DOT

A. The partition function and the free energy of a many-polaron system

We consider a system of N electrons with mutual Coulomb repulsion and interacting with the lattice vibrations following Ref. [96]. The system is assumed to be confined by a parabolic potential characterized by the frequency parameter Ω_0. The total number of electrons is represented as $N = \sum_\sigma N_\sigma$, where N_σ is the number of electrons with the spin projection $\sigma = \pm 1/2$. The electron 3D (2D) coordinates are denoted by $\mathbf{x}_{j,\sigma}$ with $j = 1, \cdots, N_\sigma$. The bulk phonons (characterized by 3D wave vectors $\mathbf{k}$ and frequencies $\omega_\mathbf{k}$) are described by the complex coordinates $Q_\mathbf{k}$, which possess the property [41]

$$Q_\mathbf{k}^* = Q_{-\mathbf{k}}. \tag{6.1}$$

The full set of the electron and phonon coordinates are denoted by $\bar{\mathbf{x}} \equiv \{\mathbf{x}_{j,\sigma}\}$ and $\bar{Q} \equiv \{Q_\mathbf{k}\}$.

Throughout the present treatment, the Euclidean time variable $\tau = it$ is used, where t is the real time variable. In this representation the Lagrangian of the system is

$$L\left(\dot{\bar{\mathbf{x}}}, \dot{\bar{Q}}; \bar{\mathbf{x}}, \bar{Q}\right) = L_e\left(\dot{\bar{\mathbf{x}}}, \bar{\mathbf{x}}\right) - V_C\left(\bar{\mathbf{x}}\right) - U_b\left(\bar{\mathbf{x}}\right) + L_{ph}\left(\dot{\bar{Q}}, \bar{Q}\right) + L_{e-ph}\left(\bar{\mathbf{x}}, \bar{Q}\right), \tag{6.2}$$

where $L_e\left(\dot{\bar{\mathbf{x}}}, \bar{\mathbf{x}}\right)$ is the Lagrangian of an electron with band mass m_b in a quantum dot:

$$L_e\left(\dot{\bar{\mathbf{x}}}, \bar{\mathbf{x}}\right) = -\sum_{\sigma=\pm 1/2} \sum_{j=1}^{N_\sigma} \frac{m_b}{2}\left(\dot{\mathbf{x}}_{j,\sigma}^2 + \Omega_0^2 \mathbf{x}_{j,\sigma}^2\right), \qquad \dot{\mathbf{x}} \equiv \frac{d\mathbf{x}}{d\tau}, \tag{6.3}$$

Ω_0 is the confinement frequency, $V_b\left(\bar{\mathbf{x}}\right)$ is the potential of a background charge (supposed to be static and uniformly distributed with the charge density en_b in a sphere of a radius R),

$$U_b\left(\bar{\mathbf{x}}\right) = \sum_\sigma \sum_{j=1}^{N} V_b\left(r_{j,\sigma}\right), \ r \equiv |\mathbf{x}|, \tag{6.4}$$

$$V_b(r) = -\frac{4\pi e^2 n_b}{3\varepsilon_0}\left[\Theta(r < R)\frac{3R^2 - r^2}{2} + \Theta(R \leq r)\frac{R^3}{r}\right], \tag{6.5}$$

where ε_0 is the static dielectric constant of a crystal, $V_C\left(\bar{\mathbf{x}}\right)$ is the potential energy of the electron-electron Coulomb repulsion in the medium with the high-frequency dielectric constant ε_∞:

$$V_C\left(\bar{\mathbf{x}}\right) = \sum_{\sigma,\sigma'=\pm 1/2} \sum_{\substack{j=1 \\ (j,\sigma)\neq(l,\sigma')}}^{N_\sigma} \sum_{l=1}^{N_{\sigma'}} \frac{e^2}{2\varepsilon_\infty} \frac{1}{\left|\mathbf{x}_{j,\sigma} - \mathbf{x}_{l,\sigma'}\right|}, \tag{6.6}$$

$L_{ph}\left(\dot{\bar{Q}},\dot{\bar{Q}}^{*};\bar{Q},\bar{Q}^{*}\right)$ is the Lagrangian of free phonons:

$$L_{ph}\left(\dot{\bar{Q}},\bar{Q}\right)=-\frac{1}{2}\sum_{\mathbf{k}}(\dot{Q}_{\mathbf{k}}^{*}\dot{Q}_{\mathbf{k}}+\omega_{\mathbf{k}}^{2}Q_{\mathbf{k}}^{*}Q_{\mathbf{k}}),\qquad \dot{Q}\equiv\frac{dQ}{d\tau}. \tag{6.7}$$

Further, $L_{e-ph}\left(\bar{\mathbf{x}},\bar{Q},\bar{Q}^{*}\right)$ is the Lagrangian of the electron-phonon interaction:

$$L_{e-ph}\left(\bar{\mathbf{x}},\bar{Q}\right)=-\sum_{\mathbf{k}}\left(\frac{2\omega_{\mathbf{k}}}{\hbar}\right)^{1/2}V_{\mathbf{k}}Q_{-\mathbf{k}}\rho_{\mathbf{k}}, \tag{6.8}$$

where $\rho_{\mathbf{k}}$ is the Fourier transform of the electron density operator:

$$\rho_{\mathbf{k}}=\sum_{\sigma=\pm1/2}\sum_{j=1}^{N_{\sigma}}e^{i\mathbf{k}\cdot\mathbf{x}_{j,\sigma}}. \tag{6.9}$$

$V_{\mathbf{k}}$ is the amplitude of the electron-phonon interaction. Here, we consider electrons interacting with the long-wavelength longitudinal optical (LO) phonons with a dispersionless frequency $\omega_{\mathbf{k}}=\omega_{\mathrm{LO}}$, for which the amplitude $V_{\mathbf{k}}$ is [24]

$$V_{\mathbf{k}}=\frac{\hbar\omega_{\mathrm{LO}}}{q}\left(\frac{2\sqrt{2}\pi\alpha}{V}\right)^{1/2}\left(\frac{\hbar}{m_{b}\omega_{\mathrm{LO}}}\right)^{1/4}, \tag{6.10}$$

where α is the electron-phonon coupling constant and V is the volume of the crystal.

We consider a *canonical* ensemble, where the numbers N_{σ} are fixed. The partition function $Z\left(\{N_{\sigma}\},\beta\right)$ of the system can be expressed as a path integral over the electron and phonon coordinates:

$$Z\left(\{N_{\sigma}\},\beta\right)=\sum_{P}\frac{(-1)^{\xi_{P}}}{N_{1/2}!N_{-1/2}!}\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}\left(\tau\right)\int d\bar{Q}\int_{\bar{Q}}^{\bar{Q}}D\bar{Q}\left(\tau\right)e^{-S\left[\bar{\mathbf{x}}(\tau),\bar{Q}(\tau)\right]}, \tag{6.11}$$

where $S\left[\bar{\mathbf{x}}\left(\tau\right),\bar{Q}\left(\tau\right)\right]$ is the "action" functional:

$$S\left[\bar{\mathbf{x}}\left(\tau\right),\bar{Q}\left(\tau\right)\right]=-\frac{1}{\hbar}\int_{0}^{\hbar\beta}L\left(\dot{\bar{\mathbf{x}}},\dot{\bar{Q}};\bar{\mathbf{x}},\bar{Q}\right)d\tau. \tag{6.12}$$

The parameter $\beta\equiv1/\left(k_{B}T\right)$ is inversely proportional to the temperature T. In order to take the Fermi-Dirac statistics into account, the integral over the electron paths $\{\bar{\mathbf{x}}\left(\tau\right)\}$ in Eq. (6.11) contains a sum over all permutations P of the electrons with the same spin projection, and ξ_{P} denotes the parity of a permutation P.

The action functional (6.12) is quadratic in the phonon coordinates $\bar{Q}$. Therefore, *the path integral over the phonon variables* in $Z\left(\{N_{\sigma}\},\beta\right)$ *can be calculated analytically* following

Ref. [41]. Let us describe this path integration in detail. First, we introduce the real phonon coordinates through the real and imaginary parts of the complex phonon coordinates $Q'_{\mathbf{k}} \equiv \operatorname{Re} Q_{\mathbf{k}}$, $Q''_{\mathbf{k}} \equiv \operatorname{Im} Q_{\mathbf{k}}$. According to the symmetry property (6.1), they obey the equalities

$$Q'_{-\mathbf{k}} = Q'_{\mathbf{k}},\ Q''_{-\mathbf{k}} = -Q''_{\mathbf{k}}. \tag{6.13}$$

$$q_{\mathbf{k}} \equiv \begin{cases} \sqrt{2} Q'_{\mathbf{k}},\ k_x \geq 0, \\ \sqrt{2} Q''_{\mathbf{k}},\ k_x < 0. \end{cases} \tag{6.14}$$

In this representation, the sum over phonon coordinates $\sum_{\mathbf{k}} |Q_{\mathbf{k}}|^2$ is transformed in the following way using the symmetry property (6.13):

$$\begin{aligned} \sum_{\mathbf{k}} |Q_{\mathbf{k}}|^2 &= \sum_{\mathbf{k}} \left[(Q'_{\mathbf{k}})^2 + (Q''_{\mathbf{k}})^2 \right] \\ &= 2 \sum_{\substack{\mathbf{k} \\ (k_x \geq 0)}} (Q'_{\mathbf{k}})^2 + 2 \sum_{\substack{\mathbf{k} \\ (k_x < 0)}} (Q''_{\mathbf{k}})^2 \\ &= \sum_{\substack{\mathbf{k} \\ (k_x \geq 0)}} q_{\mathbf{k}}^2 + \sum_{\substack{\mathbf{k} \\ (k_x < 0)}} q_{\mathbf{k}}^2 = \sum_{\mathbf{k}} q_{\mathbf{k}}^2. \end{aligned}$$

Therefore, the phonon Lagrangian (6.7) with the real phonon coordinates is

$$L_{ph} = -\frac{1}{2} \sum_{\mathbf{k}} (\dot{q}_{\mathbf{k}}^2 + \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2). \tag{6.15}$$

The Lagrangian of the electron-phonon interaction (6.8) with the real phonon coordinates is transformed in the following way using (6.13):

$$\begin{aligned} L_{e-ph} = &- \sum_{\mathbf{k}} \left(\frac{2\omega_{\mathbf{k}}}{\hbar} \right)^{1/2} V_{\mathbf{k}} \rho_{\mathbf{k}} \left(Q'_{\mathbf{k}} - i Q''_{\mathbf{k}} \right) \\ = &- \sum_{\substack{\mathbf{k} \\ (k_x \geq 0)}} \left(\frac{2\omega_{\mathbf{k}}}{\hbar} \right)^{1/2} \left(V_{\mathbf{k}} \rho_{\mathbf{k}} + V_{-\mathbf{k}} \rho_{-\mathbf{k}} \right) Q'_{\mathbf{k}} \\ &+ i \sum_{\substack{\mathbf{k} \\ (k_x < 0)}} \left(\frac{2\omega_{\mathbf{k}}}{\hbar} \right)^{1/2} \left(V_{\mathbf{k}} \rho_{\mathbf{k}} - V_{-\mathbf{k}} \rho_{-\mathbf{k}} \right) Q''_{\mathbf{k}}. \end{aligned}$$

Let us introduce the real forces:

$$\gamma_{\mathbf{k}} \equiv \begin{cases} \frac{1}{\sqrt{2}} \left(\frac{2\omega_{\mathbf{k}}}{\hbar} \right)^{1/2} \left(V_{\mathbf{k}} \rho_{\mathbf{k}} + V_{-\mathbf{k}} \rho_{-\mathbf{k}} \right),\ k_x \geq 0, \\ \frac{1}{i\sqrt{2}} \left(\frac{2\omega_{\mathbf{k}}}{\hbar} \right)^{1/2} \left(V_{\mathbf{k}} \rho_{\mathbf{k}} - V_{-\mathbf{k}} \rho_{-\mathbf{k}} \right),\ k_x > 0. \end{cases} \tag{6.16}$$

This gives us the Lagrangian of the electron-phonon interaction in terms of the real forces and real phonon coordinates:

$$L_{e-ph} = -\sum_{\mathbf{k}} \gamma_{\mathbf{k}} q_{\mathbf{k}}. \tag{6.17}$$

So, the sum of the Lagrangians of phonons and of the electron-phonon interaction is expressed through ordinary real oscillator variables:

$$L_{ph} + L_{e-ph} = -\frac{1}{2} \sum_{\mathbf{k}} (\dot{q}_{\mathbf{k}}^2 + \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2 + \gamma_{\mathbf{k}} q_{\mathbf{k}}). \tag{6.18}$$

The path integration for each phonon mode with the coordinate $q_{\mathbf{k}}$ is performed independently as described in Sec. 2 of Ref. [41] and gives the result

$$\begin{aligned} &\int_{-\infty}^{\infty} dq_{\mathbf{k}} \int_{q_{\mathbf{k}}}^{q_{\mathbf{k}}} Dq_{\mathbf{k}}(\tau) \exp\left[-\frac{1}{\hbar}\int_0^{\hbar\beta} d\tau \frac{1}{2}(\dot{q}_{\mathbf{k}}^2 + \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2 + \gamma_{\mathbf{k}} q_{\mathbf{k}})\right] \\ &= \frac{1}{2\sinh\left(\frac{\beta\hbar\omega_{\mathbf{k}}}{2}\right)} \\ &\times \exp\left\{\frac{1}{4\hbar}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'| - \hbar\beta/2\right)\right]}{\omega_{\mathbf{k}} \sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)} \gamma_{\mathbf{k}}(\tau)\, \gamma_{\mathbf{k}}(\tau'),\right\} \end{aligned}$$

where the exponential is the influence functional of a driven oscillator {[41], Eq. (3.43)}. Therefore, the path integral over all phonon modes is

$$\begin{aligned} &\int d\{q_{\mathbf{k}}\} \int_{\{q_{\mathbf{k}}\}}^{\{q_{\mathbf{k}}\}} D\{q_{\mathbf{k}}(\tau)\} \exp\left[-\frac{1}{\hbar}\int_0^{\hbar\beta} d\tau \sum_{\mathbf{k}} \frac{1}{2}(\dot{q}_{\mathbf{k}}^2 + \omega_{\mathbf{k}}^2 q_{\mathbf{k}}^2 + \gamma_{\mathbf{k}} q_{\mathbf{k}})\right] \\ &= \left(\prod_{\mathbf{k}} \frac{1}{2\sinh\left(\frac{\beta\hbar\omega_{\mathbf{k}}}{2}\right)}\right) \\ &\times \exp\left\{\frac{1}{4\hbar}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \sum_{\mathbf{k}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'| - \hbar\beta/2\right)\right]}{\omega_{\mathbf{k}} \sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)} \gamma_{\mathbf{k}}(\tau)\, \gamma_{\mathbf{k}}(\tau').\right\} \end{aligned}$$

Here, the product $\prod_{\mathbf{k}} \ldots$ is the partition function of free phonons, and the exponential is the influence functional of the phonon subsystem on the electron subsystem. This influence functional results from the above described *elimination of the phonon coordinates* and is usually written down as $e^{-\Phi}$, where Φ is

$$\Phi = -\frac{1}{4\hbar}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \sum_{\mathbf{k}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'| - \hbar\beta/2\right)\right]}{\omega_{\mathbf{k}} \sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)} \gamma_{\mathbf{k}}(\tau)\, \gamma_{\mathbf{k}}(\tau'). \tag{6.19}$$

The sum over the phonon wave vectors $\mathbf{k}$ can be simplified as follows:

$$\sum_{\mathbf{k}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\omega_{\mathbf{k}}\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)}\gamma_{\mathbf{k}}\left(\tau\right)\gamma_{\mathbf{k}}\left(\tau'\right)$$
$$= \frac{1}{\hbar}\sum_{\substack{\mathbf{k}\\(k_x\geq 0)}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)}$$
$$\times\left[V_{\mathbf{k}}\rho_{\mathbf{k}}\left(\tau\right)+V_{-\mathbf{k}}\rho_{-\mathbf{k}}\left(\tau\right)\right]\left[V_{\mathbf{k}}\rho_{\mathbf{k}}\left(\tau'\right)+V_{-\mathbf{k}}\rho_{-\mathbf{k}}\left(\tau'\right)\right]$$
$$- \frac{1}{\hbar}\sum_{\substack{\mathbf{k}\\(k_x< 0)}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)}$$
$$\times\left[V_{\mathbf{k}}\rho_{\mathbf{k}}\left(\tau\right)-V_{-\mathbf{k}}\rho_{-\mathbf{k}}\left(\tau\right)\right]\left[V_{\mathbf{k}}\rho_{\mathbf{k}}\left(\tau'\right)-V_{-\mathbf{k}}\rho_{-\mathbf{k}}\left(\tau'\right)\right]$$
$$= \frac{2}{\hbar}\sum_{\substack{\mathbf{k}\\(k_x\geq 0)}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)}V_{\mathbf{k}}V_{-\mathbf{k}}$$
$$\times\left[\rho_{\mathbf{k}}\left(\tau\right)\rho_{-\mathbf{k}}\left(\tau'\right)+\rho_{-\mathbf{k}}\left(\tau\right)\rho_{\mathbf{k}}\left(\tau'\right)\right]$$
$$= \frac{2}{\hbar}\sum_{\mathbf{k}} \frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)}\left|V_{\mathbf{k}}\right|^2\rho_{\mathbf{k}}\left(\tau\right)\rho_{-\mathbf{k}}\left(\tau'\right).$$

Herefrom, we find that

$$\Phi = -\sum_{\mathbf{k}}\frac{\left|V_{\mathbf{k}}\right|^2}{2\hbar^2}\int_0^{\hbar\beta}d\tau\int_0^{\hbar\beta}d\tau'\frac{\cosh\left[\omega_{\mathbf{k}}\left(|\tau-\tau'|-\frac{\hbar\beta}{2}\right)\right]}{\sinh\left(\frac{\beta\hbar\omega_{\mathbf{k}}}{2}\right)}\rho_{\mathbf{k}}\left(\tau\right)\rho_{-\mathbf{k}}\left(\tau'\right). \tag{6.20}$$

As a result, the partition function of the electron-phonon system (6.11) factorizes into a product

$$Z\left(\{N_\sigma\},\beta\right) = Z_p\left(\{N_\sigma\},\beta\right)\prod_{\mathbf{k}}\frac{1}{2\sinh\left(\beta\hbar\omega_{\mathbf{k}}/2\right)} \tag{6.21}$$

of the partition function of free phonons with a partition function $Z_p\left(\{N_\sigma\},\beta\right)$ of interacting polarons, which is a path integral over the electron coordinates only:

$$Z_p\left(\{N_\sigma\},\beta\right) = \sum_P\frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}\left(\tau\right)e^{-S_p\left[\bar{\mathbf{x}}(\tau)\right]}. \tag{6.22}$$

The functional

$$S_p\left[\bar{\mathbf{x}}\left(\tau\right)\right] = -\frac{1}{\hbar}\int_0^{\hbar\beta}\left[L_e\left(\dot{\bar{\mathbf{x}}}\left(\tau\right),\bar{\mathbf{x}}\left(\tau\right)\right)-V_C\left(\bar{\mathbf{x}}\left(\tau\right)\right)\right]d\tau+\Phi\left[\bar{\mathbf{x}}\left(\tau\right)\right] \tag{6.23}$$

describes the phonon-induced retarded interaction between the electrons, including the retarded self-interaction of each electron.

Using (6.3) and (6.6) we write down $S_p\left[\bar{\mathbf{x}}\left(\tau\right)\right]$ explicitly:

$$S_p\left[\bar{\mathbf{x}}\left(\tau\right)\right]=\frac{1}{\hbar}\int_0^{\hbar\beta}\left[\sum_{\sigma}\sum_{j=1}^{N_\sigma}\frac{m_b}{2}\left(\dot{\mathbf{x}}_{j,\sigma}^2+\Omega_0^2\mathbf{x}_{j,\sigma}^2\right)+\sum_{\substack{\sigma,\sigma'\\(j,\sigma)\neq(l,\sigma')}}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_{\sigma'}}\frac{e^2}{2\varepsilon_\infty\left|\mathbf{x}_{j,\sigma}-\mathbf{x}_{l,\sigma'}\right|}\right]d\tau$$

$$-\sum_{\mathbf{q}}\frac{\left|V_{\mathbf{q}}\right|^2}{2\hbar^2}\int\limits_0^{\hbar\beta}d\tau\int\limits_0^{\hbar\beta}d\tau'\frac{\cosh\left[\omega_{\text{LO}}\left(\left|\tau-\tau'\right|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\text{LO}}/2\right)}\rho_{\mathbf{q}}\left(\tau\right)\rho_{-\mathbf{q}}\left(\tau'\right). \tag{6.24}$$

The free energy of a system of interacting polarons $F_p\left(\left\{N_\sigma\right\},\beta\right)$ is related to their partition function (6.22) by the equation:

$$F_p\left(\left\{N_\sigma\right\},\beta\right)=-\frac{1}{\beta}\ln Z_p\left(\left\{N_\sigma\right\},\beta\right). \tag{6.25}$$

At present no method is known to calculate the non-gaussian path integral (6.22) analytically. For *distinguishable* particles, the Jensen-Feynman variational principle [41] provides a convenient approximation technique. It yields a lower bound to the partition function, and hence an upper bound to the free energy.

It can be shown [97] that the path-integral approach to the many-body problem for a fixed number of identical particles can be formulated as a Feynman-Kac functional on a state space for N indistinguishable particles, by imposing an ordering on the configuration space and by the introduction of a set of boundary conditions at the boundaries of this state space. The resulting variational inequality for identical particles takes the same form as the Jensen-Feynman variational principle:

$$F_p\leqslant F_{var}, \tag{6.26}$$

$$F_{var}=F_0+\frac{1}{\beta}\left\langle S_p-S_0\right\rangle_{S_0}, \tag{6.27}$$

where S_0 is a model action with the corresponding free energy F_0. The angular brackets mean a weighted average over the paths

$$\left\langle\left(\bullet\right)\right\rangle_{S_0}=\frac{\sum_P\frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}\left(\tau\right)\left(\bullet\right)e^{-S_0\left[\bar{\mathbf{x}}(\tau)\right]}}{\sum_P\frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}\left(\tau\right)e^{-S_0\left[\bar{\mathbf{x}}(\tau)\right]}}. \tag{6.28}$$

In the zero-temperature limit, the polaron ground-state energy

$$E_p^0=\lim_{\beta\to\infty}F_p \tag{6.29}$$

obeys the inequality following from (6.26) with (6.27):

$$E_p^0 \leqslant E_{var}$$

with

$$E_{var} = E_0^0 + \lim_{\beta\to\infty}\left(\frac{1}{\beta}\left\langle S_p - S_0\right\rangle_{S_0}\right), \tag{6.30}$$

$$E_0^0 = \lim_{\beta\to\infty} F_0. \tag{6.31}$$

B. Model system

We consider a model system consisting of N electrons with coordinates $\bar{\mathbf{x}} \equiv \{\mathbf{x}_{j,\sigma}\}$ and N_f fictitious particles with coordinates $\bar{\mathbf{y}} \equiv \{\mathbf{y}_j\}$ in a harmonic confinement potential with elastic interparticle interactions as studied in Refs. [96, 98]. The Lagrangian of this model system takes the form

$$\begin{aligned} L_M\left(\dot{\bar{\mathbf{x}}}, \dot{\bar{\mathbf{y}}}; \bar{\mathbf{x}}, \bar{\mathbf{y}}\right) = &-\frac{m_b}{2}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\dot{\mathbf{x}}_{j,\sigma}^2 + \Omega^2\mathbf{x}_{j,\sigma}^2\right) + \frac{m_b\omega^2}{4}\sum_{\sigma,\sigma'}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_{\sigma'}}\left(\mathbf{x}_{j,\sigma} - \mathbf{x}_{l,\sigma'}\right)^2 \\ &- \frac{m_f}{2}\sum_{j=1}^{N_f}\left(\dot{\mathbf{y}}_j^2 + \Omega_f^2\mathbf{y}_j^2\right) - \frac{k}{2}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_f}\left(\mathbf{x}_{j,\sigma} - \mathbf{y}_l\right)^2. \end{aligned} \tag{6.32}$$

The frequencies Ω, ω, Ω_f, the mass of a fictitious particle m_f, and the force constant k are variational parameters. Clearly, this Lagrangian is symmetric with respect to electron permutations. Performing the path integral over the coordinates of the fictitious particles in the same way as described above for phonons, the partition function $Z_0\left(\{N_\sigma\}, \beta\right)$ of the model system of interacting polarons becomes a path integral over the electron coordinates:

$$Z_0\left(\{N_\sigma\}, \beta\right) = \sum_P \frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}} D\bar{\mathbf{x}}(\tau)\, e^{-S_0[\bar{\mathbf{x}}(\tau)]}, \tag{6.33}$$

with the action functional $S_0\left[\bar{\mathbf{x}}(\tau)\right]$ given by

$$\begin{aligned} S_0\left[\bar{\mathbf{x}}(\tau)\right] = &\frac{1}{\hbar}\int_0^{\hbar\beta}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\frac{m_b}{2}\left[\dot{\mathbf{x}}_{j,\sigma}^2(\tau) + \Omega^2\mathbf{x}_{j,\sigma}^2(\tau)\right]d\tau \\ &- \frac{1}{\hbar}\int_0^{\hbar\beta}\sum_{\sigma,\sigma'}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_{\sigma'}}\frac{m_b\omega^2}{4}\left[\mathbf{x}_{j,\sigma}(\tau) - \mathbf{x}_{l,\sigma'}(\tau)\right]^2 d\tau \\ &- \frac{k^2N^2N_f}{4m_f\hbar\Omega_f}\int_0^{\hbar\beta}d\tau\int_0^{\hbar\beta}d\tau'\frac{\cosh\left[\Omega_f\left(|\tau-\tau'| - \hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\Omega_f/2\right)}\mathbf{X}(\tau)\cdot\mathbf{X}(\tau'), \end{aligned} \tag{6.34}$$

where $\mathbf{X}$ is the center-of-mass coordinate of the electrons,

$$\mathbf{X} = \frac{1}{N} \sum_{\sigma} \sum_{j=1}^{N_\sigma} \mathbf{x}_{j,\sigma}. \tag{6.35}$$

1. *Analytical calculation of the model partition function*

The partition function $Z_0 \left(\{N_\sigma\}, \beta \right)$ [Eq. (6.33)] for the model system of interacting polarons can be expressed in terms of the partition function $Z_M \left(\{N_\sigma\}, N_f, \beta \right)$ of the model system of interacting electrons and fictitious particles with the Lagrangian L_M [Eq. (6.32)] as follows:

$$Z_0 \left(\{N_\sigma\}, \beta \right) = \frac{Z_M \left(\{N_\sigma\}, N_f, \beta \right)}{Z_f \left(N_f, w_f, \beta \right)}, \tag{6.36}$$

where $Z_f \left(N_f, w_f, \beta \right)$ is the partition function of fictitious particles,

$$Z_f \left(N_f, \beta \right) = \frac{1}{\left(2 \sinh \frac{1}{2} \beta \hbar w_f \right)^{D N_f}}, \tag{6.37}$$

with the frequency

$$w_f = \sqrt{\Omega_f^2 + kN/m_f} \tag{6.38}$$

and D=3(2) for 3D(2D) systems. The partition function $Z_M \left(\{N_\sigma\}, N_f, \beta \right)$ is the path integral for both the electrons and the fictitious particles:

$$Z_M \left(\{N_\sigma\}, N_f, \beta \right) = \sum_P \frac{(-1)^{\xi_P}}{N_{1/2}! N_{-1/2}!} \int d\bar{\mathbf{x}} \int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}} D\bar{\mathbf{x}}(\tau) \int d\bar{\mathbf{y}} \int_{\bar{\mathbf{y}}}^{\bar{\mathbf{y}}} D\bar{\mathbf{y}}(\tau)\, e^{-S_M[\bar{\mathbf{x}}(\tau), \bar{\mathbf{y}}(\tau)]} \tag{6.39}$$

with the "action" functional

$$S_M \left[\bar{\mathbf{x}}(\tau), \bar{\mathbf{y}}(\tau) \right] = -\frac{1}{\hbar} \int_0^{\hbar\beta} L_M \left(\dot{\bar{\mathbf{x}}}, \dot{\bar{\mathbf{y}}}; \bar{\mathbf{x}}, \bar{\mathbf{y}} \right) d\tau, \tag{6.40}$$

where the Lagrangian is given by Eq. (6.32).

Let us consider an auxiliary "ghost" subsystem with the Lagrangian

$$L_g \left(\dot{\mathbf{X}}_g, \dot{\mathbf{Y}}_g, \mathbf{X}_g, \mathbf{Y}_g \right) = -\frac{m_b N}{2} \left(\dot{\mathbf{X}}_g^2 + w^2 \mathbf{X}_g^2 \right) - \frac{m_f N_f}{2} \left(\dot{\mathbf{Y}}_g^2 + w_f^2 \mathbf{Y}_g^2 \right) \tag{6.41}$$

with two frequencies w and w_f, where w is given by

$$w = \sqrt{\Omega^2 - N\omega^2 + kN_f/m_b}. \tag{6.42}$$

The partition function Z_g of this subsystem

$$Z_g = \int d\mathbf{X}_g \int d\mathbf{Y}_g \int\limits_{\mathbf{X}_g}^{\mathbf{X}_g} D\mathbf{X}_g\left(\tau\right) \int\limits_{\mathbf{Y}_g}^{\mathbf{Y}_g} D\mathbf{Y}_g\left(\tau\right) \exp\left\{-S_g\left[\mathbf{X}_g\left(\tau\right), \mathbf{Y}_g\left(\tau\right)\right]\right\}, \tag{6.43}$$

with the "action" functional

$$S_g\left[\mathbf{X}_g\left(\tau\right), \mathbf{Y}_g\left(\tau\right)\right] = -\frac{1}{\hbar} \int\limits_0^{\hbar\beta} L_g\left(\dot{\mathbf{X}}_g, \mathbf{X}_g, \dot{\mathbf{Y}}_g, \mathbf{Y}_g\right) d\tau \tag{6.44}$$

is calculated in the standard way, because its Lagrangian (6.41) has a simple oscillator form. Consequently, the partition function Z_g is

$$Z_g = \frac{1}{\left[2\sinh\left(\frac{\beta\hbar w}{2}\right)\right]^D} \frac{1}{\left[2\sinh\left(\frac{\beta\hbar w_f}{2}\right)\right]^D}. \tag{6.45}$$

The product $Z_g Z_M$ of the two partition functions Z_g and $Z_M\left(\{N_\sigma\}, N_f, \beta\right)$ is a path integral in the state space of N electrons, N_f fictitious particles and two "ghost" particles with the coordinate vectors $\mathbf{X}_g$ and $\mathbf{Y}_g$. The Lagrangian $\tilde{L}_M$ of this system is a sum of L_M and L_g,

$$\tilde{L}_M\left(\dot{\bar{\mathbf{x}}}, \dot{\bar{\mathbf{y}}}, \dot{\mathbf{X}}_g, \dot{\mathbf{Y}}_g; \bar{\mathbf{x}}, \bar{\mathbf{y}}, \mathbf{X}_g, \mathbf{Y}_g\right) \equiv L_M\left(\dot{\bar{\mathbf{x}}}, \dot{\bar{\mathbf{y}}}; \bar{\mathbf{x}}, \bar{\mathbf{y}}\right) + L_g\left(\dot{\mathbf{X}}_g, \dot{\mathbf{Y}}_g, \mathbf{X}_g, \mathbf{Y}_g\right). \tag{6.46}$$

The "ghost" subsystem is introduced because the center-of-mass coordinates in $\tilde{L}_M$ can be explicitly separated much more transparently than in L_M. This separation is realized by the linear transformation of coordinates,

$$\begin{cases} \mathbf{x}_{j,\sigma} = \mathbf{x}'_{j,\sigma} + \mathbf{X} - \mathbf{X}_g, \\ \mathbf{y}_{j\sigma} = \mathbf{y}'_{j\sigma} + \mathbf{Y} - \mathbf{Y}_g, \end{cases} \tag{6.47}$$

where $\mathbf{X}$ and $\mathbf{Y}$ are the center-of-mass coordinate vectors of the electrons and of the fictitious particles, correspondingly:

$$\mathbf{X} = \frac{1}{N} \sum_\sigma \sum_{j=1}^{N_\sigma} \mathbf{x}_{j,\sigma}, \quad \mathbf{Y} = \frac{1}{N_f} \sum_{j=1}^{N_f} \mathbf{y}_j. \tag{6.48}$$

Before the transformation (6.47), the independent variables are $\left(\bar{\mathbf{x}}, \bar{\mathbf{y}}, \mathbf{X}_g, \mathbf{Y}_g\right)$, with the center-of-mass coordinates $\mathbf{X}$ and $\mathbf{Y}$ determined by Eq. (6.48). When applying the transformation (6.47) to the centers of mass (6.48), we find that

$$\mathbf{X} = \frac{1}{N}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\mathbf{x}'_{j,\sigma} + \mathbf{X} - \mathbf{X}_g\right) = \frac{1}{N}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\mathbf{x}'_{j,\sigma} + \mathbf{X} - \mathbf{X}_g, \tag{6.49}$$

$$\mathbf{Y}=\frac{1}{N_f}\sum_{j=1}^{N_f}\left(\mathbf{y}'_{j\sigma} + \mathbf{Y} - \mathbf{Y}_g\right) = \frac{1}{N_f}\sum_{j=1}^{N_f}\mathbf{y}'_{j\sigma} + \mathbf{Y} - \mathbf{Y}_g. \tag{6.50}$$

As seen from Eqs. (6.49), (6.50), after the transformation (6.47) the independent variables are $(\bar{\mathbf{x}}', \bar{\mathbf{y}}', \mathbf{X}, \mathbf{Y})$, while the coordinates $(\mathbf{X}_g, \mathbf{Y}_g)$ obey the equations

$$\mathbf{X}_g=\frac{1}{N}\sum_{\sigma}\sum_{j=1}^{N_\sigma}\mathbf{x}'_{j,\sigma}, \quad \mathbf{Y}_g=\frac{1}{N_f}\sum_{j=1}^{N_f}\mathbf{y}'_j. \tag{6.51}$$

In order to find the explicit form of the Lagrangian (6.46) after the transformation (6.47), we use the following relations for the quadratic sums of coordinates:

$$\begin{gathered}
\sum_{\sigma}\sum_{j=1}^{N_\sigma}\mathbf{x}^2_{j,\sigma} = \sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\mathbf{x}'_{j,\sigma}\right)^2 + N\left(\mathbf{X}^2 - \mathbf{X}_g^2\right), \quad \sum_{j=1}^{N_f}\mathbf{y}_j^2 = \sum_{j=1}^{N_f}\left(\mathbf{y}'_j\right)^2 + N_f\left(\mathbf{Y}^2 - \mathbf{Y}_g^2\right), \\
\sum_{\sigma,\sigma'}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_{\sigma'}}\left(\mathbf{x}_{j,\sigma} - \mathbf{x}_{l,\sigma'}\right)^2 = 2N\sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\mathbf{x}'_{j,\sigma}\right)^2 - 2N_f^2\mathbf{X}_g^2, \\
\sum_{j=1}^{N_f}\sum_{l=1}^{N_f}\left(\mathbf{y}_j - \mathbf{y}_l\right)^2 = 2N_f\sum_{j=1}^{N_f}\left(\mathbf{y}'_j\right)^2 - 2N_f^2\mathbf{Y}_g^2, \\
\sum_{\sigma}\sum_{j=1}^{N_\sigma}\sum_{l=1}^{N_f}\left(\mathbf{x}_{j,\sigma} - \mathbf{y}_l\right)^2 = N_f\sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\mathbf{x}'_{j,\sigma}\right)^2 + N\sum_{j=1}^{N_f}\left(\mathbf{y}'_j\right)^2 + \\
NN_f\left(\mathbf{X}^2 + \mathbf{Y}^2 - 2\mathbf{X}\cdot\mathbf{Y} - \mathbf{X}_g^2 - \mathbf{Y}_g^2\right).
\end{gathered} \tag{6.52}$$

The substitution of Eq. (6.48) into Eq. (6.46) then results in the following 3 terms:

$$\tilde{L}_M\left(\dot{\bar{\mathbf{x}}}', \dot{\bar{\mathbf{y}}}', \dot{\mathbf{X}}, \dot{\mathbf{Y}}; \bar{\mathbf{x}}', \bar{\mathbf{y}}', \mathbf{X}, \mathbf{Y}\right) = L_w\left(\dot{\bar{\mathbf{x}}}', \bar{\mathbf{x}}'\right) + L_{w_f}\left(\dot{\bar{\mathbf{y}}}', \bar{\mathbf{y}}'\right) + L_C\left(\dot{\mathbf{X}}, \mathbf{X}; \dot{\mathbf{Y}}, \mathbf{Y}\right), \tag{6.53}$$

where $L_w\left(\dot{\bar{\mathbf{x}}}', \bar{\mathbf{x}}'\right)$ and $L_{w_f}\left(\dot{\bar{\mathbf{y}}}', \bar{\mathbf{y}}'\right)$ are Lagrangians of non-interacting identical oscillators with the frequencies w and w_f, respectively,

$$L_w\left(\dot{\bar{\mathbf{x}}}', \bar{\mathbf{x}}'\right) = -\frac{m_b}{2}\sum_{\sigma=\pm 1/2}\sum_{j=1}^{N_\sigma}\left[\left(\dot{\mathbf{x}}'_{j,\sigma}\right)^2 + w^2\left(\mathbf{x}'_{j,\sigma}\right)^2\right], \tag{6.54}$$

$$L_{w_f}\left(\dot{\bar{\mathbf{y}}}', \bar{\mathbf{y}}'\right) = -\frac{m_f}{2}\sum_{j=1}^{N_f}\left[\left(\dot{\mathbf{y}}'_{j,\sigma}\right)^2 + w_f^2\left(\mathbf{y}'_{j,\sigma}\right)^2\right]. \tag{6.55}$$

The Lagrangian $L_C\left(\dot{\mathbf{X}},\mathbf{X};\dot{\mathbf{Y}},\mathbf{Y}\right)$ describes the combined motion of the centers-of-mass of the electrons and of the fictitious particles,

$$L_C\left(\dot{\mathbf{X}},\mathbf{X};\dot{\mathbf{Y}},\mathbf{Y}\right)=-\frac{m_bN}{2}\left(\dot{\mathbf{X}}^2+\tilde{\Omega}^2\mathbf{X}^2\right)-\frac{m_fN_f}{2}\left(\dot{\mathbf{Y}}^2+w_f^2\mathbf{Y}^2\right)+kNN_f\mathbf{X}\cdot\mathbf{Y}, \quad (6.56)$$

with

$$\tilde{\Omega}=\sqrt{\Omega^2+kN_f/m_b}. \quad (6.57)$$

The Lagrangian (6.56) is reduced to a diagonal quadratic form in the coordinates and the velocities by a unitary transformation for two interacting oscillators using the following replacement of variables:

$$\begin{aligned}\mathbf{X}&=\frac{1}{\sqrt{m_bN}}\left(a_1\mathbf{r}+a_2\mathbf{R}\right),\\ \mathbf{Y}&=\frac{1}{\sqrt{m_fN_f}}\left(-a_2\mathbf{r}+a_1\mathbf{R}\right)\end{aligned} \quad (6.58)$$

with the coefficients

$$a_1=\left[\frac{1+\chi}{2}\right]^{1/2},\quad a_2=\left[\frac{1-\chi}{2}\right]^{1/2}, \quad (6.59)$$

$$\chi\equiv\frac{\tilde{\Omega}^2-\tilde{\Omega}_f^2}{\left[\left(\tilde{\Omega}^2-\tilde{\Omega}_f^2\right)^2+4\gamma^2\right]^{1/2}},\quad \gamma\equiv k\sqrt{\frac{NN_f}{m_bm_f}}. \quad (6.60)$$

The eigenfrequencies of the center-of-mass subsystem are then given by the expression

$$\begin{cases}\Omega_1=\sqrt{\frac{1}{2}\left[\tilde{\Omega}^2+\tilde{\Omega}_f^2+\sqrt{\left(\tilde{\Omega}^2-\tilde{\Omega}_f^2\right)^2+4\gamma^2}\right]},\\ \Omega_2=\sqrt{\frac{1}{2}\left[\tilde{\Omega}^2+\tilde{\Omega}_f^2-\sqrt{\left(\tilde{\Omega}^2-\tilde{\Omega}_f^2\right)^2+4\gamma^2}\right]}.\end{cases} \quad (6.61)$$

As a result, four independent frequencies Ω_1, Ω_2, w and w_f appear in the problem. Three of them (Ω_1, Ω_2, w) are the eigenfrequencies of the model system. Ω_1 is the frequency of the relative motion of the center of mass of the electrons with respect to the center of mass of the fictitious particles; Ω_2 is the frequency related to the center of mass of the model system as a whole; w is the frequency of the relative motion of the electrons with respect to their center of mass. The parameter w_f is an analog of the second variational parameter w of the one-polaron Feynman model. Further, the Lagrangian (6.56) takes the form

$$L_C=-\frac{1}{2}\left(\dot{\mathbf{r}}^2+\Omega_1^2\mathbf{r}^2\right)-\frac{1}{2}\left(\dot{\mathbf{R}}^2+\Omega_2^2\mathbf{R}^2\right), \quad (6.62)$$

leading to the partition function corresponding to the combined motion of the centers-of-mass of the electrons and of the fictitious particles

$$Z_C = \frac{1}{\left[2\sinh\left(\frac{\beta\hbar\Omega_1}{2}\right)\right]^D} \frac{1}{\left[2\sinh\left(\frac{\beta\hbar\Omega_2}{2}\right)\right]^D}. \tag{6.63}$$

Taking into account Eqs. (6.45) and (6.63), we obtain finally the partition function of the model system for interacting polarons

$$Z_0\left(\{N_\sigma\},\beta\right) = \left[\frac{\sinh\left(\frac{\beta\hbar w}{2}\right)\sinh\left(\frac{\beta\hbar w_f}{2}\right)}{\sinh\left(\frac{\beta\hbar\Omega_1}{2}\right)\sinh\left(\frac{\beta\hbar\Omega_2}{2}\right)}\right]^D \tilde{\mathbb{Z}}_F\left(\{N_\sigma\},w,\beta\right). \tag{6.64}$$

Here

$$\tilde{\mathbb{Z}}_F\left(\{N_\sigma\},w,\beta\right) = \mathbb{Z}_F\left(N_{1/2},w,\beta\right)\mathbb{Z}_F\left(N_{-1/2},w,\beta\right) \tag{6.65}$$

is the partition function of $N = N_{1/2}+N_{-1/2}$ non-interacting fermions in a parabolic confinement potential with the frequency w. The analytical expressions for the partition function of N_σ spin-polarized fermions $\mathbb{Z}_F\left(N_\sigma,w,\beta\right)$ were derived in Ref. [99].

C. Variational functional

In order to obtain an upper bound to the free energy E_{var}, we substitute the model action functional (6.34) into the right-hand side of the variational inequality (6.26) and consider the limit $\beta \to \infty$:

$$\begin{aligned}
&E_{var}\left(\{N_\sigma\}\right) \\
&= \mathbb{E}_F\left(\{N_\sigma\},w\right) + \frac{m_b}{2}\left(\Omega_0^2 - \Omega^2 + N\omega^2\right)\left\langle \sum_{j=1}^{N} \mathbf{x}_j^2(0)\right\rangle_{S_0} \\
&- \frac{m_b\omega^2N^2}{2}\left\langle \mathbf{X}^2(0)\right\rangle_{S_0} + \left\langle U_b(\bar{\mathbf{x}})\right\rangle_{S_0} + \sum_{\mathbf{q}\neq 0}\frac{2\pi e^2}{V\varepsilon_\infty q^2}\left[\mathcal{G}\left(\mathbf{q},0|\{N_\sigma\},\beta\to\infty\right) - N\right] \\
&+ \lim_{\beta\to\infty}\frac{k^2N^2N_f}{4m_f\beta\hbar\Omega_f}\int_0^{\hbar\beta}d\tau\int_0^{\hbar\beta}d\tau'\frac{\cosh\left[\Omega_f\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\Omega_f/2\right)}\left\langle \mathbf{X}(\tau)\cdot\mathbf{X}(\tau')\right\rangle_{S_0} \\
&- \lim_{\beta\to\infty}\sum_{\mathbf{q}}\frac{|V_{\mathbf{q}}|^2}{2\hbar^2\beta}\int_0^{\hbar\beta}d\tau\int_0^{\hbar\beta}d\tau'\frac{\cosh\left[\omega_{\mathrm{LO}}\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\omega_{\mathrm{LO}}/2\right)}\mathcal{G}\left(\mathbf{q},\tau-\tau'|\{N_\sigma\},\beta\right).
\end{aligned} \tag{6.66}$$

Here, $\mathbb{E}_F(N, w)$ is the energy of N non-interacting fermions in a parabolic confinement potential with the confinement frequency w,

$$\mathbb{E}_F(\{N_\sigma\}, w) = \hbar w \sum_{\sigma=\pm 1/2} \left\{ \sum_{n=0}^{L_\sigma - 1} \left(n + \frac{3}{2} \right) g(n) \right. \\ \left. + (N_\sigma - N_{L_\sigma}) \left(L_\sigma + \frac{3}{2} \right) \right\}, \tag{6.67}$$

where σ is the spin of an electron, L_σ is the lower partly filled or empty level for N_σ electrons with the spin projection σ. The first term in the curly brackets of Eq. (6.94) (the upper line) is the number of electrons at fully filled energy levels, while the second term (square brackets) is the number of electrons at the next upper level (which can be empty or filled partially). The energy levels of a 3D oscillator are degenerate, so that

$$g(n) = \frac{1}{2}(n+1)(n+2) \tag{6.68}$$

is the degeneracy of the n-th energy level. The parameter

$$N_{L_\sigma} = \frac{1}{6} L_\sigma (L_\sigma + 1)(L_\sigma + 2) \tag{6.69}$$

is the number of electrons at all fully filled levels. The summation in Eq. (6.67) is performed explicitly, what gives us the result

$$\mathbb{E}_F(\{N_\sigma\}, w) = \hbar w \sum_\sigma \left[\frac{1}{8} L_\sigma (L_\sigma + 1)^2 (L_\sigma + 2) + (N_\sigma - N_{L_\sigma}) \left(L_\sigma + \frac{3}{2} \right) \right]. \tag{6.70}$$

In Eq. (6.66), $\mathcal{G}(\mathbf{q}, \tau - \tau' | \{N_\sigma\}, \beta)$ is the two-point correlation function for the electron density operators:

$$\mathcal{G}(\mathbf{q}, \tau | \{N_\sigma\}, \beta) = \langle \rho_{\mathbf{q}}(\tau) \rho_{-\mathbf{q}}(0) \rangle_{S_0}. \tag{6.71}$$

The averages $\langle \mathbf{X}(\tau) \cdot \mathbf{X}(\tau') \rangle_{S_0}$ are calculated using the generating function method:

$$\langle X_k(\tau) X_k(\tau') \rangle_{S_0} = -\frac{\partial^2}{\partial \xi_k \partial \eta_k} \langle \exp[i(\xi \cdot \mathbf{X}(\tau) + \eta \cdot \mathbf{X}(\tau'))] \rangle_{S_0} \Big|_{\substack{\xi=0, \\ \eta=0}}, \tag{6.72}$$

$$\Longrightarrow \langle \mathbf{X}(\tau) \cdot \mathbf{X}(\tau') \rangle_{S_0} = \frac{3\hbar}{2mN} \sum_{i=1}^{2} \frac{a_i^2 \cosh[\Omega_i(|\tau - \sigma| - \hbar\beta/2)]}{\Omega_i \sinh(\hbar\beta\Omega_i/2)}. \tag{6.73}$$

Substituting this expression into Eq. (6.66) and performing integrations over τ and σ analytically, we obtain the result

$$\frac{k^2 N^2 N_f}{4 m_f \beta\hbar\Omega_f}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \frac{\cosh\left[\Omega_f\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\Omega_f/2\right)}\left\langle \mathbf{X}(\tau)\cdot\mathbf{X}(\tau')\right\rangle_{S_0}$$
$$=\frac{3\hbar\gamma}{4}\sum_{i=1}^{2}\frac{a_i^2}{\Omega_f^2-\Omega_i^2}\left[\frac{\coth\left(\beta\Omega_i/2\right)}{\Omega_i}-\frac{\coth\left(\beta\Omega_f/2\right)}{\Omega_f}\right], \tag{6.74}$$

and in the zero-temperature limit we have

$$\lim_{\beta\to\infty}\frac{k^2 N^2 N_f}{4 m_f \beta\hbar\Omega_f}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\tau' \frac{\cosh\left[\Omega_f\left(|\tau-\tau'|-\hbar\beta/2\right)\right]}{\sinh\left(\beta\hbar\Omega_f/2\right)}\left\langle \mathbf{X}(\tau)\cdot\mathbf{X}(\tau')\right\rangle_{S_0}$$
$$=\frac{3\hbar\gamma}{4}\sum_{i=1}^{2}\frac{a_i^2}{\Omega_f^2-\Omega_i^2}\left(\frac{1}{\Omega_i}-\frac{1}{\Omega_f}\right). \tag{6.75}$$

The average $\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_0}$ is transformed, using the described above operations with the "ghost" subsystem,

$$\mathbf{x}_j=\mathbf{x}'_j+\mathbf{X}-\mathbf{X}_g, \tag{6.76}$$

and taking into account the first of equations (6.52)

$$\sum_{j=1}^{N}\mathbf{x}_j^2=\sum_{j=1}^{N}\left(\mathbf{x}'_j\right)^2+N\left(\mathbf{X}^2-\mathbf{X}_g^2\right). \tag{6.77}$$

Consequently, averaging the left-hand side of Eq. (6.77) on the model action functional S_0, one obtains

$$\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_0}=\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_M}=\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_M+S_g}$$
$$=\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_w}+N\left(\left\langle\mathbf{X}^2\right\rangle_{S_C}-\left\langle\mathbf{X}_g^2\right\rangle_{S_g}\right). \tag{6.78}$$

The term $\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_w}$ is expressed using the virial theorem through the ground-state energy $\mathbb{E}_F(N,w)$ of N independent 3D fermion oscillators with the frequency w and with the mass m_b,

$$\left\langle \sum_{j=1}^{N}\mathbf{x}_j^2\right\rangle_{S_w}=\frac{\mathbb{E}_F(N,w)}{m_b w^2}=-\frac{1}{m_b w^2}\frac{\partial}{\partial\lambda}\ln\mathbb{Z}_I(N), \tag{6.79}$$

Two other terms in Eq. (6.78) are [cf. Eq. (6.73)]:

$$\left\langle \mathbf{X}^2 \right\rangle_{S_C} = \frac{3\hbar}{2m_bN} \sum_{i=1}^{2} \frac{a_i^2 \coth\left(\beta\Omega_i/2\right)}{\Omega_i},$$
$$\left\langle \mathbf{X}_g^2 \right\rangle_{S_g} = \frac{3\hbar}{2m_bN} \frac{\coth\left(\beta w/2\right)}{w}. \tag{6.80}$$

So, we obtain

$$\left\langle \sum_{j=1}^{N} \mathbf{x}_j^2 \right\rangle_{S_0} = \frac{\mathbb{E}_F\left(\left\{N_\sigma\right\}, w\right)}{m_b w^2} + \frac{3\hbar}{2m_b} \left(\sum_{i=1}^{2} \frac{a_i^2}{\Omega_i} - \frac{1}{w} \right). \tag{6.81}$$

The averaging of the background-charge potential gives us the result

$$\left\langle U_b\left(\bar{\mathbf{x}}\right)\right\rangle_{S_0} = \frac{3\sqrt{2}\alpha\eta}{\pi\left(1-\eta\right)} \sum_{\sigma} \sum_{n=0}^{\infty} \left. f_1\left(n,\sigma|\beta,N_\sigma\right)\right|_{\beta\to\infty} \sum_{k=0}^{n} \frac{(-1)^k}{k!} \binom{n+2}{n-k} \left(\frac{1}{2w}\right)^k$$
$$\times \left\{ \frac{\Gamma\left(k-\frac{1}{2}\right)}{A^{k-1/2}} \left[{}_1F_1\left(k-\frac{1}{2};\frac{1}{2};-\frac{R^2}{4A}\right) - {}_1F_1\left(k-\frac{1}{2};\frac{3}{2};-\frac{R^2}{4A}\right) \right] \right\}, \tag{6.82}$$
$$\eta \equiv \varepsilon_\infty/\varepsilon_0, \quad A \equiv \frac{\hbar}{4m_bN} \left(\sum_{i=1}^{2} \frac{a_i^2}{\Omega_i} + \frac{N-1}{w} \right),$$

where $f_1\left(n,\sigma|\beta,N_\sigma\right)$ is the one-particle distribution function of fermions (the distribution functions are considered in more details in the next subsection).

Collecting all terms together, we arrive at the variational functional

$$E_{var}\left(\Omega_1,\Omega_2,w,\Omega_f\right) = \hbar \left\{ \frac{\Omega_0^2+w^2}{2w^2} \left[\frac{\tilde{E}\left(w,N\right)}{\hbar} - \frac{3}{2}w \right] + \frac{3}{2}\left(\Omega_1+\Omega_2-\Omega_f\right) \right.$$
$$\left. + \frac{3}{4}\left(\Omega_0^2-\Omega_1^2-\Omega_2^2+\Omega_f^2\right) \sum_{i=1}^{2} \frac{a_i^2}{\Omega_i} + \frac{3\gamma^2}{4\Omega_f} \sum_{i=1}^{2} \frac{a_i^2}{\Omega_i\left(\Omega_i+\Omega_f\right)} \right\}$$
$$+ \left\langle U_b\left(\bar{\mathbf{x}}\right)\right\rangle_{S_0} + E_C + E_{e-ph}, \tag{6.83}$$

where E_C and E_{e-ph} are the Coulomb and polaron contributions, respectively:

$$E_C = \frac{e^2}{4\pi^2\varepsilon_\infty} \int d\mathbf{q} \frac{1}{q^2} \left[\left. \mathcal{G}\left(\mathbf{q},0|\left\{N_\sigma\right\},\beta\right)\right|_{\beta\to\infty} - N \right], \tag{6.84}$$

$$E_{e-ph} = -\frac{\sqrt{2}\alpha}{4\pi^2\hbar} \int d\mathbf{q} \frac{1}{q^2} \int_0^\infty d\tau \exp\left(-\omega_{\mathrm{LO}}\tau\right) \left. \mathcal{G}\left(\mathbf{q},\tau|\left\{N_\sigma\right\},\beta\right)\right|_{\beta\to\infty}. \tag{6.85}$$

The correlation function (6.71) is calculated analytically in the next subsection. With this correlation function, the variational ground-state energy is calculated and minimized numerically.

D. Two-point correlation functions

The two-point correlation function (6.71) is represented as the following path integral:

$$\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)=\frac{1}{Z_0(\{N_\sigma\},\beta)}\sum_P\frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\times\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}(\tau)\,e^{-S_0[\bar{\mathbf{x}}(\tau)]}\rho_{\mathbf{q}}(\tau)\,\rho_{-\mathbf{q}}(0)\,. \tag{6.86}$$

We observe that $\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)$ can be rewritten as an average within the model "action" $S_M[\bar{\mathbf{x}}(\tau),\bar{\mathbf{y}}(\tau)]$ of interacting electrons and fictitious particles:

$$\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)=\frac{1}{Z_M(\{N_\sigma\},N_f,\beta)}\sum_P\frac{(-1)^{\xi_P}}{N_{1/2}!N_{-1/2}!}\times\int d\bar{\mathbf{x}}\int_{\bar{\mathbf{x}}}^{P\bar{\mathbf{x}}}D\bar{\mathbf{x}}(\tau)\int d\bar{\mathbf{y}}\int_{\bar{\mathbf{y}}}^{\bar{\mathbf{y}}}D\bar{\mathbf{y}}(\tau)\,e^{-S_M[\bar{\mathbf{x}}(\tau),\bar{\mathbf{y}}(\tau)]}\times\rho_{\mathbf{q}}(\tau)\,\rho_{-\mathbf{q}}(0)\,. \tag{6.87}$$

Indeed, one readily derives that the elimination of the fictitious particles in (6.87) leads to (6.86). The representation (6.87) allows one to calculate the correlation function $\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)$ in a much simpler way than through Eq. (6.86), using the separation of the coordinates of the centers of mass of the electrons and of the fictitious particles. This separation is performed for the two-point correlation function (6.87) by the same method as it has been done for the partition function (6.39). As a result, one obtains

$$\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)=\tilde{g}(\mathbf{q},\tau|\{N_\sigma\},\beta)\frac{\langle\exp[i\mathbf{q}\cdot(\mathbf{X}(\tau)-\mathbf{X}(\sigma))]\rangle_{S_C}}{\langle\exp[i\mathbf{q}\cdot(\mathbf{X}_g(\tau)-\mathbf{X}_g(\sigma))]\rangle_{S_g}}, \tag{6.88}$$

where $\tilde{g}(\mathbf{q},\tau|\{N_\sigma\},\beta)$ is the time-dependent correlation function of N non-interacting electrons in a parabolic confinement potential with the frequency w,

$$\tilde{g}(\mathbf{q},\tau|\{N_\sigma\},\beta)=\langle\rho_{\mathbf{q}}(\tau)\,\rho_{-\mathbf{q}}(0)\rangle_{S_w}\,. \tag{6.89}$$

The action functional $S_w[\bar{\mathbf{x}}_\tau]$ is related to the Lagrangian $L_w\left(\dot{\bar{\mathbf{x}}},\bar{\mathbf{x}}\right)$ [Eq. (6.54)]

$$S_w[\bar{\mathbf{x}}_\tau]=\frac{1}{\hbar}\int_0^{\hbar\beta}L_w\left(\dot{\bar{\mathbf{x}}},\bar{\mathbf{x}}\right)\,d\tau. \tag{6.90}$$

The averages in (6.88) are calculated using Feynman's method of generating functions [41]. Namely, according to [41], the average

$$G[f(\tau)]\equiv\left\langle\exp\left\{\frac{i}{\hbar}\int_0^{\beta}f(\tau)\,x_\tau\,d\tau\right\}\right\rangle_{S_\omega}, \tag{6.91}$$

where S_ω is the action functional of a one-dimensional harmonic oscillator with the frequency ω and with the mass m, results in

$$G\left[f\left(\tau\right)\right]=\exp\left\{-\frac{1}{4m\hbar\omega}\int\limits_0^\beta d\tau\int\limits_0^\beta d\sigma\frac{\cosh\left[\omega\left(\left|\tau-\sigma\right|-\beta/2\right)\right]}{\sinh\left(\beta\omega/2\right)}f\left(\tau\right)f\left(\sigma\right)\right\}. \tag{6.92}$$

The diagonalization procedure for the Lagrangian L_C (6.56) allows us to represent that Lagrangian as a sum of Lagrangians of independent harmonic oscillators, what gives the following explicit expressions for averages in Eq. (6.88):

$$\begin{aligned}&\left\langle \exp\left[i\mathbf{q}\cdot\left(\mathbf{X}\left(\tau\right)-\mathbf{X}\left(\sigma\right)\right)\right]\right\rangle_{S_C}\\&=\exp\left\{-\frac{\hbar q^2}{Nm_b}\left[\sum_{i=1}^{2}a_i^2\frac{\sinh\left(\frac{\Omega_i\left|\tau-\sigma\right|}{2}\right)\sinh\left(\frac{\Omega_i\left(\hbar\beta-\left|\tau-\sigma\right|\right)}{2}\right)}{\Omega_i\sinh\left(\frac{\beta\hbar\Omega_i}{2}\right)}\right]\right\},\\&\left\langle \exp\left[i\mathbf{q}\cdot\left(\mathbf{X}_g\left(\tau\right)-\mathbf{X}_g\left(\sigma\right)\right)\right]\right\rangle_{S_g}\\&=\exp\left[-\frac{\hbar q^2}{Nm_b}\frac{\sinh\left(\frac{w\left|\tau-\sigma\right|}{2}\right)\sinh\left(\frac{w\left(\hbar\beta-\left|\tau-\sigma\right|\right)}{2}\right)}{w\sinh\left(\frac{\beta\hbar w}{2}\right)}\right].\end{aligned}$$

1. The correlation function $\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta)$

As seen from the formula (6.89), $\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta)$ is the time-dependent correlation function of N non-interacting fermions in a parabolic confinement potential with the frequency w. Let us consider first of all a system of N identical spin-polarized oscillators with the Lagrangian

$$L=\frac{m}{2}\sum_{j=1}^{N}\left(\dot{\mathbf{x}}_j^2-\omega^2\mathbf{x}_j^2\right). \tag{6.93}$$

The corresponding Hamiltonian is

$$\hat{H}=\sum_{j=1}^{N}\left(\frac{\hat{\mathbf{p}}_j^2}{2m}+\frac{m\omega^2\hat{\mathbf{x}}_j^2}{2}\right), \tag{6.94}$$

$$\hat{H}=\sum_{j=1}^{N}\hat{h}_j,\quad \hat{h}\equiv\frac{\hat{\mathbf{p}}^2}{2m}+\frac{m\omega^2\hat{\mathbf{x}}^2}{2}. \tag{6.95}$$

A set of eigenfunctions of the one-particle Hamiltonian $\hat{h}$ is determined as follows:

$$\hat{h}\psi_{\mathbf{n}}\left(\mathbf{x}\right)=\varepsilon_n\psi_{\mathbf{n}}\left(\mathbf{x}\right), \tag{6.96}$$

where

$$\mathbf{n} \equiv (n_1, n_2, n_3)\,, \quad n \equiv n_1 + n_2 + n_3,$$
$$\varepsilon_{\mathbf{n}} = \varepsilon_n = \hbar\omega\left(n + \frac{3}{2}\right), \quad \psi_{\mathbf{n}}(\mathbf{x}) = \varphi_{n_1}(x_1)\,\varphi_{n_2}(x_2)\,\varphi_{n_3}(x_3)\,, \tag{6.97}$$

$\varphi_n(x)$ is the n-th eigenfunction of a one-dimensional oscillator with the frequency ω.

The Hamiltonian (6.94) can be written down in terms of the annihilation ($\hat{a}_{\mathbf{n}}$) and creation ($\hat{a}^+_{\mathbf{n}}$) operators:

$$\hat{H} = \sum_{\mathbf{n}} \varepsilon_{\mathbf{n}} \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}} = \sum_{\mathbf{n}} \varepsilon_{\mathbf{n}} \hat{N}_{\mathbf{n}}, \quad \hat{N}_{\mathbf{n}} \equiv \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}}. \tag{6.98}$$

The many-particle quantum states in the representation of "occupation numbers" are written down as $|\ldots N_{\mathbf{n}} \ldots\rangle$, where $N_{\mathbf{n}}$ is the number of particles in the $\mathbf{n}$-th one-particle quantum state. The states $|\ldots N_{\mathbf{n}} \ldots\rangle$ are defined as the eigenstates of the operator of the number of particles in the $\mathbf{n}$-th state $\hat{N}_{\mathbf{n}}$:

$$\hat{N}_{\mathbf{n}} |\ldots N_{\mathbf{n}} \ldots\rangle = N_{\mathbf{n}} |\ldots N_{\mathbf{n}} \ldots\rangle\,. \tag{6.99}$$

Let us determine a set of quantum states with a *finite* total number of particles

$$\sum_{\mathbf{n}} N_{\mathbf{n}} = N \tag{6.100}$$

as follows:

$$|\ldots N_{\mathbf{n}} \ldots\rangle|_{\sum_{\mathbf{n}} N_{\mathbf{n}} = N} \equiv \left|\Psi_{N,\{N_{\mathbf{n}}\}}\right\rangle. \tag{6.101}$$

Further on, we use the basis set of quantum states (6.101) for the derivation of the partition function, of the density function and of the two-point correlation function.

Partition function

The density matrix of the *canonical* Hibbs ensemble is

$$\hat{\rho} = \exp\left(-\beta\hat{H}\right), \quad \beta \equiv \frac{1}{k_B T}.$$

The partition function of this ensemble is the trace of the density matrix on the set of quantum states (6.101):

$$\begin{aligned}\mathbb{Z}_I(\beta|N) &= \sum_{\{N_{\mathbf{n}}\}} \left\langle \Psi_{N,\{N_{\mathbf{n}}\}} \left| \exp\left(-\beta\hat{H}\right) \right| \Psi_{N,\{N_{\mathbf{n}}\}} \right\rangle \\ &= \left[\sum_{\{N_{\mathbf{n}}\}} \exp\left(-\beta \sum_{\mathbf{n}} \varepsilon_{\mathbf{n}} N_{\mathbf{n}}\right)\right]_{\sum_{\mathbf{n}} N_{\mathbf{n}} = N}.\end{aligned} \tag{6.102}$$

This expression can be written down also in the form

$$\mathbb{Z}_I(\beta|N) = \sum_{\{N_\mathbf{n}\}} \exp\left(-\beta \sum_\mathbf{n} \varepsilon_\mathbf{n} N_\mathbf{n}\right) \delta_{N,\sum_\mathbf{n} N_\mathbf{n}}, \tag{6.103}$$

where

$$\delta_{j,k} = \begin{cases} 1, & j = k \\ 0, & j \neq k \end{cases}$$

is the delta symbol.

Let us introduce the generating function for the partition function in the same way as in Ref. [99]:

$$\begin{aligned} \Xi(\beta, u) &= \sum_{N=0}^{\infty} u^N \mathbb{Z}_I(\beta|N) = \sum_{N=0}^{\infty} u^N \sum_{\{N_\mathbf{n}\}} \exp\left(-\beta \sum_\mathbf{n} \varepsilon_\mathbf{n} N_\mathbf{n}\right) \delta_{N,\sum_\mathbf{n} N_\mathbf{n}} \\ &= \sum_{\{N_\mathbf{n}\}} \exp\left(-\beta \sum_\mathbf{n} \varepsilon_\mathbf{n} N_\mathbf{n}\right) \sum_{N=0}^{\infty} u^{\sum_\mathbf{n} N_\mathbf{n}} \delta_{N,\sum_\mathbf{n} N_\mathbf{n}} \\ &= \sum_{\{N_\mathbf{n}\}} \exp\left(-\beta \sum_\mathbf{n} \varepsilon_\mathbf{n} N_\mathbf{n}\right) u^{\sum_\mathbf{n} N_\mathbf{n}} \quad \Longrightarrow \end{aligned}$$

$$\Xi(\beta, u) = \prod_\mathbf{n} \left\{ \sum_{N_\mathbf{n}} \left[u \exp(-\beta \varepsilon_\mathbf{n})\right]^{N_\mathbf{n}} \right\}. \tag{6.104}$$

Fermions

For fermions, the number $N_\mathbf{n}$ can take only values $N_\mathbf{n} = 0$ and $N_\mathbf{n} = 1$. Hence, for fermions (denoted by the index F), we obtain:

$$\Xi_F(\beta, u) = \prod_\mathbf{n} \left[1 + u \exp(-\beta \varepsilon_\mathbf{n})\right].$$

Since the n-th level of a 3D oscillator is degenerate with the degeneracy

$$g(n) = \frac{(n+1)(n+2)}{2},$$

we find that the generating function $\Xi_F(\beta, u)$ is given by

$$\Xi_F(\beta, u) = \prod_{n=0}^{\infty} \left[1 + u \exp(-\beta \varepsilon_n)\right]^{g(n)}. \tag{6.105}$$

Bosons

For bosons (denoted by the index B), $N_{\mathbf{n}} = 0, 1, \ldots, \infty$. The summations over $\{N_{\mathbf{n}}\}$ in Eq. (6.104) gives:

$$\Xi_B(\beta, u) = \prod_{n=0}^{\infty} \left[\frac{1}{1 - u \exp(-\beta \varepsilon_n)} \right]^{g(n)}. \tag{6.106}$$

The results (6.104) and (6.106) prove (for the partition function) the equivalence of the path-integral approach for identical particles [99] and of the second-quantization method.

Integral representation

Let us use the Fourier representation for the delta symbol:

$$\delta_{N, \sum_{\mathbf{n}} N_{\mathbf{n}}} = \frac{1}{2\pi} \int_0^{2\pi} \exp\left[i \left(\sum_{\mathbf{n}} N_{\mathbf{n}} - N \right) (\theta - i\zeta) \right] d\theta, \tag{6.107}$$

where ζ is an arbitrary constant. Substituting Eq. (6.107) into Eq. (6.103) we obtain

$$\begin{aligned} \mathbb{Z}_I(\beta|N) &= \sum_{\{N_{\mathbf{n}}\}} \exp\left(-\beta \sum_{\mathbf{n}} \varepsilon_{\mathbf{n}} N_{\mathbf{n}} \right) \frac{1}{2\pi} \int_0^{2\pi} \exp\left[i \left(\sum_{\mathbf{n}} N_{\mathbf{n}} - N \right) (\theta - i\zeta) \right] d\theta \\ &= \frac{1}{2\pi} \int_0^{2\pi} d\theta \exp\left[-iN(\theta - i\zeta) \right] \sum_{\{N_{\mathbf{n}}\}} \exp\left(-\beta \sum_{\mathbf{n}} \varepsilon_{\mathbf{n}} N_{\mathbf{n}} + i \sum_{\mathbf{n}} N_{\mathbf{n}} (\theta - i\zeta) \right) \\ &= \frac{1}{2\pi} \int_0^{2\pi} d\theta \exp(-iN\theta - N\zeta)\, \Xi\left(\beta, e^{i\theta + \zeta}\right) \quad \Longrightarrow \end{aligned}$$

$$\mathbb{Z}_I(\beta|N) = \frac{1}{2\pi} \int_0^{2\pi} d\theta \exp\left[\ln \Xi\left(\beta, e^{i\theta+\zeta}\right) - N\zeta - iN\theta \right]. \tag{6.108}$$

The partition function for a finite number of particles can be obtained from the generation function also by the inversion formula [100]

$$\begin{aligned} \mathbb{Z}_I(\beta|N) &= \frac{1}{2\pi i} \oint \frac{\Xi(\beta, z)}{z^{N+1}} dz \qquad (6.109) \\ &= \frac{1}{2\pi} \int_0^{2\pi} e^{\left[\ln \Xi\left(\beta, u e^{i\theta}\right) - N \ln u \right]} e^{-iN\theta} d\theta. \qquad (6.110) \end{aligned}$$

Let us denote in Eq. (6.108):

$$\zeta \equiv \ln u. \tag{6.111}$$

In these notations, Eqs. (6.108) and (6.110) are *identical*. For the numerical calculation, it is more convenient to choose in Eq. (6.107) the interval of the integration over θ as $[-\pi, \pi]$

instead of $[0, 2\pi]$, what gives:

$$\mathbb{Z}_I(\beta|N) = \frac{1}{2\pi}\int_{-\pi}^{\pi} \Phi_N(\theta)\, d\theta, \tag{6.112}$$

with the function

$$\Phi_N(\theta) = \exp\left[\ln \Xi\left(\beta, ue^{i\theta}\right) - N \ln u - iN\theta\right]. \tag{6.113}$$

The aforesaid method of derivation of the partition function [Eqs. (6.107) to (6.108)] is heuristically useful, because it allows a simple generalization to spin-mixed systems with various polarization distributions.

The two-point density-density correlation function in the operator formalism is

$$\tilde{g}(\mathbf{q}, \tau | \{N_\sigma\}, \beta) = \langle \hat{\rho}_{\mathbf{q}}(\tau)\, \hat{\rho}_{-\mathbf{q}}(0) \rangle, \tag{6.114}$$

where $\hat{\rho}_{\mathbf{q}}(t)$ is the density operator in the Heisenberg representation:

$$\hat{\rho}_{\mathbf{q}}(\tau) = \exp\left(\frac{\tau}{\hbar}\hat{H}\right) \rho_{\mathbf{q}} \exp\left(-\frac{\tau}{\hbar}\hat{H}\right). \tag{6.115}$$

In the "second-quantization" representation, $\hat{\rho}_{\mathbf{q}}(t)$ is

$$\begin{aligned}\hat{\rho}_{\mathbf{q}}(\tau) &= \sum_{\mathbf{n},\mathbf{n}'} \left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{n}'} \hat{a}^+_{\mathbf{n}}(\tau)\, \hat{a}_{\mathbf{n}'}(\tau) \\ &= \sum_{\mathbf{n},\mathbf{n}'} \left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{n}'} \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}'} \exp\left[\frac{\tau}{\hbar}\left(\varepsilon_{\mathbf{n}} - \varepsilon_{\mathbf{n}'}\right)\right].\end{aligned} \tag{6.116}$$

After substituting Eq. (6.116) into (6.114), we find that

$$\tilde{g}(\mathbf{q}, \tau | \{N_\sigma\}, \beta) = \sum_{\mathbf{n},\mathbf{n}'} \sum_{\mathbf{m},\mathbf{m}'} \left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{n}'} \left(e^{-i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{m}\mathbf{m}'} \exp\left[\frac{\tau}{\hbar}\left(\varepsilon_{\mathbf{n}} - \varepsilon_{\mathbf{n}'}\right)\right] \left\langle \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}'} \hat{a}^+_{\mathbf{m}} \hat{a}_{\mathbf{m}'} \right\rangle. \tag{6.117}$$

The operator $\hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}'} \hat{a}^+_{\mathbf{m}} \hat{a}_{\mathbf{m}'}$ has non-zero diagonal matrix elements in the basis of quantum states $\left|\Psi_{N,\{N_{\mathbf{n}}\}}\right\rangle$ only in the cases

$$\begin{cases} \mathbf{n} = \mathbf{n}' \\ \mathbf{m} = \mathbf{m}' \end{cases} \quad \text{or} \quad \begin{cases} \mathbf{n} = \mathbf{m}' \\ \mathbf{m} = \mathbf{n}' \end{cases}. \tag{6.118}$$

Hence, the average

$$\left\langle \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}'} \hat{a}^+_{\mathbf{m}} \hat{a}_{\mathbf{m}'} \right\rangle = \frac{1}{\mathbb{Z}_I(\beta|N)} \sum_{\{N_{\mathbf{n}}\}} \left\langle \Psi_{N,\{N_{\mathbf{n}}\}} \left| \exp\left(-\beta\hat{H}\right) \hat{a}^+_{\mathbf{n}} \hat{a}_{\mathbf{n}'} \hat{a}^+_{\mathbf{m}} \hat{a}_{\mathbf{m}'} \right| \Psi_{N,\{N_{\mathbf{n}}\}} \right\rangle \tag{6.119}$$

is not equal to zero only when the condition (6.118) is fulfilled. This allows us to write down the average (6.119) as

$$\begin{aligned}\left\langle \hat{a}^+_{\mathbf{n}}\hat{a}_{\mathbf{n}'}\hat{a}^+_{\mathbf{m}}\hat{a}_{\mathbf{m}'}\right\rangle &= \delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\left(1-\delta_{\mathbf{mn}}\right)\left\langle \hat{a}^+_{\mathbf{n}}\hat{a}_{\mathbf{n}}\hat{a}^+_{\mathbf{m}}\hat{a}_{\mathbf{m}}\right\rangle + \delta_{\mathbf{m}'\mathbf{n}}\delta_{\mathbf{n}'\mathbf{m}}\left(1-\delta_{\mathbf{mn}}\right)\left\langle \hat{a}^+_{\mathbf{n}}\hat{a}_{\mathbf{m}}\hat{a}^+_{\mathbf{m}}\hat{a}_{\mathbf{n}}\right\rangle \\ &+ \delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\delta_{\mathbf{mn}}\left\langle \hat{a}^+_{\mathbf{n}}\hat{a}_{\mathbf{n}}\hat{a}^+_{\mathbf{n}}\hat{a}_{\mathbf{n}}\right\rangle \\ &= \delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\left(1-\delta_{\mathbf{mn}}\right)\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle + \delta_{\mathbf{m}'\mathbf{n}}\delta_{\mathbf{n}'\mathbf{m}}\left(1-\delta_{\mathbf{mn}}\right)\left(\left\langle \hat{N}_{\mathbf{n}}\right\rangle - \left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle\right) \\ &+ \delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\delta_{\mathbf{mn}}\bar{N}_{\mathbf{n}} \\ &= \delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle + \delta_{\mathbf{m}'\mathbf{n}}\delta_{\mathbf{n}'\mathbf{m}}\left\langle \hat{N}_{\mathbf{n}}\left(1-\hat{N}_{\mathbf{m}}\right)\right\rangle. \qquad (6.120)\end{aligned}$$

Here, the notation is used for the average occupation number $\bar{N}_{\mathbf{n}}$:

$$\left\langle \hat{N}_{\mathbf{n}}\right\rangle = \frac{1}{\mathbb{Z}_I\left(\beta|N\right)}\sum_{\{N_{\mathbf{n}}\}}\left\langle \Psi_{N,\{N_{\mathbf{n}}\}}\left|\exp\left(-\beta\hat{H}\right)\hat{N}_{\mathbf{n}}\right|\Psi_{N,\{N_{\mathbf{n}}\}}\right\rangle. \qquad (6.121)$$

$$\begin{aligned}\mathbb{Z}_I\left(\beta|N\right) &= \sum_{\{N_{\mathbf{n}}\}}\left\langle \Psi_{N,\{N_{\mathbf{n}}\}}\left|\exp\left(-\beta\hat{H}\right)\right|\Psi_{N,\{N_{\mathbf{n}}\}}\right\rangle \\ &= \left[\sum_{\{N_{\mathbf{n}}\}}\exp\left(-\beta\sum_{\mathbf{n}}\varepsilon_{\mathbf{n}}N_{\mathbf{n}}\right)\right]_{\sum_{\mathbf{n}}N_{\mathbf{n}}=N}. \qquad (6.122)\end{aligned}$$

In the same way as Eq. (6.102), the average (6.121) can be written down in the form

$$\left\langle \hat{N}_{\mathbf{n}}\right\rangle = \frac{1}{\mathbb{Z}_I\left(\beta|N\right)}\left[\sum_{\{N_{\mathbf{n}'}\}}N_{\mathbf{n}}\exp\left(-\beta\sum_{\mathbf{n}'}\varepsilon_{\mathbf{n}'}N_{\mathbf{n}'}\right)\right]_{\sum_{\mathbf{n}'}N_{\mathbf{n}'}=N}.$$

$$\left\langle \hat{N}_{\mathbf{n}}\right\rangle = -\frac{1}{\beta\mathbb{Z}_I\left(\beta|N\right)}\frac{\delta\mathbb{Z}_I\left(\beta|N\right)}{\delta\varepsilon_{\mathbf{n}}},$$

$$\left\langle \hat{N}_{\mathbf{n}}\right\rangle = \frac{1}{2\pi\mathbb{Z}_I\left(\beta|N\right)}\int_{-\pi}^{\pi}\frac{\Phi_N\left(\theta\right)}{\exp\left(\beta\varepsilon_{\mathbf{n}}-\zeta-i\theta\right)+1}d\theta. \qquad (6.123)$$

Since $\varepsilon_{\mathbf{n}} = \varepsilon_n$, $\left\langle \hat{N}_{\mathbf{n}}\right\rangle$ depends only on n.

Using Eq. (6.111), we can write $\bar{N}_n$ as

$$\begin{aligned}\left\langle \hat{N}_{\mathbf{n}}\right\rangle &= \frac{1}{2\pi\mathbb{Z}_I\left(\beta|N\right)}\int_{-\pi}^{\pi}\frac{\Phi_N\left(\theta\right)}{\frac{1}{u}\exp\left(\beta\varepsilon_n-i\theta\right)+1}d\theta \\ &= \frac{1}{2\pi\mathbb{Z}_I\left(\beta|N\right)}\int_{-\pi}^{\pi}\frac{\exp\left[\ln\Xi\left(\beta,ue^{i\theta}\right)-N\ln u-iN\theta\right]}{\frac{1}{u}\exp\left(\beta\varepsilon_n-i\theta\right)+1}d\theta \\ &= \frac{1}{2\pi\mathbb{Z}_I\left(\beta|N\right)}\int_{-\pi}^{\pi}\frac{\Xi\left(\beta,ue^{i\theta}\right)}{u^{N-1}}\frac{\exp\left[-i\theta\left(N-1\right)-\beta\varepsilon_n\right]}{1+u\exp\left(i\theta-\beta\varepsilon_n\right)}d\theta. \qquad (6.124)\end{aligned}$$

The averages $\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle$ for $\mathbf{m}\neq\mathbf{n}$ can be also expressed in terms of the integral representation:

$$\begin{aligned}
\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle\Big|_{\mathbf{m}\neq\mathbf{n}} &= \frac{1}{\mathbb{Z}_I\left(\beta|N\right)}\sum_{\{N_{\mathbf{n}}\}}\left\langle \Psi_{N,\{N_{\mathbf{n}}\}}\left|\exp\left(-\beta\hat{H}\right)\hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right|\Psi_{N,\{N_{\mathbf{n}}\}}\right\rangle\Bigg|_{\mathbf{m}\neq\mathbf{n}} \\
&= \frac{1}{\mathbb{Z}_I\left(\beta|N\right)}\left[\sum_{\{N_{\mathbf{n}'}\}}N_{\mathbf{n}}N_{\mathbf{m}}\exp\left(-\beta\sum_{\mathbf{n}'}\varepsilon_{\mathbf{n}'}N_{\mathbf{n}'}\right)\right]_{\sum_{\mathbf{n}'}N_{\mathbf{n}'}=N,\quad \mathbf{m}\neq\mathbf{n}} \\
&= \frac{1}{\mathbb{Z}_I\left(\beta|N\right)\beta^2}\frac{\delta^2\mathbb{Z}_I\left(\beta|N\right)}{\delta\varepsilon_{\mathbf{m}}\delta\varepsilon_{\mathbf{n}}} \\
&= \frac{1}{\mathbb{Z}_I\left(\beta|N\right)\beta^2}\frac{\delta^2}{\delta\varepsilon_{\mathbf{m}}\delta\varepsilon_{\mathbf{n}}}\left(\frac{1}{2\pi}\int_0^{2\pi}d\theta\exp\left[\ln\Xi\left(\beta,e^{i\theta+\zeta}\right)-N\zeta-iN\theta\right]\right)\quad\Rightarrow
\end{aligned}$$

We obtain the integral representation for the average of the product of operators $\hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}$ for $\mathbf{m}\neq\mathbf{n}$:

$$\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle\Big|_{\mathbf{m}\neq\mathbf{n}} = \frac{1}{2\pi\mathbb{Z}_I\left(N\right)}\int_{-\pi}^{\pi}\frac{\Phi_N\left(\theta\right)}{\left[\exp\left(\beta\varepsilon_{\mathbf{n}}-\zeta-i\theta\right)+1\right]\left[\exp\left(\beta\varepsilon_{\mathbf{m}}-\zeta-i\theta\right)+1\right]}d\theta. \quad (6.125)$$

Let us introduce the notation

$$f\left(\varepsilon,\theta\right)\equiv\frac{1}{\exp\left(\beta\varepsilon-\zeta-i\theta\right)+1}, \quad (6.126)$$

which formally coincides with the Fermi distribution function of the energy ε with the "chemical potential" $\left(\zeta+i\theta\right)/\beta$. Using this notation, the averages (6.123) and (6.125) can be written down in the form

$$\left\langle \hat{N}_{\mathbf{n}}\right\rangle = \frac{1}{2\pi\mathbb{Z}_I\left(\beta|N\right)}\int_{-\pi}^{\pi}f\left(\varepsilon_{\mathbf{n}},\theta\right)\Phi_N\left(\theta\right)d\theta, \quad (6.127)$$

$$\left\langle \hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle\Big|_{\mathbf{m}\neq\mathbf{n}} = \frac{1}{2\pi\mathbb{Z}_I\left(N\right)}\int_{-\pi}^{\pi}f\left(\varepsilon_{\mathbf{n}},\theta\right)f\left(\varepsilon_{\mathbf{m}},\theta\right)\Phi_N\left(\theta\right)d\theta. \quad (6.128)$$

We can develop the aforesaid procedure for the average of a product of any number of operators $\hat{N}_{\mathbf{n}_1}\hat{N}_{\mathbf{n}_2}\ldots\hat{N}_{\mathbf{n}_K}$, where all quantum numbers $\mathbf{n}_1,\mathbf{n}_2,\ldots,\mathbf{n}_K$ are *different*. The result is:

$$\left\langle \hat{N}_{\mathbf{n}_1}\hat{N}_{\mathbf{n}_2}\ldots\hat{N}_{\mathbf{n}_K}\right\rangle\Big|_{\mathbf{n}_j\neq\mathbf{n}_l} = \frac{1}{2\pi\mathbb{Z}_I\left(N\right)}\int_{-\pi}^{\pi}f\left(\varepsilon_{\mathbf{n}_1},\theta\right)f\left(\varepsilon_{\mathbf{n}_2},\theta\right)\ldots f\left(\varepsilon_{\mathbf{n}_K},\theta\right)\Phi_N\left(\theta\right)d\theta. \quad (6.129)$$

It should be emphasized, that all expressions above [including Eq. (6.129)] are derived for a *canonical* Hibbs ensemble (i. e., for a fixed number of particles) and for both closed-shell and open-shell systems.

Let us substitute the average (6.120) into the correlation function $\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta)$:

$$\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta) = \sum_{\mathbf{n},\mathbf{n}'}\sum_{\mathbf{m},\mathbf{m}'}\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{n}'}\left(e^{-i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{m}\mathbf{m}'}\exp\left[\frac{\tau}{\hbar}\left(\varepsilon_{\mathbf{n}}-\varepsilon_{\mathbf{n}'}\right)\right]$$
$$\times\left(\delta_{\mathbf{n}'\mathbf{n}}\delta_{\mathbf{m}'\mathbf{m}}\left\langle\hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle+\delta_{\mathbf{m}'\mathbf{n}}\delta_{\mathbf{n}'\mathbf{m}}\left\langle\hat{N}_{\mathbf{n}}\left(1-\hat{N}_{\mathbf{m}}\right)\right\rangle\right)$$

$\Longrightarrow$

$$\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_\sigma\right\},\beta) = \sum_{\mathbf{n},\mathbf{m}}\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{n}}\left(e^{-i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{m}\mathbf{m}}\left\langle\hat{N}_{\mathbf{n}}\hat{N}_{\mathbf{m}}\right\rangle$$
$$+\sum_{\mathbf{n},\mathbf{m}}\left|\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{m}}\right|^2\exp\left[\frac{\tau}{\hbar}\left(\varepsilon_{\mathbf{n}}-\varepsilon_{\mathbf{m}}\right)\right]\left\langle\hat{N}_{\mathbf{n}}\left(1-\hat{N}_{\mathbf{m}}\right)\right\rangle. \qquad (6.130)$$

The matrix elements $\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{m}}$ has the following form

$$\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{n}\mathbf{m}} = \left\langle m_1\left|e^{iq_1\hat{x}_1}\right|n_1\right\rangle\left\langle m_2\left|e^{iq_2\hat{x}_2}\right|n_2\right\rangle\left\langle m_3\left|e^{iq_3\hat{x}_3}\right|n_3\right\rangle,$$

where $\left\langle m\left|e^{iq\hat{x}}\right|n\right\rangle$ is the matrix element of a one-dimensional oscillator with the frequency w:

$$\left\langle m\left|e^{iq\hat{x}}\right|n\right\rangle = \exp\left(-\frac{\gamma^2}{2}\right)\left(i\gamma\right)^{n_>-n_<}\sqrt{\frac{n_<!}{n_>!}}L_{n_<}^{\left(n_>-n_<\right)}\left(\gamma^2\right),$$
$$\begin{cases} n_> \equiv \max\left(n,m\right); \\ n_< \equiv \min\left(n,m\right), \end{cases} \qquad (6.131)$$

$\gamma\equiv q\sqrt{\frac{\hbar}{2mw}}$, $\left|n\right\rangle$ are the quantum states of the one-dimensional oscillator with the frequency w, $L_n^{(\alpha)}\left(z\right)$ is the generalized Laguerre polynomial.

System with mixed spins

The correlation functions for a system with mixed spins can be explicitly derived by the generalization of Eqs. (6.112) and (6.129) to the case of the particles with the non-zero spin. We use the fact, that the derivation of Eqs. (6.112) and (6.129), performed in this section, does not depend on the concrete form of the energy spectrum $\varepsilon_{\mathbf{n}}$. Hence, in the formulae, derived above, the replacement should be made:

$$|\mathbf{n}\rangle \to |\mathbf{n},\sigma\rangle\,, \quad \hat{N}_{\mathbf{n}} \to \hat{N}_{\mathbf{n},\sigma} = a^{+}_{\mathbf{n},\sigma} a_{\mathbf{n},\sigma}, \tag{6.132}$$

where σ is the electron spin projection. Consequently, the matrix elements $\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{mn}}$ are replaced by

$$\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{mn}} \to \left\langle \mathbf{m},\sigma\left|e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right|\mathbf{n},\sigma'\right\rangle = \delta_{\sigma\sigma'}\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{mn}}. \tag{6.133}$$

Taking into account Eqs. (6.132) and (6.133), the two-point correlation function (6.130) becomes

$$\begin{aligned} \tilde{g}\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta) = &\sum_{\mathbf{n},\mathbf{m}}\sum_{\sigma_1,\sigma_2}\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{nn}}\left(e^{-i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{mm}}\left\langle \hat{N}_{\mathbf{n},\sigma_1}\hat{N}_{\mathbf{m},\sigma_2}\right\rangle \\ &+\sum_{\mathbf{n},\mathbf{m}}\sum_{\sigma}\left|\left(e^{i\mathbf{q}\cdot\hat{\mathbf{x}}}\right)_{\mathbf{nm}}\right|^{2}\exp\left[\frac{\tau}{\hbar}\left(\varepsilon_{\mathbf{n}}-\varepsilon_{\mathbf{m}}\right)\right]\left\langle \hat{N}_{\mathbf{n},\sigma}\left(1-\hat{N}_{\mathbf{m},\sigma}\right)\right\rangle. \end{aligned} \tag{6.134}$$

The averages $\left\langle \hat{N}_{\mathbf{n},\sigma}\right\rangle$ and $\left\langle \hat{N}_{\mathbf{n},\sigma_1}\hat{N}_{\mathbf{m},\sigma_2}\right\rangle$ are, respectively, one-particle and two-particle distribution functions,

$$\left\langle \hat{N}_{\mathbf{n},\sigma}\right\rangle \equiv f_1\left(n,\sigma|N_{\sigma},\beta\right), \tag{6.135}$$

$$\left\langle \hat{N}_{\mathbf{n},\sigma}\hat{N}_{\mathbf{n}',\sigma'}\right\rangle \equiv f_2\left(n,\sigma;n',\sigma'|\left\{N_{\sigma}\right\},\beta\right). \tag{6.136}$$

The one-electron distribution function $f_1\left(n,\sigma|N_{\sigma},\beta\right)$ is the average number of electrons with the spin projection σ at the n-th energy level, while the two-electron distribution function $f_2\left(n,\sigma;n',\sigma'|\left\{N_{\sigma}\right\},\beta\right)$ is the average product of the numbers of electrons with the spin projections σ and σ' at the levels n and n'. These functions are expressed through the following integrals [see (6.127), (6.128)]:

$$f_1\left(n,\sigma|N_{\sigma},\beta\right) = \frac{1}{2\pi\mathbb{Z}_F\left(N_{\sigma},w,\beta\right)}\int_{-\pi}^{\pi} f\left(\varepsilon_n,\theta\right)\Phi\left(\theta,\beta,N_{\sigma}\right)d\theta, \tag{6.137}$$

$$f_2\left(n,\sigma;n',\sigma'|\left\{N_{\sigma}\right\},\beta\right) = \begin{cases} \frac{1}{2\pi\mathbb{Z}_F\left(N_{\sigma},w,\beta\right)}\int_{-\pi}^{\pi} f\left(\varepsilon_n,\theta\right)f\left(\varepsilon_{n'},\theta\right)\Phi\left(\theta,\beta,N_{\sigma}\right)d\theta, & \text{if } \sigma'=\sigma; \\ f_1\left(n,\sigma|N_{\sigma},\beta\right)f_1\left(n,\sigma'|N_{\sigma'},\beta\right), & \text{if } \sigma'\neq\sigma \end{cases} \tag{6.138}$$

with the notations

$$\Phi\left(\theta,\beta,N_{\sigma}\right) = \exp\left[\sum_{n=0}^{\infty}\ln\left(1+e^{i\theta+\xi-\beta\varepsilon_n}\right) - N_{\sigma}\left(\xi+i\theta\right)\right], \tag{6.139}$$

$$f\left(\varepsilon,\theta\right) \equiv \frac{1}{\exp\left(\beta\varepsilon-\xi-i\theta\right)+1}. \tag{6.140}$$

The function $f\left(\varepsilon,\theta\right)$ formally coincides with the Fermi-Dirac distribution function of the energy ε with the "chemical potential" $\left(\xi+i\theta\right)/\beta$.

Here we consider the zero-temperature limit, for which the integrals (6.137) and (6.138) can be calculated analytically. The result for the one-electron distribution function is

$$\left. f_1\left(n,\sigma|\beta,N_\sigma\right)\right|_{\beta\to\infty} = \begin{cases} 1, & n<L_\sigma; \\ 0, & n>L_\sigma; \\ \frac{N_\sigma-N_{L_\sigma}}{g_{L_\sigma}}, & n=L_\sigma. \end{cases} \tag{6.141}$$

According to (6.141), L_σ is the number of the lowest open shell, and

$$g_n = \begin{cases} \frac{1}{2}\left(n+1\right)\left(n+2\right) & \left(3D\right), \\ n+1 & \left(2D\right). \end{cases}$$

is the degeneracy of the n-th shell. N_{L_σ} is the number of electrons in all the closed shells with the spin projection σ,

$$N_{L_\sigma} \equiv \sum_{n=0}^{L_\sigma-1} g_n = \begin{cases} \frac{1}{6}L_\sigma\left(L_\sigma+1\right)\left(L_\sigma+2\right) & \left(3D\right), \\ \frac{1}{2}L_\sigma\left(L_\sigma+1\right) & \left(2D\right). \end{cases} \tag{6.142}$$

The two-electron distribution function $f_2\left(n,\sigma;n',\sigma'|\left\{N_\sigma\right\},\beta\right)$ at $T=0$ takes the form

$$\begin{aligned} &\left. f_2\left(n,\sigma;n',\sigma'|\beta,\left\{N_\sigma\right\}\right)\right|_{\beta\to\infty} \\ &= \begin{cases} \left. f_1\left(n,\sigma|\beta,N_\sigma\right)\right|_{\beta\to\infty} \left. f_1\left(n',\sigma'|\beta,N_{\sigma'}\right)\right|_{\beta\to\infty}, & n\neq n' \text{ or } \sigma\neq\sigma' \\ 1, & \sigma=\sigma' \text{ and } n=n'<L_\sigma; \\ 0, & \sigma=\sigma' \text{ and } n=n'>L_\sigma; \\ \frac{N_\sigma-N_{L_\sigma}}{g_{L_\sigma}}\frac{N_\sigma-N_{L_\sigma}-1}{g_{L_\sigma}-1}, & \sigma=\sigma' \text{ and } n=n'=L_\sigma. \end{cases} \end{aligned} \tag{6.143}$$

In summary, we have obtained the following expression for $\tilde{g}\left(\mathbf{q},\tau|\left\{N_\sigma\right\},\beta\right)$:

$$\begin{aligned} \tilde{g}\left(\mathbf{q},\tau|\left\{N_\sigma\right\},\beta\right) = & \sum_{\mathbf{n},\sigma,\mathbf{n}',\sigma'} \left(e^{i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{nn}} \left(e^{-i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{n}'\mathbf{n}'} f_2\left(n,\sigma;n',\sigma'|\left\{N_\sigma\right\},\beta\right) \\ & + \sum_{\mathbf{n},\mathbf{n}',\sigma} \left|\left(e^{i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{nn}'}\right|^2 \exp\left[\frac{\tau}{\hbar}\left(\varepsilon_n-\varepsilon_{n'}\right)\right] \\ & \times \left[f_1\left(n,\sigma|\left\{N_\sigma\right\},\beta\right) - f_2\left(n,\sigma;n',\sigma|\left\{N_\sigma\right\},\beta\right)\right]. \end{aligned} \tag{6.144}$$

This formula is valid for both closed and open shells. The correlation functions derived in this subsection are used both for the calculation of the ground-state energy and, a shown below, for the calculation of the optical conductivity of an N-polaron system in a quantum dot.

E. Many-polaron ground state in a quantum dot: extrapolation to the homogeneous limit and comparison to the results for a polaron gas in bulk [88]

The correlation function given by Eq. (6.144) can be subdivided as

$$\tilde{g}\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta\right)=\tilde{g}_1\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta\right)+\tilde{g}_2\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta\right), \tag{6.145}$$

with

$$\begin{aligned}\tilde{g}_1\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta\right)\equiv&\sum_{\mathbf{n},\mathbf{n}',\sigma}\left|\left(e^{i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{n}\mathbf{n}'}\right|^2\exp\left[\frac{\tau}{\hbar}\left(\varepsilon_n-\varepsilon_{n'}\right)\right]\\&\times\left[f_1\left(n,\sigma\right|\left\{N_{\sigma}\right\},\beta\right)-f_2\left(n,\sigma;n',\sigma\right|\left\{N_{\sigma}\right\},\beta\right)\right],\end{aligned} \tag{6.146}$$

$$\tilde{g}_2\left(\mathbf{q},\tau\right|\left\{N_{\sigma}\right\},\beta\right)\equiv\sum_{\mathbf{n},\sigma,\mathbf{n}',\sigma'}\left(e^{i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{n}\mathbf{n}}\left(e^{-i\mathbf{q}\cdot\mathbf{x}}\right)_{\mathbf{n}'\mathbf{n}'}f_2\left(n,\sigma;n',\sigma'\right|\left\{N_{\sigma}\right\},\beta\right). \tag{6.147}$$

In accordance with the subdivision (6.145) of the correlation function, we subdivide the Coulomb and polaron contributions:

$$E_C=E_C^{(1)}+E_C^{(2)}, \tag{6.148}$$

$$E_{e-ph}=E_{e-ph}^{(1)}+E_{e-ph}^{(2)}. \tag{6.149}$$

We have numerically checked whether the polaron contribution per particle $E_{e-ph}^{(1)}/N$ tends to a finite value at $N\to\infty$. In Figs. 14 and 15, we have plotted the polaron contributions $E_{e-ph}^{(1)}/N$ as a function of N for a quantum dot in ZnO and in a polar medium with $\alpha=5$, $\eta=0.3$, respectively. [3] As seen from Fig. 14, the polaron contribution $E_{e-ph}^{(1)}/N$ in ZnO as a function of N oscillates taking expressed maxima for N corresponding to the closed shells $N=2,8,20,40,\ldots$. There exist kinks of $E_{e-ph}^{(1)}/N$ at N corresponding to the half-filled shells, but these kinks are extremely small. In the case of the medium with $\alpha=5$, $\eta=0.3$, for $r_s^*=20$ (what corresponds to the density $n_0\approx1.14\times10^{18}$ cm^{-3}), the polaron contribution $E^{(1)}_{e-ph}/N$ oscillates taking maximal values at the numbers of fermions, which correspond to the closed shells for a spin-polarized system with parallel spins.

---

[3] Since, as discussed above, for a single polaron only the whole polaron contribution $E_{e-ph}=E_{e-ph}^{(1)}+E_{e-ph}^{(2)}$ has a physical meaning, the plots for $E_{e-ph}^{(1)}$ in Figs. 3 and 4 start from $N=2$. The total polaron contribution E_{e-ph} for $N=1$ is plotted below, in Fig. 9.

In Figs. 14 and 15, the dashed curves are the envelopes for local maxima (closed shells) and local minima of $E^{(1)}_{e-ph}/N$.We see that when these envelopes are *extrapolated* to larger number of fermions, the distance between the envelopes decreases. Therefore, the magnitude of the variations of $E^{(1)}_{e-ph}/N$ related to the shell filling diminishes with increasing N, and it is safe to suppose that in the limit of large N, the envelopes tend to one and the same value. That value corresponds to the *homogeneous* ("bulk") limit $\lim_{N\to\infty}\left(E^{(1)}_{e-ph}/N\right)$.



FIG. 14: Polaron contribution $E^{(1)}_{e-ph}/N$ as a function of N for a ZnO quantum dot.The material parameters for ZnO are taken from Ref. [89]. The value $r_s^* = 2$ corresponds to $n_0 = 4.34 \times 10^{19}$ cm^{-2}. Inset: the total spin as a function of N.



FIG. 15: Polaron contribution $E^{(1)}_{e-ph}/N$ as a function of N for a quantum dot of a polar medium with $\alpha = 5$, $\eta = 0.3$. The value $r_s^* = 20$ corresponds to $n_0 = 1.14 \times 10^{18}$ cm^{-3}. Inset: the total spin as a function of N.

In Fig. 16, we compare the polaron contribution $E^{(1)}_{e-ph}/N$ calculated within our variational path-integral method for different numbers of fermions with the polaron contribution to the ground-state energy per particle for a polaron gas in bulk, calculated (i) in Ref. [101] within an intermediate-coupling approach (the thin solid curve), (ii) in Ref. [102], using a variational approach developed first in Ref. [89]. As seen from this figure, our all-coupling variational method provides lower values for the polaron contribution than those obtained in Refs. [101, 102]. The difference between the polaron contribution calculated within our method and that of Ref. [101] is smaller at low densities and increases in magnitude with increasing density. The difference between the polaron contribution calculated within our method and that of Ref. [102] very slightly depends on the density. The result of Ref. [102] becomes closer to our result only at high densities.



FIG. 16: Polaron contribution to the polaron ground-state energy per particle $E^{(1)}_{e-ph}$ in an N-polaron quantum dot as a function of the effective density. The parameters are taken for ZnO (see Ref. [89]): $\alpha = 0.849$, $\varepsilon_0 = 8.15$, $\varepsilon_\infty = 4.0$, $\hbar\omega_{\mathrm{LO}} = 73.27$ meV, $m_b = 0.24 m_e$, where m_e is the electron mass in the vacuum. This polaron contribution is compared with the polaron contribution to the ground-state energy of a polaron gas in bulk calculated in Refs. [101, 102].

F. Optical conductivity

In Ref. [96] we have extended the memory-function approach to a system of arbitrary-coupling interacting polarons confined to a parabolic confinement potential. The optical conductivity relates the current $\mathbf{J}\left(t\right)$ per electron to a time-dependent uniform electric field $\mathbf{E}\left(t\right)$ in the framework of linear response theory. Further on, the Fourier components of the electric field are denoted by $\mathbf{E}\left(\omega\right)$:

$$\mathbf{E}\left(t\right)=\frac{1}{2\pi}\int_{-\infty}^{\infty}\mathbf{E}\left(\omega\right)e^{-i\omega t}d\omega, \tag{6.150}$$

and the similar denotations are used for other time-dependent quantities. The electric current per electron $\mathbf{J}\left(t\right)$ is related to the mean electron coordinate response $\mathbf{R}\left(t\right)$ by

$$\mathbf{J}\left(t\right)=-e\frac{d\mathbf{R}\left(t\right)}{dt}, \tag{6.151}$$

and hence

$$\mathbf{J}\left(\omega\right)=ie\omega\mathbf{R}\left(\omega\right). \tag{6.152}$$

Within the linear-response theory, both the electric current and the coordinate response are proportional to $\mathbf{E}\left(\omega\right)$:

$$\mathbf{J}\left(\omega\right)=\sigma\left(\omega\right)\mathbf{E}\left(\omega\right), \quad \mathbf{R}\left(\omega\right)=\frac{\sigma\left(\omega\right)}{ie\omega}\mathbf{E}\left(\omega\right), \tag{6.153}$$

where $\sigma\left(\omega\right)$ is the conductivity per electron. Because we treat an isotropic electron-phonon system, $\sigma\left(\omega\right)$ is a scalar function. It is determined from the time evolution of the center-of-mass coordinate:

$$\mathbf{R}\left(t\right)\equiv\frac{1}{N}\left\langle\left\langle\sum_{j=1}^{N}\mathbf{x}_{j}\left(t\right)\right\rangle\right\rangle_{S}. \tag{6.154}$$

The symbol $\langle\langle(\bullet)\rangle\rangle_{S}$ denotes an average in the *real-time* representation for a system with action functional S:

$$\langle\langle(\bullet)\rangle\rangle_{S}\equiv\int d\bar{\mathbf{x}}\int d\bar{\mathbf{x}}_{0}\int d\bar{\mathbf{x}}_{0}^{\prime}\int_{\bar{\mathbf{x}}_{0}}^{\bar{\mathbf{x}}}D\bar{\mathbf{x}}\left(t\right)\int_{\bar{\mathbf{x}}_{0}^{\prime}}^{\bar{\mathbf{x}}}D\bar{\mathbf{x}}^{\prime}\left(t\right)e^{\frac{i}{\hbar}S\left[\bar{\mathbf{x}}(t),\bar{\mathbf{x}}^{\prime}(t)\right]}\left(\bullet\right)\left\langle\bar{\mathbf{x}}_{0}\left|\hat{\rho}\left(t_{0}\right)\right|\bar{\mathbf{x}}_{0}^{\prime}\right\rangle\Big|_{t_{0}\rightarrow-\infty}, \tag{6.155}$$

where $\left\langle\bar{\mathbf{x}}_{0}\left|\hat{\rho}\left(t_{0}\right)\right|\bar{\mathbf{x}}_{0}^{\prime}\right\rangle$ is the density matrix before the onset of the electric field in the infinite past ($t_{0}\rightarrow-\infty$). The corresponding action functional is

$$S\left[\bar{\mathbf{x}}\left(t\right),\bar{\mathbf{x}}^{\prime}\left(t\right)\right]=\int_{-\infty}^{t}\left[L_{e}\left(\dot{\bar{\mathbf{x}}}\left(t\right),\bar{\mathbf{x}}\left(t\right),t\right)-L_{e}\left(\dot{\bar{\mathbf{x}}}^{\prime}\left(t\right),\bar{\mathbf{x}}^{\prime}\left(t\right),t\right)\right]dt^{\prime}-i\hbar\Phi\left[\bar{\mathbf{x}}\left(t\right),\bar{\mathbf{x}}^{\prime}\left(t\right)\right], \tag{6.156}$$

where $L_e\left(\dot{\bar{\mathbf{x}}}, \bar{\mathbf{x}}, t\right)$ is the Lagrangian of N interacting electrons in a time-dependent uniform electric field $\mathbf{E}(t)$

$$L_e\left(\dot{\bar{\mathbf{x}}}, \bar{\mathbf{x}}, t\right) = \sum_{\sigma}\sum_{j=1}^{N_\sigma}\left(\frac{m_b \dot{\mathbf{x}}_{j,\sigma}^2}{2} - \frac{m_b\Omega_0^2\mathbf{x}_{j,\sigma}^2}{2} - e\mathbf{x}_{j,\sigma}\cdot\mathbf{E}(t)\right) - \sum_{\sigma,\sigma'}\sum_{j=1}^{N_\sigma}\sum_{\substack{l=1 \\ (j,\sigma)\neq(l,\sigma')}}^{N_{\sigma'}} \frac{e^2}{2\varepsilon_\infty\left|\mathbf{x}_{j,\sigma}-\mathbf{x}_{l,\sigma'}\right|}. \tag{6.157}$$

The influence phase of the phonons

$$\Phi\left[\bar{\mathbf{x}}(s), \bar{\mathbf{x}}'(s)\right] = -\sum_{\mathbf{q}}\frac{\left|V_{\mathbf{q}}\right|^2}{\hbar^2}\int_{-\infty}^{t} ds \int_{-\infty}^{s} ds' \left[\rho_{\mathbf{q}}(s) - \rho'_{\mathbf{q}}(s)\right]$$
$$\times\left[T^*_{\omega_{\mathbf{q}}}(s-s')\rho_{\mathbf{q}}(s') - T_{\omega_{\mathbf{q}}}(s-s')\rho'_{\mathbf{q}}(s')\right] \tag{6.158}$$

describes both a retarded interaction between different electrons and a retarded self-interaction of each electron due to the elimination of the phonon coordinates. This functional contains the free-phonon Green's function:

$$T_\omega(t) = \frac{e^{i\omega t}}{1-e^{-\beta\hbar\omega}} + \frac{e^{-i\omega t}}{e^{\beta\hbar\omega}-1}. \tag{6.159}$$

The equation of motion for $\mathbf{R}(t)$ is

$$m_b\frac{d^2\mathbf{R}(t)}{dt^2} + m_b\Omega_0^2\mathbf{R}(t) + e\mathbf{E}(t) = \mathbf{F}_{ph}(t), \tag{6.160}$$

where $\mathbf{F}_{ph}(t)$ is the average force due to the electron-phonon interaction,

$$\mathbf{F}_{ph}(t) = -\operatorname{Re}\sum_{\mathbf{q}}\frac{2\left|V_{\mathbf{q}}\right|^2\mathbf{q}}{N\hbar}\int_{-\infty}^{t} ds\, T^*_{\omega_{\mathrm{LO}}}(t-s)\left\langle\left\langle\rho_{\mathbf{q}}(t)\rho_{-\mathbf{q}}(s)\right\rangle\right\rangle_S. \tag{6.161}$$

The two-point correlation function $\left\langle\left\langle\rho_{\mathbf{q}}(t)\rho_{-\mathbf{q}}(s)\right\rangle\right\rangle_S$ should be calculated from Eq. (6.155) using the exact action (6.156), but like for the free energy above, this path integral cannot be calculated analytically. Instead, we perform an approximate calculation, replacing the two-point correlation function in Eq. (6.161) by $\left\langle\left\langle\rho_{\mathbf{q}}(t)\rho_{-\mathbf{q}}(s)\right\rangle\right\rangle_{S_0}$, where $S_0\left[\bar{\mathbf{x}}(t), \bar{\mathbf{x}}'(t)\right]$ is the action functional with the optimal values of the variational parameters for the model system considered in the previous section in the presence of the electric field $\mathbf{E}(t)$. The functional $S_0\left[\bar{\mathbf{x}}(t), \bar{\mathbf{x}}'(t)\right]$ is quadratic and describes a system of coupled harmonic oscillators in the uniform electric field $\mathbf{E}(t)$. This field enters the term $-e\mathbf{E}(t)\cdot\sum_\sigma\sum_{j=1}^{N_\sigma}\mathbf{x}_{j,\sigma}$ in the

Lagrangian, which only affects the center-of-mass coordinate. Hence, a shift of variables to the frame of reference with the origin at the center of mass

$$\begin{cases} \mathbf{x}_n(t) = \tilde{\mathbf{x}}_n(t) + \mathbf{R}(t), \\ \mathbf{x}'_n(t) = \tilde{\mathbf{x}}'_n(t) + \mathbf{R}(t), \end{cases} \tag{6.162}$$

results in

$$\langle\langle \rho_q(t)\,\rho_{-q}(s)\rangle\rangle_{S_0} = \langle\langle \rho_q(t)\,\rho_{-q}(s)\rangle\rangle_{S_0}\Big|_{E=0} e^{i\mathbf{q}\cdot[\mathbf{R}(t)-\mathbf{R}(s)]}. \tag{6.163}$$

This result (6.163) is valid for any *quadratic* model action S_0.

The applicability of the parabolic approximation is confirmed by the fact that a self-induced polaronic potential, created by the polarization cloud around an electron, is rather well described by a parabolic potential whose parameters are determined by a variational method. For weak coupling, our variational method is at least of the same accuracy as the perturbation theory, which results from our approach at a special choice of the variational parameters. For strong coupling, an interplay of the electron-phonon interaction and the Coulomb correlations within a confinement potential can lead to the assemblage of polarons in multi-polaron systems. Our choice of the model variational system is reasonable because of this trend, apparently occurring in a many-polaron system with arbitrary N for a finite confinement strength.

The correlation function $\langle \rho_{\mathbf{q}}(t)\,\rho_{-\mathbf{q}}(s)\rangle_{S_0}\big|_{\mathbf{E}=0}$ corresponds to the model system in the absence of an electric field. For $t > s$, this function is related to the imaginary-time correlation function $\mathcal{G}(\mathbf{q},\tau|\{N_\sigma\},\beta)$, described in the previous section:

$$\langle\langle \rho_{\mathbf{q}}(t)\,\rho_{-\mathbf{q}}(s)\rangle\rangle_{S_0}\Big|_{\mathbf{E}=0,t>s} = \mathcal{G}(\mathbf{q}, i(t-s)\,|\,\{N_\sigma\},\beta). \tag{6.164}$$

Using the transformation (6.162) and the relation (6.164), we obtain from Eq. (6.161)

$$\mathbf{F}_{ph}(t) = -\operatorname{Re}\sum_{\mathbf{q}} \frac{2|V_{\mathbf{q}}|^2\mathbf{q}}{N\hbar} \int_{-\infty}^{t} T^*_{\omega_{\mathrm{LO}}}(t-s)\,\mathrm{e}^{\mathrm{i}\mathbf{q}\cdot[\mathbf{R}(t)-\mathbf{R}(s)]}\mathcal{G}(\mathbf{q}, i(t-s)\,|\,\{N_\sigma\},\beta)\,ds. \tag{6.165}$$

Within the linear-response theory, we expand the function $\mathrm{e}^{\mathrm{i}\mathbf{q}\cdot[\mathbf{R}(t)-\mathbf{R}(s)]}$ in Eq. (6.165) as a Taylor series in $[\mathbf{R}(t)-\mathbf{R}(s)]$ up to the first-order term. The zeroth-order term gives no contribution into $\mathbf{F}_{ph}(t)$ due to the symmetry of $|V_{\mathbf{q}}|^2$ and of $f_{\mathbf{q}}(t-s)$ with respect to the inversion $\mathbf{q}\to-\mathbf{q}$. In this approach, the Cartesian coordinates of the force $(j = 1,2,3)$

become

$$\left(\mathbf{F}_{ph}\left(t\right)\right)_{j} = \sum_{k=1}^{3}\sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^{2} q_{j}q_{k}}{N\hbar} \int\limits_{-\infty}^{t} \left[R_{k}\left(t\right) - R_{k}\left(s\right)\right]$$
$$\times \operatorname{Im}\left[T_{\omega_{\mathrm{LO}}}^{*}\left(t-s\right)\mathcal{G}\left(\mathbf{q}, i\left(t-s\right)\right|\left\{N_{\sigma}\right\},\beta)\right] ds. \tag{6.166}$$

Further on, we perform the Fourier expansion:

$$\mathbf{R}\left(t\right) = \frac{1}{2\pi}\int_{-\infty}^{\infty}\mathbf{R}\left(\omega\right)e^{-i\omega t}d\omega. \tag{6.167}$$

In Eq. (6.166), we make the replacement

$$\tau \equiv t - s, \quad \Longrightarrow \quad s = t - \tau,$$

what gives

$$\begin{aligned}
\left(\mathbf{F}_{ph}\left(t\right)\right)_{j} &= \sum_{k=1}^{3}\sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^{2} q_{j}q_{k}}{N\hbar} \int\limits_{0}^{\infty} d\tau \ \left[R_{k}\left(t\right) - R_{k}\left(t-\tau\right)\right] \operatorname{Im}\left[T_{\omega_{\mathrm{LO}}}^{*}\left(\tau\right)\mathcal{G}\left(\mathbf{q}, it\right|\left\{N_{\sigma}\right\},\beta)\right] \\
&= \sum_{k=1}^{3}\sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^{2} q_{j}q_{k}}{N\hbar}\frac{1}{2\pi}\int_{-\infty}^{\infty} d\omega R_{k}\left(\omega\right)e^{-i\omega t} \int\limits_{0}^{\infty} d\tau \ \left(1 - e^{i\omega\tau}\right) \operatorname{Im}\left[T_{\omega_{\mathrm{LO}}}^{*}\left(\tau\right) f_{\mathbf{q}}\left(\tau\right)\right] \\
&= \frac{1}{2\pi}\int_{-\infty}^{\infty} d\omega F_{j}\left(\omega\right)e^{-i\omega t},
\end{aligned}$$

where the Fourier component of the force is

$$\left(\mathbf{F}_{ph}\left(\omega\right)\right)_{j} = \sum_{k=1}^{3}\sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^{2} q_{j}q_{k}}{N\hbar} \int\limits_{0}^{\infty} dt \ \left(1 - e^{i\omega t}\right) \operatorname{Im}\left[T_{\omega_{\mathrm{LO}}}^{*}\left(\tau\right)\mathcal{G}\left(\mathbf{q}, it\right|\left\{N_{\sigma}\right\},\beta)\right] R_{k}\left(\omega\right). \tag{6.168}$$

The expression (6.168) can be written down as

$$\left(\mathbf{F}_{ph}\left(\omega\right)\right)_{j} = -m_{b}\sum_{k=1}^{3}\chi_{jk}\left(\omega\right)R_{k}\left(\omega\right), \tag{6.169}$$

where $\chi_{jk}\left(\omega\right)$ are components of the tensor

$$\chi_{jk}\left(\omega\right) = \sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^{2} q_{j}q_{k}}{N\hbar m_{b}} \int\limits_{0}^{\infty} dt \ \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[T_{\omega_{\mathrm{LO}}}^{*}\left(t\right)\mathcal{G}\left(\mathbf{q}, it\right|\left\{N_{\sigma}\right\},\beta)\right]. \tag{6.170}$$

In the abstract tensor form, Eq. (6.169) is

$$\mathbf{F}\left(\omega\right) = -\overleftrightarrow{\chi}\left(\omega\right)\mathbf{R}\left(\omega\right). \tag{6.171}$$

In particular, for the isotropic electron-phonon interaction and in the absence of the magnetic field, the tensor $\overleftrightarrow{\chi}(\omega)$ is proportional to the unity tensor $\mathbb{I}$,

$$\overleftrightarrow{\chi}(\omega) = \chi(\omega)\,\mathbb{I}, \tag{6.172}$$

where $\chi(\omega)$ is the scalar memory function:

$$\chi(\omega) = \sum_{\mathbf{q}} \frac{2|V_{\mathbf{q}}|^2 q^2}{3N\hbar m_b} \int_0^\infty dt\ \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[T^*_{\omega_{\mathrm{LO}}}(t)\,\mathcal{G}(\mathbf{q}, it|\{N_\sigma\}, \beta)\right]. \tag{6.173}$$

Let us perform the Fourier transformation of the equation of motion (6.160):

$$m_b\left(\Omega_0^2 - \omega^2\right)\mathbf{R}(\omega) + e\mathbf{E}(\omega) = \mathbf{F}_{ph}(\omega). \tag{6.174}$$

With Eq. (6.171), this equation takes the form

$$m_b\left(\Omega_0^2 - \omega^2\right)\mathbf{R}(\omega) + e\mathbf{E}(\omega) = -m_b\overleftrightarrow{\chi}(\omega)\,\mathbf{R}(\omega)$$

$$\Downarrow$$

$$m_b\left[\omega^2 - \Omega_0^2 - \overleftrightarrow{\chi}(\omega)\right]\mathbf{R}(\omega) = e\mathbf{E}(\omega). \tag{6.175}$$

Comparing Eqs. (6.153) and (6.175) between each other, we find that

$$m_b\left[\omega^2 - \Omega_0^2 - \overleftrightarrow{\chi}(\omega)\right]\frac{\sigma(\omega)}{ie\omega}\mathbf{E}(\omega) = e\mathbf{E}(\omega),$$

so that

$$\sigma(\omega) = \frac{ie^2\omega}{m_b}\left[\omega^2 - \Omega_0^2 - \overleftrightarrow{\chi}(\omega)\right]^{-1}.$$

In the case when Eq. (6.172) is valid, we obtain the conductivity in the scalar form

$$\sigma(\omega) = \frac{ie^2\omega}{m_b\left[\omega^2 - \Omega_0^2 - \chi(\omega)\right]}.$$

The real part of the conductivity is

$$\begin{aligned}\operatorname{Re}\sigma(\omega) &= \operatorname{Re}\frac{ie^2\omega\left[(\omega^2 - \Omega_0^2) - \operatorname{Re}\chi(\omega) + i\operatorname{Im}\chi(\omega)\right]}{m_b\left\{\left[(\omega^2 - \Omega_0^2) - \operatorname{Re}\chi(\omega)\right]^2 + \left[\operatorname{Im}\chi(\omega)\right]^2\right\}} \\ &= -\frac{e^2\omega}{m_b}\frac{\operatorname{Im}\chi(\omega)}{\left[(\omega^2 - \Omega_0^2) - \operatorname{Re}\chi(\omega)\right]^2 + \left[\operatorname{Im}\chi(\omega)\right]^2}.\end{aligned}$$

In summary, the optical conductivity can be expressed in terms of the memory function $\chi(\omega)$ (cf. Ref. [48]),

$$\operatorname{Re}\sigma(\omega) = -\frac{e^2}{m_b}\frac{\omega\operatorname{Im}\chi(\omega)}{\left[\omega^2 - \Omega_0^2 - \operatorname{Re}\chi(\omega)\right]^2 + \left[\operatorname{Im}\chi(\omega)\right]^2}, \tag{6.176}$$

where $\chi(\omega)$ is given by

$$\chi(\omega) = \sum_{\mathbf{q}} \frac{2\left|V_{\mathbf{q}}\right|^2 q^2}{3N\hbar m_b} \int\limits_0^\infty dt\, \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[T^*_{\omega_{\mathrm{LO}}}(t)\, \mathcal{G}\left(\mathbf{q}, it|\left\{N_\sigma\right\}, \beta\right)\right]. \tag{6.177}$$

It is worth noting that the optical conductivity (6.176) differs from that for a translationally invariant polaron system both by the explicit form of $\chi(\omega)$ and by the presence of the term Ω_0^2 in the denominator. For $\alpha \to 0$, the optical conductivity tends to a δ-like peak at $\omega = \Omega_0$,

$$\lim_{\alpha \to 0} \operatorname{Re}\sigma(\omega) = \frac{\pi e^2}{2m_b}\delta\left(\omega - \Omega_0\right). \tag{6.178}$$

For a translationally invariant system $\Omega_0 \to 0$, and this weak-coupling expression (6.178) reproduces the "central peak" of the polaron optical conductivity [80].

The further simplification of the memory function (6.177) is performed in the following way. With the Fröhlich amplitudes of the electron-phonon interaction, we transform the summation over $\mathbf{q}$ to the integral and use the Feynman units ($\hbar = 1$, $\omega_{\mathrm{LO}} = 1$, $m_b = 1$), in which $\left|V_{\mathbf{q}}\right|^2 = \frac{2\sqrt{2}\pi\alpha}{q^2 V}$. We also use the fact that in an isotropic crystal, $\mathcal{G}\left(\mathbf{q}, it|\left\{N_\sigma\right\}, \beta\right) = \mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)$. As a result, we find

$$\begin{aligned} \chi(\omega) &= \frac{V}{(2\pi)^3} \int_0^\infty 4\pi q^2 dq \frac{2q^2}{3N} \frac{2\sqrt{2}\pi\alpha}{q^2 V} \\ &\times \int\limits_0^\infty dt\, \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[T^*_{\omega_{\mathrm{LO}}}(t)\, \mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)\right] \\ &= \frac{2\sqrt{2}\alpha}{3\pi N} \int_0^\infty q^2 dq \int\limits_0^\infty dt\, \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[T^*_{\omega_{\mathrm{LO}}}(t)\, \mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)\right]. \end{aligned}$$

In the zero-temperature case, $T^*_{\omega_{\mathrm{LO}}}(t) \to e^{-it}$, and we arrive at the expression

$$\chi(\omega) = \frac{2\sqrt{2}\alpha}{3\pi N} \int_0^\infty q^2 dq \int\limits_0^\infty dt\, \left(e^{i\omega t} - 1\right) \operatorname{Im}\left[e^{-it} \mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)\right]. \tag{6.179}$$

Substituting the two-point correlation function $\mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)$ with the one-electron (6.141) and the two-electron (6.143) distribution functions into the memory function (6.179) and expanding $\mathcal{G}\left(q, it|\left\{N_\sigma\right\}, \beta\right)$ in powers of e^{-iwt}, $e^{-i\Omega_1 t}$ and $e^{-i\Omega_2 t}$, the integrations over q and t in Eq. (6.179) are performed analytically. The similar transformations are performed also in the 2D case. As a result, the memory function (6.177) is represented in the unified form for 3D and 2D interacting polarons:

$$
\begin{aligned}
\chi\left(\omega\right) = {} & \lim_{\varepsilon\to+0} \frac{2\alpha}{3\pi N} \left(\frac{3\pi}{4}\right)^{3-D} \left(\frac{\omega_{\mathrm{LO}}}{A}\right)^{3/2} \\
& \times \sum_{p_1=0}^{\infty}\sum_{p_2=0}^{\infty}\sum_{p_3=0}^{\infty} \frac{(-1)^{p_3}}{p_1!p_2!p_3!} \left(\frac{a_1^2}{N\Omega_1 A}\right)^{p_1} \left(\frac{a_2^2}{N\Omega_2 A}\right)^{p_2} \left(\frac{1}{NwA}\right)^{p_3} \\
& \times \Bigg\{ \Bigg[\sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\sum_{\sigma} \left[f_1\left(n,\sigma|\left\{N_\sigma\right\},\beta\right) - f_2\left(n,\sigma;m,\sigma|\left\{N_\sigma\right\},\beta\right)\right]\Big|_{\beta\to\infty} \\
& \times \left(\begin{array}{c} \frac{1}{\omega-\omega_{\mathrm{LO}}-[p_1\Omega_1+p_2\Omega_2+(p_3-m+n)w]+\mathrm{i}\varepsilon} - \frac{1}{\omega+\omega_{\mathrm{LO}}+p_1\Omega_1+p_2\Omega_2+(p_3-m+n)w+\mathrm{i}\varepsilon} \\ +\mathcal{P}\left(\frac{2}{\omega_{\mathrm{LO}}+p_1\Omega_1+p_2\Omega_2+(p_3-m+n)w}\right) \end{array} \right) \\
& \times \sum_{l=0}^{m} \sum_{k=n-m+l}^{n} \frac{(-1)^{n-m+l+k}\,\Gamma\left(p_1+p_2+p_3+k+l+\frac{3}{2}\right)}{k!l!} \left(\frac{1}{wA}\right)^{l+k} \\
& \times \binom{n+D-1}{n-k} \binom{2k}{k-l-n+m} \Bigg] \\
& + \Bigg[\left(\begin{array}{c} \frac{1}{\omega-\omega_{\mathrm{LO}}-(p_1\Omega_1+p_2\Omega_2+p_3 w)+\mathrm{i}\varepsilon} - \frac{1}{\omega+\omega_{\mathrm{LO}}+p_1\Omega_1+p_2\Omega_2+p_3 w+\mathrm{i}\varepsilon} \\ +\mathcal{P}\left(\frac{2}{\omega_{\mathrm{LO}}+p_1\Omega_1+p_2\Omega_2+p_3 w}\right) \end{array} \right) \\
& \times \sum_{m=0}^{\infty}\sum_{n=0}^{\infty}\sum_{\sigma,\sigma'} f_2\left(n,\sigma;m,\sigma'|\left\{N_\sigma\right\},\beta\right)\Big|_{\beta\to\infty} \\
& \times \sum_{k=0}^{n}\sum_{l=0}^{m} \frac{(-1)^{k+l}\,\Gamma\left(p_1+p_2+p_3+k+l+\frac{3}{2}\right)}{k!l!} \left(\frac{1}{wA}\right)^{k+l} \\
& \times \binom{n+D-1}{n-k} \binom{m+D-1}{m-l} \Bigg] \Bigg\}, \qquad (6.180)
\end{aligned}
$$

where $D = 2, 3$ is the dimensionality of the space, $\mathcal{P}$ denotes the principal value, A is defined as $A \equiv \left[\sum_{i=1}^{2} a_i^2/\Omega_i + (N-1)/w\right]/N$, Ω_1, Ω_2, and w are the eigenfrequencies of the model system, a_1 and a_2 are the coefficients of the canonical transformation which diagonalizes the model Lagrangian (6.32).

1. *Selected results: the manifestations of the shell filling in optical conductivity*

The shell filling schemes for an N-polaron system in a quantum dot can manifest themselves in the spectra of the optical conductivity. In Fig. 17, optical conductivity spectra for $N = 20$ polarons are presented for a quantum dot with the parameters of CdSe: $\alpha = 0.46$,

$\eta = 0.656$ [5] and with different values of the confinement energy $\hbar\Omega_0$. [4] In this case, the spin-polarized ground state changes to the ground state satisfying Hund's rule with increasing $\hbar\Omega_0$ in the interval $0.0421H^* < \hbar\Omega_0 < 0.0422H^*$.



FIG. 17: Optical conductivity spectra of $N = 20$ interacting polarons in CdSe quantum dots with $\alpha = 0.46$, $\eta = 0.656$ for different confinement energies close to the transition from a spin-polarized ground state to a ground state obeying Hund's rule. *Inset*: the first frequency moment $\langle\omega\rangle$ of the optical conductivity as a function of the confinement energy. (From Ref. [96].)

In the inset to Fig. 17, the first frequency moment of the optical conductivity

$$\langle\omega\rangle \equiv \frac{\int_0^\infty \omega \operatorname{Re}\sigma(\omega)\, d\omega}{\int_0^\infty \operatorname{Re}\sigma(\omega)\, d\omega}, \tag{6.181}$$

as a function of $\hbar\Omega_0$ shows a *discontinuity*, at the value of the confinement energy corresponding to the change of the shell filling schemes from the spin-polarized ground state to the ground state obeying Hund's rule. This discontinuity might be observable in optical measurements.

---

[4] For the numerical calculations, we use effective atomic units, where $\hbar$, the electron band mass m_b and $e/\sqrt{\varepsilon_\infty}$ have the numerical value of 1. This means that the unit of length is the effective Bohr radius $a_B^* = \hbar^2\varepsilon_\infty/\left(m_b e^2\right)$, while the unit of energy is the effective Hartree $H^* = m_b e^4/\left(\hbar^2\varepsilon_\infty^2\right)$.

The shell structure for a system of interacting polarons in a quantum dot is clearly revealed when analyzing the addition energy and the first frequency moment of the optical conductivity in parallel. In Fig 18, we show both the function

$$\Theta\left(N\right) \equiv \left.\left\langle\omega\right\rangle\right|_{N+1} - 2\left.\left\langle\omega\right\rangle\right|_{N} + \left.\left\langle\omega\right\rangle\right|_{N-1}, \tag{6.182}$$

and the addition energy

$$\Delta\left(N\right) = E^{0}\left(N+1\right) - 2E^{0}\left(N\right) + E^{0}\left(N-1\right). \tag{6.183}$$

for interacting polarons in a 3D CdSe quantum dot.



FIG. 18: The function $\Theta\left(N\right)$ and the addition energy $\Delta\left(N\right)$ for systems of interacting polarons in CdSe quantum dots with $\alpha = 0.46$, $\eta = 0.656$ for $\hbar\Omega_0 = 0.1H^*$. (From Ref. [96].)

As seen from Fig 18, distinct peaks appear in $\Theta\left(N\right)$ and $\Delta\left(N\right)$ at the “magic numbers” corresponding to closed-shell configurations at $N = 8, 20$ and to half-filled-shell configurations at $N = 5, 14$. We see that each of the peaks of $\Theta\left(N\right)$ corresponds to a peak of the addition energy. The filling patterns for a many-polaron system in a quantum dot can be therefore determined from the analysis of the first moment of the optical absorption for different numbers of polarons.

VII. VARIATIONAL PATH-INTEGRAL TREATMENT OF A TRANSLATION INVARIANT N-POLARON SYSTEM

A. The many-polaron system

In the present section, the ground-state properties of a translation invariant N-polaron system are theoretically studied in the framework of the variational path-integral method for identical particles, using a further development [103] of the model introduced in Refs. [96, 98, 104].

In order to describe a many-polaron system, we start from the translation invariant N-polaron Hamiltonian

$$H=\sum_{j=1}^{N}\frac{\mathbf{p}_j^2}{2m}+\frac{1}{2}\sum_{j=1}^{N}\sum_{l=1,\neq j}^{N}\frac{e^2}{\epsilon_\infty\left|\mathbf{r}_j-\mathbf{r}_l\right|}+\sum_{\mathbf{k}}\hbar\omega_{\mathrm{LO}}a_{\mathbf{k}}^{\dagger}a_{\mathbf{k}}+\left(\sum_{j=1}^{N}\sum_{\mathbf{k}}V_k a_{\mathbf{k}}e^{i\mathbf{k}\cdot\mathbf{r}_j}+H.c.\right), \quad (7.1)$$

where m is the band mass, e is the electron charge, ω_{LO} is the longitudinal optical (LO) phonon frequency, and V_k are the amplitudes of the Fröhlich electron-LO-phonon interaction

$$V_k=i\frac{\hbar\omega_{\mathrm{LO}}}{k}\left(\frac{4\pi\alpha}{V}\right)^{1/2}\left(\frac{\hbar}{2m\omega_{\mathrm{LO}}}\right)^{1/4}, \quad \alpha=\frac{e^2}{2\hbar\omega_{\mathrm{LO}}}\left(\frac{2m\omega_{\mathrm{LO}}}{\hbar}\right)^{1/2}\left(\frac{1}{\epsilon_\infty}-\frac{1}{\epsilon_0}\right), \quad (7.2)$$

with the electron-phonon coupling constant $\alpha>0$, the high-frequency dielectric constant $\epsilon_\infty>0$ and the static dielectric constant $\epsilon_0>0$, and consequently

$$\frac{e^2}{\epsilon_\infty}>\hbar\left(\frac{2\hbar\omega_{\mathrm{LO}}}{m}\right)^{1/2}\alpha\Longleftrightarrow\alpha\sqrt{2}<\left(\frac{H^*}{\hbar\omega_{\mathrm{LO}}}\right)^{1/2}\equiv U. \quad (7.3)$$

In the expression (7.3), H^* is the effective Hartree

$$H^*=\frac{e^2}{\epsilon_\infty a_B^*}, \quad a_B^*=\frac{\hbar^2}{me^2/\epsilon_\infty} \quad (7.4)$$

where a_B^* is the effective Bohr radius. The partition function of the system can be expressed as a path integral over all electron and phonon coordinates. The path integral over the phonon variables can be calculated analytically [41]. Feynman's phonon elimination technique for this system is well known and leads to the partition function, which is a path integral over the electron coordinates only:

$$Z=\left(\prod_{\mathbf{k}}\frac{e^{\frac{1}{2}\beta\hbar\omega_{\mathrm{LO}}}}{2\sinh\frac{1}{2}\beta\hbar\omega_{\mathrm{LO}}}\right)\oint e^{S}\mathcal{D}\bar{\mathbf{r}} \quad (7.5)$$

where $\bar{\mathbf{r}} = \{\mathbf{r}_1, \cdots, \mathbf{r}_N\}$ denotes the set of electron coordinates, and $\oint \mathcal{D}\bar{\mathbf{r}}$ denotes the path integral over all the electron coordinates, integrated over equal initial and final points, i.e.

$$\oint e^S \mathcal{D}\bar{\mathbf{r}} \equiv \int d\bar{\mathbf{r}} \int_{\bar{\mathbf{r}}(0)=\bar{\mathbf{r}}}^{\bar{\mathbf{r}}(\beta)=\bar{\mathbf{r}}} e^S \mathcal{D}\bar{\mathbf{r}}\left(\tau\right).$$

Throughout this paper, imaginary time variables are used. The effective action for the N-polaron system is retarded and given by

$$S = -\int_0^\beta \left(\frac{m}{2} \sum_{j=1}^{N} \left(\frac{d\mathbf{r}_j\left(\tau\right)}{d\tau} \right)^2 + \frac{1}{2} \sum_{j=1}^{N} \sum_{l=1,\neq j}^{N} \frac{e^2}{\epsilon_\infty \left| \mathbf{r}_j\left(\tau\right) - \mathbf{r}_l\left(\tau\right) \right|} \right) d\tau$$
$$+ \frac{1}{2} \int_0^\beta \int_0^\beta \sum_{j,l=1}^{N} \sum_{\mathbf{k}} \left| V_k \right|^2 e^{i\mathbf{k}\cdot\left(\mathbf{r}_j(\tau) - \mathbf{r}_l(\sigma)\right)} \frac{\cosh \hbar\omega_{\mathrm{LO}} \left(\frac{1}{2}\beta - \left| \tau - \sigma \right| \right)}{\sinh \frac{1}{2}\beta\hbar\omega_{\mathrm{LO}}} d\sigma d\tau. \tag{7.6}$$

Note that the electrons are fermions. Therefore the path integral for the electrons with parallel spin has to be interpreted as the required antisymmetric projection of the propagators for distinguishable particles.

We below use units in which $\hbar = 1$, $m = 1$, and $\omega_{\mathrm{LO}} = 1$. The units of distance and energy are thus the effective polaron radius $\left[\hbar / \left(m\omega_{\mathrm{LO}}\right)\right]^{1/2}$ and the LO-phonon energy $\hbar\omega_{\mathrm{LO}}$.

B. Variational principle

For distinguishable particles, it is well known that the Jensen-Feynman inequality [41] provides a lower bound on the partition function Z (and consequently an upper bound on the free energy F)

$$Z = \oint e^S \mathcal{D}\bar{\mathbf{r}} = \left(\oint e^{S_0} \mathcal{D}\bar{\mathbf{r}} \right) \left\langle e^{S-S_0} \right\rangle_0 \geq \left(\oint e^{S_0} \mathcal{D}\bar{\mathbf{r}} \right) e^{\left\langle S-S_0 \right\rangle_0} \text{ with } \left\langle A \right\rangle_0 \equiv \frac{\oint A\left(\bar{\mathbf{r}}\right) e^{S_0} \mathcal{D}\bar{\mathbf{r}}}{\oint e^{S_0} \mathcal{D}\bar{\mathbf{r}}}, \tag{7.7}$$

$$e^{-\beta F} \geq e^{-\beta F_0} e^{\left\langle S-S_0 \right\rangle_0} \Longrightarrow F \leq F_0 - \frac{\left\langle S-S_0 \right\rangle_0}{\beta} \tag{7.8}$$

for a system with real action S and a real trial action S_0.The many-body extension (Ref. [97, 105]) of the Jensen-Feynman inequality, requires that the potentials are symmetric with respect to all particle permutations, and that the exact propagator as well as the model propagator are defined on the same state space. Within this interpretation we consider the following generalization of Feynman's trial action

$$\begin{aligned} S_0 = &-\int_0^\beta \left(\frac{1}{2}\sum_{j=1}^{N}\left(\frac{d\mathbf{r}_j(\tau)}{d\tau}\right)^2 + \frac{\omega^2+w^2-v^2}{4N}\sum_{j,l=1}^{N}\left(\mathbf{r}_j(\tau)-\mathbf{r}_l(\tau)\right)^2\right)d\tau \\ &-\frac{w}{8}\frac{v^2-w^2}{N}\sum_{j,l=1}^{N}\int_0^\beta\int_0^\beta \left(\mathbf{r}_j(\tau)-\mathbf{r}_l(\sigma)\right)^2\frac{\cosh w\left(\frac{1}{2}\beta-|\tau-\sigma|\right)}{\sinh\frac{1}{2}\beta w}d\sigma d\tau \end{aligned} \tag{7.9}$$

with the variational frequency parameters v, w, ω.

Using the explicit forms of the exact (7.6) and the trial (7.9) actions, the variational inequality (7.8) takes the form

$$\begin{aligned} F(\beta|N_\uparrow,N_\downarrow) \le\ & F_0(\beta|N_\uparrow,N_\downarrow) + \frac{U}{2\beta}\int_0^\beta\left\langle\sum_{j,l=1,\neq j}^{N}\frac{1}{|\mathbf{r}_j(\tau)-\mathbf{r}_l(\tau)|}\right\rangle_0 d\tau \\ &-\frac{\omega^2+w^2-v^2}{4N\beta}\int_0^\beta\left\langle\sum_{j,l=1}^{N}\left(\mathbf{r}_j(\tau)-\mathbf{r}_l(\tau)\right)^2\right\rangle_0 d\tau \\ &-\frac{w}{8}\frac{v^2-w^2}{N\beta}\int_0^\beta\int_0^\beta\left\langle\sum_{j,l=1}^{N}\left(\mathbf{r}_j(\tau)-\mathbf{r}_l(\sigma)\right)^2\right\rangle_0\frac{\cosh w\left(\frac{1}{2}\beta-|\tau-\sigma|\right)}{\sinh\frac{1}{2}\beta w}d\sigma d\tau \\ &-\frac{1}{2\beta}\int_0^\beta\int_0^\beta\sum_{\mathbf{k}}|V_k|^2\left\langle\sum_{j,l=1}^{N}e^{i\mathbf{k}\cdot(\mathbf{r}_j(\tau)-\mathbf{r}_l(\sigma))}\right\rangle_0\frac{\cosh\omega_{\mathrm{LO}}\left(\frac{1}{2}\beta-|\tau-\sigma|\right)}{\sinh\frac{1}{2}\beta\omega_{\mathrm{LO}}}d\sigma d\tau. \end{aligned} \tag{7.10}$$

In the zero-temperature limit ($\beta\to\infty$), we arrive at the following upper bound for the ground-state energy $E^0(N_\uparrow,N_\downarrow)$ of a translation invariant N-polaron system

$$E^0(N_\uparrow,N_\downarrow) \le E_{var}(N_\uparrow,N_\downarrow|v,w,\omega),$$

with

$$\begin{aligned} E_{var}(N_\uparrow,N_\downarrow|v,w,\omega) = &\frac{3}{4}\frac{(v-w)^2}{v}-\frac{3}{4}\omega+\frac{1}{2}\mathbb{E}_F(N_\downarrow)+\frac{1}{2}\mathbb{E}_F(N_\downarrow) \\ &+E_{C\|}(N_\uparrow)+E_{C\|}(N_\downarrow)+E_{C\uparrow\downarrow}(N_\uparrow,N_\downarrow) \\ &+E_{\alpha\|}(N_\uparrow)+E_{\alpha\|}(N_\downarrow)+E_{\alpha\uparrow\downarrow}(N_\uparrow,N_\downarrow), \end{aligned} \tag{7.11}$$

where $\mathbb{E}_F(N)$ is the energy of N spin-polarized fermions confined to a parabolic potential with the confinement frequency ω, $E_{C\|}\left(N_{\uparrow(\downarrow)}\right)$ is the Coulomb energy of the electrons with parallel spins, $E_{C\uparrow\downarrow}(N_\uparrow,N_\downarrow)$ is the Coulomb energy of the electrons with opposite spins, $E_{\alpha\|}\left(N_{\uparrow(\downarrow)}\right)$ is the electron-phonon energy of the electrons with parallel spins, and $E_{\alpha\uparrow\downarrow}(N_\uparrow,N_\downarrow)$ is the electron-phonon energy of the electrons with opposite spins.

C. Results

Here, we discuss some results of the numerical minimization of $E_{var}\left(N_\uparrow, N_\downarrow | v, w, \omega\right)$ with respect to the three variational parameters v, w, and ω. The Fröhlich constant α and the Coulomb parameter

$$\alpha_0 \equiv \frac{U}{\sqrt{2}} \equiv \frac{\alpha}{1-\eta} \text{ with } \frac{1}{\eta} = \frac{\varepsilon_0}{\varepsilon_\infty} \tag{7.12}$$

characterize the strength of the electron-phonon and of the Coulomb interaction, obeying the physical condition $\alpha \geq \alpha_0$ [see (7.3)]. The optimal values of the variational parameters v,w,and ω are denoted v_{op},w_{op},and ω_{op}, respectively. The optimal value of the total spin was always determined by choosing the combination $(N_\uparrow, N_\downarrow)$ for fixed $N = N_\uparrow + N_\downarrow$which corresponds to the lowest value $E^0(N)$ of the variational functional

$$E^0(N) \equiv \min_{N_\uparrow} E_{var}\left(N_\uparrow, N - N_\uparrow | v_{op}, w_{op}, \omega_{op}\right). \tag{7.13}$$



FIG. 19: The "phase diagrams" of a translation invariant N-polaron system. The grey area is the non-physical region, for which $\alpha > \alpha_0$. The stability region for each number of electrons is determined by the equation $\alpha_c < \alpha < \alpha_0$. (From Ref. [103].)

In Fig. 19, the "phase diagrams" analogous to the bipolaron "phase diagram" of Ref. [106] are plotted for an N-polaron system in bulk with $N = 2, 3, 5$, and 10. The area where

$\alpha_0 \leq \alpha$ is the non-physical region. For $\alpha > \alpha_0$, each sector between a curve corresponding to a well defined N and the line indicating $\alpha_0 = \alpha$ shows the stability region where $\omega_{op} \neq 0$, while the white area corresponds to the regime with $\omega_{op} = 0$. When comparing the stability region for $N = 2$ from Fig. 19 with the bipolaron "phase diagram" of Ref. [106], the stability region in the present work starts from the value $\alpha_c \approx 4.1$ (instead of $\alpha_c \approx 6.9$ in Ref. [106]). The width of the stability region within the present model is also larger than the width of the stability region within the model of Ref. [106]. Also, the absolute values of the ground-state energy of a two-polaron system given by the present model are smaller than those given by the approach of Ref. [106].

The difference between the numerical results of the present work and of Ref. [106] is due to the following distinction between the used model systems. The model system of Ref. [106] consists of two electrons interacting with two fictitious particles and with each other through quadratic interactions. But the trial Hamiltonian given by Eq. (6) of Ref. [106] is not symmetric with respect to the permutation of the electrons. It is only symmetric under the permutation of the pairs "electron + fictitious particle". As a consequence, this trial system is only applicable if the electrons are distinguishable, i.e. have opposite spin. In contrast to the model of Ref. [106], the model used in the present paper is described by the trial action (9), which is fully symmetric with respect to the permutations of the electrons, as is required to describe identical particles.

The "phase diagrams" for $N > 2$ demonstrate the existence of stable multipolaron states (see Ref. [107]). As distinct from Ref. [107], here the ground state of an N-polaron system is investigated supposing that the electrons are fermions. As seen from Fig. 20, for $N > 2$, the stability region for a multipolaron state is narrower than the stability region for $N = 2$, and its width decreases with increasing N.

A consequence of the Fermi statistics is the dependence of the polaron characteristics and of the total spin of an N-polaron system on the parameters (α, α_0,N). In Fig. 20, we present the ground-state energy per particle, the confinement frequency ω_{op} and the total spin S as a function of the coupling constant α for $\alpha_0/\alpha = 1.05$ and for a different numbers of polarons. The ground-state energy turns out to be a continuous function of α, while $\omega_{op}$and S reveal jumps. For $N = 2$ (the case of a bipolaron), we see from Fig. 20 that the ground state has a total spin $S = 0$ for all values of α, i. e., the ground state of a bipolaron is a singlet. This result is in agreement with earlier investigations on the large-bipolaron



FIG. 20: The ground-state energy per particle (a), the optimal value ω_{op} of the confinement frequency (b), and the total spin (c) of a translation invariant N-polaron system as a function of the coupling strength α for $\alpha_0/\alpha = 0.5$. The vertical dashed lines in the panel c indicate the critical values α_c separating the regimes of $\alpha > \alpha_c$, where the multipolaron ground state with $\omega_{op} \neq 0$ exists, and $\alpha < \alpha_c$, where $\omega_{op} = 0$. (From Ref. [103].)

problem (see, e. g., [108]).

In summary, using the extension of the Jensen-Feynman variational principle to the systems of identical particles, we have derived a rigorous upper bound for the free energy of a translation invariant system of N interacting polarons. The developed approach is valid for an arbitrary coupling strength. The resulting ground-state energy is obtained taking into account the Fermi statistics of electrons.

VIII. RIPPLONIC POLARONS IN MULTIELECTRON BUBBLES

A. Ripplon-phonon modes of a MEB

Spherical shells of charged particles appear in a variety of physical systems, such as fullerenes, metallic nanoshells, charged droplets and neutron stars. A particularly interesting physical realization of the spherical electron gas is found in multielectron bubbles (MEBs) in liquid helium-4. These MEBs are 0.1 μm – 100 μm sized cavities inside liquid helium, that contain helium vapor at vapor pressure and a nanometer-thick electron layer anchored to the surface of the bubble [110]. They exist as a result of equilibrium between the surface tension of liquid helium and the Coulomb repulsion of the electrons [111]. Recently proposed experimental schemes to stabilize MEBs [112] have stimulated theoretical investigation of their properties.

We describe the dynamical modes of an MEB by considering the motion of the helium surface ("ripplons") and the vibrational modes of the electrons together. In particular, we analyze the case when the ripplopolarons form a Wigner lattice [109].

First, we derive the Lagrangian of interacting ripplons and phonons within a continuum approach. The shape of the surface of a bubble is described by the function $R(\theta,\varphi) = R_\mathrm{b} + u(\theta,\varphi)$, where $u(\theta,\varphi)$ is the deformation of the surface from a sphere with radius R_b. The deformation can be expanded in a series of spherical harmonics $Y_{lm}(\theta,\varphi)$ with amplitudes Q_{lm},

$$u(\theta,\varphi) = \sum_{l=1}^{\infty}\sum_{m=-l}^{l} Q_{lm} Y_{lm}(\theta,\varphi). \tag{8.1}$$

We suppose that the amplitudes are small in such a way that $\sqrt{l(l+1)}\,|Q_{lm}| \ll R_\mathrm{b}$.

The ripplon contribution (T_r) to the kinetic energy of an MEB, and the contributions to the potential energy due to the surface tension (U_σ) and due to the pressure (U_V) were described in Ref. [113]:

$$\begin{aligned} T_\mathrm{r} &= \frac{\rho}{2} R_\mathrm{b}^3 \sum_{l=1}^{\infty}\sum_{m=-l}^{l} \frac{1}{l+1} \left|\dot{Q}_{lm}\right|^2, \\ U_\sigma &= 4\pi\sigma R_\mathrm{b}^2 + \frac{\sigma}{2}\sum_{l=1}^{\infty}\sum_{m=-l}^{l} (l^2+l+2)\,|Q_{lm}|^2, \\ U_\mathrm{V} &= \frac{4\pi}{3} p R_\mathrm{b}^3 + p R_\mathrm{b}\sum_{l=1}^{\infty}\sum_{m=-l}^{l} |Q_{lm}|^2. \end{aligned} \tag{8.2}$$

Here $\rho \approx 145$ kg/m^3 is the density of liquid helium, $\sigma \approx 3.6 \times 10^{-4}$ J/m^2 is its surface tension, and p is the difference of pressures outside and inside the bubble.

Expanding the surface electron density $n\left(\theta, \varphi\right)$ in a series of spherical harmonics with amplitudes n_{lm},

$$n\left(\theta,\varphi\right)=\sum_{l=0}^{\infty}\sum_{m=-l}^{l} n_{lm}Y_{lm}\left(\theta,\varphi\right), \tag{8.3}$$

the kinetic energy of the motion of electrons can be written as

$$T_{\mathrm{p}}=\frac{1}{2}\sum_{l=1}^{\infty}\sum_{m=-l}^{l}\frac{4\pi m_{\mathrm{e}}R_{\mathrm{b}}^{6}}{l(l+1)N}\left|\dot{n}_{lm}\right|^{2}, \tag{8.4}$$

where m_{e} is the bare electron mass and N is the number of electrons. Finally, the electrostatic energy (U_{C}) of the deformed MEB with a non-uniform surface electron density (8.3) is calculated using the Maxwell equations and the electrostatic boundary conditions at the surface. The result is:

$$\begin{aligned} U_{\mathrm{C}}= & \frac{e^{2}N^{2}}{2\varepsilon R_{\mathrm{b}}}+2\pi e^{2}R_{\mathrm{b}}^{3}\sum_{l=1}^{\infty}\sum_{m=-l}^{l}\frac{\left|n_{lm}\right|^{2}}{l+\varepsilon\left(l+1\right)} \\ & -\frac{e^{2}N^{2}}{8\pi\varepsilon R_{\mathrm{b}}^{3}}\sum_{l=1}^{\infty}\sum_{m=-l}^{l}\frac{l^{2}-\varepsilon\left(l+1\right)}{l+\varepsilon\left(l+1\right)}\left|Q_{lm}\right|^{2} \\ & -e^{2}N\sum_{l=1}^{\infty}\sum_{m=-l}^{l}\frac{l+1}{l+\varepsilon\left(l+1\right)}n_{lm}Q_{lm}^{*}, \end{aligned} \tag{8.5}$$

with the dielectric constant of liquid helium $\varepsilon \approx 1.0572$. The last term in Eq. (8.5) describes the ripplon-phonon mixing. Only ripplon and phonon modes which have the same angular momentum couple to each other. After the diagonalization of the Lagrangian of this ripplon-phonon system, we arrive at the eigenfrequencies:

$$\begin{aligned} \Omega_{1,2}\left(l\right)= & \left\{\frac{1}{2}\left[\omega_{\mathrm{p}}^{2}\left(l\right)+\omega_{\mathrm{r}}^{2}\left(l\right)\right.\right. \\ & \left.\left.\pm\sqrt{\left[\omega_{\mathrm{p}}^{2}\left(l\right)-\omega_{\mathrm{r}}^{2}\left(l\right)\right]^{2}+4\gamma^{2}\left(l\right)}\right]\right\}^{1/2}, \end{aligned} \tag{8.6}$$

where $\omega_{\mathrm{r}}\left(l\right)$ is the bare ripplon frequency,

$$\begin{aligned} \omega_{\mathrm{r}}\left(l\right)= & \left\{\frac{l+1}{\rho R_{\mathrm{b}}^{3}}\left[\sigma\left(l^{2}+l+2\right)\right.\right. \\ & \left.\left.-\frac{e^{2}N^{2}}{4\pi\varepsilon R_{\mathrm{b}}^{3}}\frac{l^{2}-\varepsilon\left(l+1\right)}{l+\varepsilon\left(l+1\right)}+2pR_{\mathrm{b}}\right]\right\}^{1/2}, \end{aligned} \tag{8.7}$$

while $\omega_{\mathrm{p}}(l)$ is the bare phonon frequency,

$$\omega_{\mathrm{p}}(l)=\left(\frac{e^{2}N}{m_{\mathrm{e}}R_{\mathrm{b}}^{3}}\frac{l(l+1)}{l+\varepsilon(l+1)}\right)^{1/2}, \tag{8.8}$$

and $\gamma(l)$ describes the ripplon-phonon coupling:

$$\gamma(l)=\frac{e^{2}N}{R_{\mathrm{b}}^{3}}\left(\frac{Nl}{4\pi m_{\mathrm{e}}\rho R_{\mathrm{b}}^{3}}\right)^{1/2}\frac{(l+1)^{2}}{l+\varepsilon(l+1)}. \tag{8.9}$$

B. Electron-ripplon interaction in the MEB

The interaction energy between the ripplons and the electrons in the multielectron bubble can be derived from the following considerations: (i) the distance between the layer electrons and the helium surface is fixed (the electrons find themselves confined to an effectively 2D surface anchored to the helium surface) and (ii) the electrons are subjected to a force field, arising from the electric field of the other electrons. For a spherical bubble, this electric field lies along the radial direction and equals

$$\mathbf{E}=-\frac{Ne}{2R_{b}^{2}}\mathbf{e_{r}}. \tag{8.10}$$

A bubble shape oscillation will displace the layer of electrons anchored to the surface. The interaction energy which arises from this, equals the displacement of the electrons times the force $e\mathbf{E}$ acting on them. Thus, we get for the interaction Hamiltonian

$$\hat{H}_{int}=\sum_{j}e|\mathbf{E}|\times u(\hat{\Omega}_{j}). \tag{8.11}$$

Here $u(\Omega)$ is the radial displacement of the surface in the direction given by the spherical angle Ω; and $\hat{\Omega}_{j}$ is the (angular) position operator for electron j. The displacement can be rewritten using (8.1) and we find

$$\hat{H}_{int}=\sum_{j}e|\mathbf{E}|\sum_{\ell,m}\hat{Q}_{\ell m}Y_{\ell m}(\hat{\Omega}_{j}). \tag{8.12}$$

Using the relation

$$\hat{Q}_{\ell,m}=(-1)^{(m-|m|)/2}\sqrt{\tfrac{\hbar(\ell+1)}{2\rho R_{b}^{3}\omega_{\ell}}}(\hat{a}_{\ell,m}+\hat{a}_{\ell,-m}^{+}), \tag{8.13}$$

the interaction Hamiltonian can be written in the suggestive form

$$\hat{H}_{int}=\sum_{\ell,m}\sum_{j}M_{\ell,m}Y_{\ell,m}(\hat{\Omega}_{j})(\hat{a}_{\ell,m}+\hat{a}_{\ell,-m}^{+}), \tag{8.14}$$

with the electron-ripplon coupling amplitude for a MEB given by

$$M_{\ell,m} = (-1)^{(m-|m|)/2} \frac{Ne^2}{2R_b^2} \sqrt{\frac{\hbar(\ell+1)}{2\rho R_b^3 \omega_\ell}} \tag{8.15}$$

C. Locally flat approximation

Substituting $M_{\ell,m}$ into (8.14), we get

$$\hat{H}_{int} = \sum_{\ell,m} \sum_j \frac{Ne^2}{2R_b^2} \sqrt{\frac{\hbar(\ell+1)}{2\rho R_b^3 \omega_\ell}} \tag{8.16}$$
$$\times \left[(-1)^{(m-|m|)/2} \frac{Y_{\ell,m}(\hat{\Omega}_j)}{R_b} \right] (\hat{a}_{\ell,m} + \hat{a}^+_{\ell,-m}).$$

In this expression, we consider the limit of a bubble so large that the surface becomes flat on all length scales of interest. Hence we let $R_b \to \infty$ but keep $\ell/R_b = q$ a constant. This means we have to let $\ell \to \infty$ as well. In this limit,

$$\lim_{\ell\to\infty} Y_{\ell,0}(\theta) = \frac{i^\ell}{\pi\sqrt{\sin\theta}} \sin[(\ell+1/2)\theta + \pi/4], \tag{8.17}$$

and $Y_{\ell,0}(\theta)$ varies locally as a plane wave with wave vector $q = \ell/R_b$. The wave function $Y_{\ell,m}(\hat{\Omega}_j)/R_b$ is furthermore normalized with respect to integration over the surface (with total area $4\pi R_b^2$). Thus, we get in the locally flat approximation

$$\hat{H}_{int} = \sum_{\mathbf{q}} \sum_j \frac{Ne^2}{2R_b^2} \sqrt{\frac{\hbar q}{2\rho\omega(q)}} e^{i\mathbf{q}.\hat{\mathbf{r}}_j} (\hat{a}_{\mathbf{q}} + \hat{a}^+_{-\mathbf{q}}), \tag{8.18}$$

or

$$\hat{H}_{int} = \sum_{\mathbf{q}} \sum_j M_q e^{i\mathbf{q}.\hat{\mathbf{r}}_j} (\hat{a}_{\mathbf{q}} + \hat{a}^+_{-\mathbf{q}}),$$
$$M_q = e|\mathbf{E}| \sqrt{\frac{\hbar q}{2\rho\omega(q)}}. \tag{8.19}$$

This corresponds in the limit of large bubbles to the interaction Hamiltonian expected for a flat surface.

D. Ripplopolaron in a Wigner lattice: the mean-field approach

In their treatment of the electron Wigner lattice embedded in a polarizable medium such as a semiconductors or an ionic solid, Fratini and Quémerais [114] described the effect of the electrons on a particular electron through a mean-field lattice potential. The (classical) lattice potential V_{lat} is obtained by approximating all the electrons acting on one particular electron by a homogenous charge density in which a hole is punched out; this hole is centered in the lattice point of the particular electron under investigation and has a radius given by the lattice distance d.

Within this particular mean-field approximation, the lattice potential can be calculated from classical electrostatics and we find that for a 2D electron gas it can be expressed in terms of the elliptic functions of first and second kind, $E\left(x\right)$ and $K\left(x\right)$,

$$\begin{aligned} V_{lat}\left(\mathbf{r}\right) = -\frac{2e^2}{\pi d^2}\Bigg\{ & |d-r|\, E\left[-\frac{4rd}{\left(d-r\right)^2}\right] \\ & + \left(d+r\right)\operatorname{sgn}\left(d-r\right) K\left[-\frac{4rd}{\left(d-r\right)^2}\right]\Bigg\}. \end{aligned} \tag{8.20}$$

Here, $\mathbf{r}$ is the position vector measured from the lattice position. We can expand this potential around the origin to find the small-amplitude oscillation frequency of the electron lattice:

$$\lim_{r\ll d} V_{lat}\left(\mathbf{r}\right) = -\frac{2e^2}{d} + \frac{1}{2} m_e \omega_{lat}^2 r^2 + \mathcal{O}\left(r^4\right), \tag{8.21}$$

with the confinement frequency

$$\omega_{lat} = \sqrt{\frac{e^2}{m_e d^3}}. \tag{8.22}$$

In the mean-field approximation, the Hamiltonian for a ripplopolaron in a lattice on a *locally flat* helium surface is given by

$$\begin{aligned} \hat{H} = & \frac{\hat{p}^2}{2m_e} + V_{lat}\left(\hat{\mathbf{r}}\right) + \sum_{\mathbf{q}} \hbar\omega(q) \hat{a}_{\mathbf{q}}^{+} \hat{a}_{\mathbf{q}} \\ & + \sum_{\mathbf{q}} M_q e^{-i\mathbf{q}\cdot\mathbf{r}} \left(\hat{a}_{\mathbf{q}} + \hat{a}_{-\mathbf{q}}^{+}\right), \end{aligned} \tag{8.23}$$

where $\hat{\mathbf{r}}$ is the electron position operator.

Now that the lattice potential has been introduced, we can move on and include effects of the bubble geometry. If we restrict our treatment to the case of large bubbles (with $N > 10^5$ electrons), then both the ripplopolaron radius and the inter-electron distance d are much smaller than the radius of the bubble R_b. This gives us ground to use the locally flat approximation using the auxiliary model of a ripplonic polaron in a planar system described by (8.23), but with a modified ripplon dispersion relation and an modified pressing field.

We find for the modified ripplon dispersion relation in the MEB:

$$\omega(q) = \sqrt{\frac{\sigma}{\rho}q^3 + \frac{p}{\rho R_b}q}, \tag{8.24}$$

where R_b is the equilibrium bubble radius which depends on the pressure and the number of electrons. The bubble radius is found by balancing the surface tension and the pressure with the Coulomb repulsion. The modified electron-ripplon interaction amplitude in an MEB is given by

$$M_{\mathbf{q}} = e|\mathbf{E}|\sqrt{\frac{\hbar q}{2\rho\omega(q)}}. \tag{8.25}$$

The effective electric pressing field pushing the electrons against the helium surface and determining the strength of the electron-ripplon interaction is

$$\mathbf{E} = -\frac{Ne}{2R_b^2}\mathbf{e_r}. \tag{8.26}$$

E. Ripplopolaron Wigner lattice at finite temperature

To study the ripplopolaron Wigner lattice at finite temperature and for any value of the electron-ripplon coupling, we use the variational path-integral approach [41]. This variational principle distinguishes itself from Rayleigh-Ritz variation in that it uses a trial action functional S_{trial} instead of a trial wave function.

The action functional of the system described by Hamiltonian (8.23), becomes, after elimination of the ripplon degrees of freedom,

$$\begin{aligned} S = -\frac{1}{\hbar}\int_0^{\hbar\beta} d\tau \left\{ \frac{m_e}{2}\dot{r}^2(\tau) + V_{lat}[r(\tau)] \right\} + \sum_{\mathbf{q}} |M_q|^2 \\ \times \int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\sigma G_{\omega(q)}(\tau - \sigma) e^{i\mathbf{q}\cdot[\mathbf{r}(\tau)-\mathbf{r}(\sigma)]}, \end{aligned} \tag{8.27}$$

with

$$G_\nu(\tau - \sigma) = \frac{\cosh[\nu(|\tau - \sigma| - \hbar\beta/2)]}{\sinh(\beta\hbar\nu/2)}. \tag{8.28}$$

In preparation of its customary use in the Jensen-Feynman inequality, the action functional (8.27) is written in imaginary time $t = i\tau$ with $\beta = 1/(k_B T)$ where T is the temperature.

We introduce a quadratic trial action of the form

$$S_{trial} = -\frac{1}{\hbar}\int_0^{\hbar\beta} d\tau \left[\frac{m_e}{2}\dot{r}^2(\tau) + \frac{m_e\Omega^2}{2}r^2(\tau)\right]$$
$$-\frac{Mw^2}{4\hbar}\int_0^{\hbar\beta} d\tau \int_0^{\hbar\beta} d\sigma G_w(\tau-\sigma)\mathbf{r}(\tau)\cdot\mathbf{r}(\sigma). \tag{8.29}$$

where M, w, and Ω are the variationally adjustable parameters. This trial action corresponds to the Lagrangian

$$\mathcal{L}_0 = \frac{m_e}{2}\dot{r}^2 + \frac{M}{2}\dot{R}^2 - \frac{\kappa}{2}r^2 - \frac{K}{2}(\mathbf{r}-\mathbf{R})^2, \tag{8.30}$$

from which the degrees of freedom associated with $\mathbf{R}$ have been integrated out. This Lagrangian can be interpreted as describing an electron with mass m_e at position $\mathbf{r}$, coupled through a spring with spring constant κ to its lattice site, and to which a fictitious mass M at position $\mathbf{R}$ has been attached with another spring, with spring constant K. The relation between the spring constants in (8.30) and the variational parameters w, Ω is given by

$$w = \sqrt{K/m_e}, \tag{8.31}$$
$$\Omega = \sqrt{(\kappa+K)/m_e}. \tag{8.32}$$

Based on the trial action S_{trial}, Feynman's variational method allows one to obtain an upper bound for the free energy F of the system (at temperature T) described by the action functional S by minimizing the following function:

$$F = F_0 - \frac{1}{\beta}\left\langle S - S_{trial}\right\rangle, \tag{8.33}$$

with respect to the variational parameters of the trial action. In this expression, F_0 is the free energy of the trial system characterized by the Lagrangian $\mathcal{L}_0$, $\beta = 1/(k_bT)$ is the inverse temperature, and the expectation value $\langle S - S_{trial}\rangle$ is to be taken with respect to the ground state of this trial system. The evaluation of expression (8.33) is straightforward though lengthy. We find

$$
\begin{aligned}
F = {} & \frac{2}{\beta}\ln\left[2\sinh\left(\frac{\beta\hbar\Omega_1}{2}\right)\right] + \frac{2}{\beta}\ln\left[2\sinh\left(\frac{\beta\hbar\Omega_2}{2}\right)\right] \\
& - \frac{2}{\beta}\ln\left[2\sinh\left(\frac{\beta\hbar w}{2}\right)\right] - \frac{\hbar}{2}\sum_{i=1}^{2} a_i^2\Omega_i\coth\left(\frac{\beta\hbar\Omega_i}{2}\right) \\
& - \frac{\sqrt{\pi}e^2}{D}e^{-d^2/(2D)}\left[I_0\left(\frac{d^2}{2D}\right) + I_1\left(\frac{d^2}{2D}\right)\right] \\
& - \frac{1}{2\pi\hbar\beta}\int_{1/R_b}^{\infty} dqq|M_q|^2\int_0^{\hbar\beta/2} d\tau\frac{\cosh[\omega(q)(\tau-\hbar\beta/2)]}{\sinh[\beta\hbar\omega(q)/2]} \\
& \times\exp\left[-\tfrac{\hbar q^2}{2m_e}\sum_{j=1}^{2} a_j^2\tfrac{\cosh(\hbar\Omega_j\beta/2)-\cosh[\hbar\Omega_j(\tau-\beta/2)]}{\Omega_j\sinh(\hbar\Omega_j\beta/2)}\right].
\end{aligned}
\tag{8.34}
$$

In this expression, I_0 and I_1 are Bessel functions of imaginary argument, and

$$D = \frac{\hbar}{m_e}\sum_{j=1}^{2}\frac{a_j^2}{\Omega_j}\coth\left(\hbar\Omega_j\beta/2\right), \tag{8.35}$$

$$a_1 = \sqrt{\frac{\Omega_1^2 - w^2}{\Omega_1^2 - \Omega_2^2}};\, a_2 = \sqrt{\frac{w^2 - \Omega_2^2}{\Omega_1^2 - \Omega_2^2}}. \tag{8.36}$$

Finally, Ω_1 and Ω_2 are the eigenfrequencies of the trial system, given by

$$\Omega_{1,2}^2 = \frac{1}{2}\left[\Omega^2 + w^2 \pm \sqrt{\left(\Omega^2 - w^2\right)^2 + 4K/(Mm_e)}\right]. \tag{8.37}$$

Optimal values of the variational parameters are determined by the numerical minimization of the variational functional F as given by expression (8.34).

F. Melting of the ripplopolaron Wigner lattice

The Lindemann melting criterion [115] states in general that a crystal lattice of objects (be it atoms, molecules, electrons, or ripplopolarons) will melt when the average motion of the objects around their lattice site is larger than a critical fraction δ_0 of the lattice parameter d. It would be a strenuous task to calculate from first principles the exact value of the critical fraction δ_0, but for the particular case of electrons on a helium surface, we can make use of an experimental determination. Grimes and Adams [116] found that the Wigner lattice melts when $\Gamma = 137 \pm 15$, where Γ is the ratio of potential energy to the kinetic energy per electron. At temperature T the average kinetic energy in a lattice potential V_{lat} is

$$E_{kin} = \frac{\hbar\omega_{lat}}{2}\coth\left(\frac{\hbar\omega_{lat}}{2k_BT}\right), \tag{8.38}$$

and the average distance that an electron moves out of the lattice site is determined by

$$\left\langle \mathbf{r}^2 \right\rangle = \frac{\hbar}{m_e \omega_{lat}} \coth\left(\frac{\hbar\omega_{lat}}{2k_B T}\right) = \frac{2E_{kin}}{m_e \omega_{lat}^2}. \tag{8.39}$$

From this we find that for the melting transition in Grimes and Adams' experiment [116], the critical fraction equals $\delta_0 \approx 0.13$. This estimate is in agreement with previous (empirical) estimates yielding $\delta_0 \approx 0.1$ [117], and we shall use it in the rest of this section.

Within the approach of Fratini and Quémerais [114], the Wigner lattice of (ripplo)polarons melts when at least one of the two following Lindemann criteria are met:

$$\delta_r = \frac{\sqrt{\langle \mathbf{R}_{cms}^2 \rangle}}{d} > \delta_0, \tag{8.40}$$

$$\delta_\rho = \frac{\sqrt{\langle \rho^2 \rangle}}{d} > \delta_0. \tag{8.41}$$

where ρ and $\mathbf{R}_{cms}$ are, respectively, the relative coordinate and the center of mass coordinate of the model system (8.30): if $\mathbf{r}$ is the electron coordinate and $\mathbf{R}$ is the position coordinate of the fictitious ripplon mass M, this is

$$\mathbf{R}_{cms} = \frac{m_e \mathbf{r} + M\mathbf{R}}{m_e + M}; \rho = \mathbf{r} - \mathbf{R}. \tag{8.42}$$

The appearance of two Lindemann criteria takes into account the composite nature of (ripplo)polarons. As follows from the physical sense of the coordinates ρ and $\mathbf{R}_{cms}$, the first criterion (8.40) is related to the melting of the ripplopolaron Wigner lattice towards a ripplopolaron liquid, where the ripplopolarons move as a whole, the electron together with its dimple. The second criterion (8.41) is related to the dissociation of ripplopolarons: the electrons shed their dimple.

The path-integral variational formalism allows us to calculate the expectation values $\langle \mathbf{R}_{cms}^2 \rangle$ and $\langle \rho^2 \rangle$ with respect to the ground state of the variationally optimal model system. We find

$$\begin{aligned} \left\langle \mathbf{R}_{cms}^2 \right\rangle = {} & \frac{\hbar w^4}{m_e \left[w^2(\Omega_1^2 + \Omega_2^2) - \Omega_1^2\Omega_2^2\right](\Omega_1^2 - \Omega_2^2)} \\ & \times \left[\Omega_2^4(\Omega_1^2 - w^2)\coth(\hbar\Omega_1\beta/2)/\Omega_1 \right. \\ & \left. + \Omega_1^4(w^2 - \Omega_2^2)\coth(\hbar\Omega_2\beta/2)/\Omega_2\right], \end{aligned} \tag{8.43}$$

$$
\begin{aligned}
\left\langle \rho^2 \right\rangle = & \frac{\hbar}{m_e\left(\Omega_1^2-\Omega_2^2\right)\left(\Omega_1^2-w^2\right)\left(w^2-\Omega_2^2\right)} \\
& \times\left[\Omega_1^3(w^2-\Omega_2^2)\coth\left(\hbar\Omega_1\beta/2\right)\right. \\
& \left.+\Omega_2^3(\Omega_1^2-w^2)\coth(\hbar\Omega_2\beta/2)\right]. \qquad (8.44)
\end{aligned}
$$

Numerical calculation shows that for ripplopolarons in an MEB the inequality $\Omega_1 \gg w$ is fulfilled ($w/\Omega_1 \approx 10^{-3}$ to 10^{-2}) so that the strong-coupling regime is realized. Owing to this inequality, we find from Eqs. (8.43),(8.44) that

$$
\left\langle \mathbf{R}_{cms}^2 \right\rangle \ll \left\langle \rho^2 \right\rangle. \qquad (8.45)
$$

So, the destruction of the ripplopolaron Wigner lattice in an MEB occurs through the dissociation of ripplopolarons, since the second criterion (8.41) will be fulfilled before the first (8.40). The results for the melting of the ripplopolaron Wigner lattice are summarized in the phase diagram shown in Fig. 21.



FIG. 21: The phase diagram for the spherical 2D layer of electrons in the MEB. Above a critical pressure, a ripplopolaron solid (a Wigner lattice of electrons with dimples in the helium surface underneath them) is formed. Below the critical pressure, the ripplopolaron solid melts into an electron liquid through dissociation of ripplopolarons. (From Ref. [109].)

For every value of N, pressure p and temperature T in an experimentally accessible range, this figure shows whether the ripplopolaron Wigner lattice is present (points above the surface) or molten (points below the surface). Below a critical pressure (on the order of 10^4 Pa) the ripplopolaron solid will melt into an electron liquid. This critical pressure is nearly independent of the number of electrons (except for the smallest bubbles) and is weakly temperature dependent, up to the helium critical temperature 5.2 K. This can be understood since the typical lattice potential well in which the ripplopolaron resides has frequencies of the order of THz or larger, which correspond to ~ 10 K.

The new phase that we predict, the ripplopolaron Wigner lattice, will not be present for electrons on a flat helium surface. At the values of the pressing field necessary to obtain a strong enough electron-ripplon coupling, the flat helium surface is no longer stable against long-wavelength deformations [118]. Multielectron bubbles, with their different ripplon dispersion and the presence of stabilizing factors such as the energy barrier against fissioning [119], allow for much larger electric fields pressing the electrons against the helium surface. The regime of N, p, T parameters suitable for the creation of a ripplopolaron Wigner lattice lies within the regime that would be achievable in recently proposed experiments aimed at stabilizing multielectron bubbles [112]. The ripplopolaron Wigner lattice and its melting transition might be detected by spectroscopic techniques [116, 120] probing for example the transverse phonon modes of the lattice [121].

---

[1] L. D. Landau, *Phys. Z. Sowjetunion* **3**, 664 (1933) [English translation in *Collected Papers*, Gordon and Breach, New York, 1965, pp. 67-68].

[2] J. T. Devreese, in *Lectures on the Physics of Highly Correlated Electron Systems VII,* edited A. Avella and F. Mancini Proceedings of the 7th Training Course in the Physics of Correlated Electron Systems&High-T_c Superconductors, Vietri sul Mare, Italy, October 14-16, 2002, AIP, Melville (2003), pp. 3 - 56.

[3] H. Fröhlich, *Adv. Phys.* **3**, 325 (1954).

[4] R. P. Feynman, R. B. Leiton and M. Sands, The Feynman Lectures on Physics (Addison-Wesley, 1972), Vol. II.

[5] E. Kartheuser, in *Polarons in Ionic Crystals and Polar Semiconductors* edited J. T. Devreese,

North-Holland, Amsterdam (1972), pp. 717 - 733.

[6] M. Grynberg, S. Huant, G. Martinez, J. Kossut, T. Wojtowicz, G. Karczewski, J. M. Shi, F. M. Peeters, J. T. Devreese, *Phys. Rev. B* **54**, 1467 (1996).

[7] The spectra of the infrared-active LO (and TO) phonons in α-Al_2O_3 (sapphire) contain six modes. The values of the LO and TO phonon frequencies and of the high-frequency dielectric constants $\varepsilon_{\infty\|} = 3.072$, $\varepsilon_{\infty\perp} = 3.077$ are taken from Ref. [8]. Using these parameters and the electron band mass $m_b = 0.25m_e$ as estimated in Ref. [9], the effective value of the electron-phonon coupling constant α in Al_2O_3

has been calculated as

$$\alpha = \sum_j \alpha_j \left\langle \left(\mathbf{e}_j \cdot \frac{\mathbf{k}}{k} \right)^2 \right\rangle ,$$

where $\mathbf{e}_j$ is the polarization vector of the j-th LO-phonon branch, $\mathbf{k}$ is the phonon wave vector, $\langle\rangle$ denote the angular averaging, and the coupling constants α_j for each branch are obtained using the method [10]. The resulting value of the polaron coupling constant is $\alpha \approx 1.25$

[8] M. Schubert, T. E. Tiwald, and C. M. Herzinger, *Phys. Rev. B* **61**, 8187 (2000).

[9] J. Shan, F. Wang, E. Knoesel, M. Bonn, and T. F. Heinz, *Phys. Rev. Lett.* **90**, 247401 (2003).

[10] S. N. Klimin, V. M. Fomin, and J. T. Devreese, *to be published*.

[11] J. W. Hodby, G. P. Russell, F. Peeters, J. T. Devreese, and D. M. Larsen, *Phys. Rev. Lett.* **58**, 1471 (1987).

[12] The spectra of the LO (and TO) phonons in α-SiO_2 contain ten modes. The values of the LO and TO phonon frequencies are taken from Refs. [13, 14]. Using these frequencies and the value $\varepsilon_\infty = 2.40$ from Ref. [15] for the high-frequency dielectric constant, the effective value of the electron-phonon coupling constant α in SiO_2 has been calculated using the method of Ref. [10] as indicated in Ref. [7]. We use the estimated value of the electron band mass $m_b = 0.5m_e$ as in Refs. [16–18]. The resulting value of the polaron coupling constant is $\alpha \approx 1.59$.

[13] F. Gervais and B. Piriou, *Phys. Rev. B* **11**, 3944 (1975).

[14] J. L. Duarte, J. A. Sanjurjo, and R. S. Katiyar, *Phys. Rev. B* **36**, 3368 (1987).

[15] S. T. Pantelides and W. A. Harrison, *Phys. Rev. B* **13**, 2667 (1976).

[16] M. V. Fischetti, D. J. DiMaria, L. Dori, J. Batey, E. Tierney, and J. Stasiak, *Phys. Rev. B*

35, 4404 (1987).

[17] D. Arnold, E. Cartier, and D. J. DiMaria, *Phys. Rev. B* **49**, 10278 (1994).

[18] P. Martin, S. Guizard, Ph. Daguzan, G. Petite, P. D'Oliveira, P. Meynadier, and M. Perdrix, *Phys. Rev. B* **55**, 5799 (1997).

[19] I. Biaggio, R. W. Hellwarth, and J. P. Partanen, *Phys. Rev. Lett.* **78**, 891 (1997) (for $m_b/m_\mathrm{e} = 2$).

[20] G. Verbist, F. M. Peeters, and J. T. Devreese, *Ferroelectrics*, **130**, 27 (1992) (for $m_b/m_\mathrm{e} = 2.6$).

[21] S. I. Pekar, *Issledovanija po Ekektronnoj Teorii Kristallov*, Gostekhizdat, Moskva, 1951 (in Russian) [German translation: *Untersuchungen über die Elektronentheorie der Kristalle*, Akademie Verlag, Berlin, 1951].

[22] G. C. Kuper and G. D. Whitfield (eds.), *Polarons and Excitons*, Oliver and Boyd, Edinburgh, 1963.

[23] J. Appel, in *Solid State Physics*, edited by F. Seitz, D. Turnbull, and H. Ehrenreich, Academic Press, New York, 1968, vol. 21, pp. 193-391.

[24] J. T. Devreese (ed.), *Polarons in Ionic Crystals and Polar Semiconductors*, North-Holland, Amsterdam, 1972.

[25] T. K. Mitra, A. Chatterjee, and S. Mukhopadhyay, *Phys. Rep.* **153**, 91 (1987).

[26] J. T. Devreese, in *Encyclopedia of Applied Physics*, edited by G. L. Trigg, VCH, Weinheim, 1996, vol. 14, pp. 383 - 413.

[27] A. S. Alexandrov and Sir Nevill Mott, *Polarons and Bipolarons*, World Scientific, Singapore, 1996.

[28] A. S. Mishchenko, N. V. Prokof'ev, A. Sakamoto, and B. V. Svistunov, *Phys. Rev. B* **62**, 6317 (2000).

[29] T. D. Lee, F. E. Low, and D. Pines, Phys. Rev. **90**, 297 (1953).

[30] J. Röseler, Phys. Stat. Sol.(b) **25**, 311 (1968)

[31] M. A. Smondyrev, Teor. Math. Fiz. **68**, 29 (1986) [English translation: Theor. Math. Phys. **68**, 653 (1986)]

[32] O. V. Selyugin and M. A. Smondyrev, Phys. Stat. Sol.(b)

[33] L. D. Landau and S. I. Pekar, Zh. Eksper. Teor. Fiz. **18**, 419 (1948)

[34] N. N. Bogolubov and S. V. Tyablikov, Zh. Eksp. i Teor. Fiz. **19**, 256 (1949)

[35] N. N. Bogolubov, Ukr. Matem. Zh. **2**, 3 (1950)

[36] S. V. Tyablikov, Zh. Eksp. i Theor. Phys. **21**, 377 (1951)

[37] R. Evrard, Phys. Letters **14**, 295 (1965)

[38] S. J. Miyake, J. Phys. Soc. Japan **38**, 181 (1975)

[39] J. T. Devreese and R. Evrard, Phys. Letters **11**, 278 (1964).

[40] E. Kartheuser, R. Evrard, and J. Devreese, in *Optical Properties of Solids*, edited by E. D. Haidemenakis, Gordon and Breach, New York, 1970, pp. 433-459 (Table 1).

[41] R. P. Feynman, *Phys. Rev.* **97** 660 (1955).

[42] J. T. Devreese and R. Evrard, in *Proceedings of the British Ceramic Society* **10**, 151 (1968).

[43] J. T. Titantah, C. Pierleoni, and S. Ciuchi, *Phys. Rev. Lett.* **87**, 206406 (2001).

[44] G. De Filippis, V. Cataudella, V. Marigliano Ramaglia, C. A. Perroni, and D. Bercioux, *Eur. Phys. J. B* **36**, 65 (2003).

[45] R. P. Feynman, R. W. Hellwarth, C. K. Iddings, and P. M. Platzman, *Phys. Rev.* **127**, 1004 (1962).

[46] K. K. Thornber and R. P. Feynman, Phys. Rev. B **1**, 4099 (1970)

[47] E. Kartheuser, R. Evrard, and J. Devreese, Phys. Rev. Lett.**22**, 94 (1969).

[48] J. T. Devreese, J. De Sitter, and M. Goovaerts, Phys. Rev. B **5**, 2367 (1972)

[49] F. M. Peeters and J. T. Devreese *Phys. Rev. B* **34**, 7246 (1986).

[50] V. Cataudella, G. De Filippis, and G. Iadonisi, *Eur. Phys. J. B* **12**, 17 (1999).

[51] J. Tempere and J. T. Devreese, *Phys. Rev. B* **64**, 104504 (2001).

[52] J. Devreese, R. Evrard, and E. Kartheuser, *Phys. Rev. B* **12** 3353 (1975).

[53] W. Becker, B. Gerlach, H. Schliffke, Phys. Rev. B **28**, 5735 (1983).

[54] F. M. Peeters, J. T. Devreese, Phys. Rev. B **31**, 6826 (1985).

[55] Y. Osaka, Prog. Theor. Phys. **22**, 437 (1959).

[56] J. T. Devreese, *Contribution to the polaron theory*, Ph.D. Thesis, KU Leuven, 1964.

[57] J. T. Devreese and R. Evrard, in *Proceedings of the British Ceramic Society* **10**, 151 (1968). Reprinted in: *Path Integrals and Their Applications in Quantum, Statistical, and Solid State Physics*, edited by G. J. Papadopoulos and J. T. Devreese, NATO ASI Series B, Physics, vol. 34, Plenum, New York, 1977, pp. 344-357.

[58] F. M. Peeters and J. T. Devreese, in *Solid State Physics*, edited by F. Seitz and D. Turnbull, Academic Press, New York, 1984, vol. 38, pp. 81 - 133.

[59] H. Fröhlich, *Proc. R. Soc. (London) Ser. A* **160**, 230 (1937).

[60] D. J. Howarth and E. H. Sondheimer, *Proc. R. Soc. (London) Ser. A* **219**, 53 (1953).

[61] Y. Osaka, *Progr. Theoret. Phys.* **25**, 517 (1961).

[62] F. E. Low and D. Pines, *Phys. Rev.* **98**, 414 (1955).

[63] L. P. Kadanoff, *Phys. Rev.* **130**, 1364 (1963).

[64] D. C. Langreth and L. P. Kadanoff, *Phys. Rev.* **133**, A1070 (1964).

[65] F.M. Peeters, J.T. Devreese, *Phys. Stat. Sol. (b)* **115**, 539 (1983).

[66] R. W. Hellwarth and I. Biaggio, *Phys. Rev. B* **60**, 299 (1999).

[67] F. C. Brown, in *Point Defects in Solids*, edited by J. H. Crawford and L. M. Slifkin, Plenum, New York, 1972, vol. 1, p. 537.

[68] E. Hendry, F. Wang, J. Shan, T. F. Heinz, and M. Bonn, *Phys. Rev. B* **69**, 081101(R) (2004).

[69] V. L. Gurevich, I. G. Lang, and Yu. A. Firsov, *Fiz. Tverd. Tela* **4**, 1252 (1962) [English translation: *Sov. Phys. — Solid St.* **4**, 918 (1962)].

[70] G. D. Mahan, Many-Particle Physics, Kluwer/Plenum, New York, 2000.

[71] J. Devreese, W. Huybrechts, and L. Lemmens, Phys. Stat. Sol. (b) **48**, 77 (1971).

[72] H. Finkenrath, N. Uhle, and W. Waidelich, *Solid State Commun.* **7**, 11 (1969).

[73] M. J. Goovaerts, J. De Sitter, and J. T. Devreese, *Phys. Rev.* **7**, 2639 (1973).

[74] F. M. Peeters and J. T. Devreese, Phys. Rev. B **28**, 6051 (1983).

[75] D. Forster, *Hydrodynamic Fluctuations, Broken Symmetry and Correlation Functions,* Benjamin, New York, 1975.

[76] F. M. Peeters and J. T. Devreese, Phys. Rev. B 23, 1936 (1981).

[77] A. S. Mishchenko, N. Nagaosa, N. V. Prokof'ev, A. Sakamoto, and B. V. Svistunov, *Phys. Rev. Lett.* **91**, 236401 (2003).

[78] W. Huybrechts and J.T. Devreese, Phys. Rev. B **8**, 5754 (1973).

[79] D. M. Eagles, R. P. S. M. Lobo, and F. Gervais, *Phys. Rev. B* **52**, 6440 (1995).

[80] J. T. Devreese, L. Lemmens, and J. Van Royen, Phys. Rev. B **15**, 1212 (1977).

[81] L. F. Lemmens, J. De Sitter, and J. T. Devreese, Phys. Rev. B **8**, 2717 (1973).

[82] F. M. Peeters, Wu Xiaoguang, J. T. Devreese, Phys. Rev. B **33**, 3926 (1986).

[83] Wu Xiaoguang, F. M. Peeters, J. T. Devreese, Phys. Rev. B **31**, 3420 (1985).

[84] F. M. Peeters, J. T. Devreese, Phys. Rev. B **36**, 4442 (1987).

[85] F. M. Peeters, J. T. Devreese, Phys. Rev. B **28**, 6051 (1983).

[86] J. T. Devreese, L. F. Lemmens, J. Van Royen, Phys. Rev. B **2**, 1212 (1977).

[87] F. M. Peeters, J. T. Devreese, Solid State Phys. **38**, 81 (1984).

[88] F. Brosens, S. N. Klimin, and J. T. Devreese, Phys. Rev. B **77**, 085308 (2008).

[89] L. F. Lemmens, J. T. Devreese, and F. Brosens, Phys. Stat. Sol. (b) **82**, 439 (1977).

[90] S. Lupi, P. Maselli, M. Capizzi, P. Calvani, P. Giura and P. Roy, Phys. Rev. Lett. **83**, 4852 (1999).

[91] Ch. Hartinger, F. Mayr, J. Deisenhofer, A. Loidl, and T. Kopp, Phys. Rev. B **69**, 100403(R) (2004).

[92] Ch. Hartinger, F. Mayr, A. Loidl, and T. Kopp, cond-mat/0406123.

[93] D. Emin, Phys. Rev. B **48**, 13691 (1993).

[94] G. P. Zhang, T. A. Callcott, G. T. Woods, L. Lin, B. Sales, D. Mandrus, and J. He, Phys. Rev. Lett. **88**, 077401 (2002).

[95] T. Hotta, Phys. Rev. B **67**, 104428 (2003).

[96] S. N. Klimin, V. M. Fomin, F. Brosens, and J. T. Devreese, Phys. Rev. B **69**, 235324 (2004).

[97] L. F. Lemmens, F. Brosens, and J. T. Devreese, Phys. Rev. E **53**, 4467 (1996).

[98] J. T. Devreese, S. N. Klimin, V. M. Fomin, and F. Brosens, Solid State Communications **114**, 305 (2000).

[99] F. Brosens, J. T. Devreese, and L. F. Lemmens, Phys. Rev. E **55**, 227 (1997); **55**, 6795 (1997); **58**, 1634 (1998).

[100] L. F. Lemmens, F. Brosens, and J. T. Devreese, Solid State Communications **109**, 615 (1999).

[101] G. De Filippis, V. Cataudella and G. Iadonisi, Eur. Phys. J. B **8**, 339 (1999).

[102] W.B. da Costa, N. Studart, Phys. Rev. B **47**, 6356 (1993).

[103] F. Brosens, S. N. Klimin, and J. T. Devreese, Phys. Rev. B **71**, 214301 (2005).

[104] S. N. Klimin, V. M. Fomin, F. Brosens, and J. T. Devreese, Physica E **22**, 494 (2004).

[105] J. T. Devreese, in: *Fluctuating Paths and Fields* (World Scientific, Singapore, 2001), pp. 289-304.

[106] G. Verbist, F. M. Peeters and J. T. Devreese, Phys. Rev. B **43**, 2712 (1991).

[107] M. A. Smondyrev, G. Verbist, F. M. Peeters, and J. T. Devreese, Phys. Rev. B **47**, 2596 (1993).

[108] N. I. Kashirina, V. D. Lakhno, and V. V. Sychyov, Phys. Stat. Sol. (b) **239**, 174 (2003).

[109] J. Tempere, S. N. Klimin, I. F. Silvera, J. T. Devreese, Eur. Phys. J. **32**, 329 (2003).

[110] A.P. Volodin, M.S. Khaikin, and V.S. Edelman, JETP Lett. **26**, 543 (1977); U. Albrecht and P. Leiderer, Europhys. Lett. **3**, 705 (1987).

[111] V.B. Shikin, JETP Lett. **27**, 39 (1978); M.M. Salomaa and G.A. Williams, Phys. Rev. Lett. **47**, 1730 (1981).

[112] I.F. Silvera, Bull. Am. Phys. Soc. **46**, 1016 (2001).

[113] J. Tempere, I.F. Silvera, and J.T. Devreese, Phys. Rev. Lett. **87**, 275301 (2001).

[114] S. Fratini and P. Quémerais, Eur. Phys. J. B **14**, 99 (2000).

[115] F. Lindemann, Z. Phys. **11**, 609 (1910); C.M. Care and N.H. March, Adv. Phys. **24**, 101 (1975).

[116] C.C. Grimes and G. Adams, Phys. Rev. Lett. **42**, 795 (1979).

[117] V.M. Bedanov and F.M. Peeters, Phys. Rev. B **49**, 2667 (1994).

[118] L.P. Gor'kov and D.M. Chernikova, Pis'ma Zh. Eksp. Teor. Fiz. **18**, 119 (1973) [JETP Lett. **18**, 68 (1973)].

[119] J. Tempere, I.F. Silvera, and J.T. Devreese, Phys. Rev. B **67**, 035402 (2003).

[120] D.S. Fisher, B.I. Halperin, and P.M. Platzman, Phys. Rev. Lett. **42**, 798 (1979).

[121] G. Deville *et al.*, Phys. Rev. Lett. **53**, 588 (1984).